

# thm\_2EquantHeuristics\_2ESIMPLE\_GUESS\_EXISTS\_AND\_1 (TMaGHm82NCLivQ6VFeJawf9xsahBsKobwvc)

October 26, 2020

**Definition 1** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define  $c\_2Ebool\_2E\_2T$  to be  $(ap (ap (c\_2Emin\_2E\_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

**Definition 3** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A\_27a : \iota.(\lambda V0P \in (2^{A\_27a}).(ap (ap (c\_2Emin\_2E\_3D (2^{A\_27a}))$

**Definition 4** We define  $c\_2Ebool\_2E\_2F$  to be  $(ap (c\_2Ebool\_2E\_21 2) (\lambda V0t \in 2.V0t))$ .

**Definition 5** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o (p \Rightarrow q)$  of type  $\iota$ .

**Definition 6** We define  $c\_2Ebool\_2E\_27E$  to be  $(\lambda V0t \in 2.(ap (ap c\_2Emin\_2E\_3D\_3D\_3E V0t) c\_2Ebool\_2E\_2F$

**Definition 7** We define  $c\_2Ebool\_2E\_2F\_5C$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_2Ebool\_2E\_21 2) (\lambda V2t \in 2.V2t$

**Definition 8** We define  $c\_2EquantHeuristics\_2ESIMPLE\_GUESS\_EXISTS$  to be  $\lambda A\_27a : \iota.\lambda V0v \in A\_27a.\lambda V1i \in A\_27a.\lambda V2P \in 2.(ap (ap c\_2Emin\_2E\_3D\_3D\_3E V2P) (ap (ap (c\_2Emin\_2E\_3D$

Assume the following.

$$True \tag{1}$$

Assume the following.

$$\forall A\_27a.nonempty A\_27a \Rightarrow (\forall V0t \in 2.((\forall V1x \in A\_27a.(p V0t) \Leftrightarrow (p V0t)))) \tag{2}$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2.(((True \Rightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Rightarrow True) \Leftrightarrow \\ & True) \wedge (((False \Rightarrow (p V0t)) \Leftrightarrow True) \wedge (((p V0t) \Rightarrow (p V0t)) \Leftrightarrow True) \wedge (( \\ & (p V0t) \Rightarrow False) \Leftrightarrow (\neg (p V0t)))))) \end{aligned} \tag{3}$$

Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow (\forall V0x \in A\_27a. ((V0x = V0x) \Leftrightarrow True)) \quad (4)$$

Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow (\forall V0x \in A\_27a. (\forall V1y \in A\_27a. ((V0x = V1y) \Leftrightarrow (V1y = V0x)))) \quad (5)$$

Assume the following.

$$(\forall V0t \in 2. (((True \Leftrightarrow (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \Leftrightarrow True) \Leftrightarrow (p\ V0t)) \wedge (((False \Leftrightarrow (p\ V0t)) \Leftrightarrow \neg(p\ V0t)) \wedge (((p\ V0t) \Leftrightarrow False) \Leftrightarrow \neg(p\ V0t)))))) \quad (6)$$

Assume the following.

$$(\forall V0t1 \in 2. (\forall V1t2 \in 2. (\forall V2t3 \in 2. (((p\ V0t1) \Rightarrow ((p\ V1t2) \Rightarrow (p\ V2t3))) \Leftrightarrow (((p\ V0t1) \wedge (p\ V1t2)) \Rightarrow (p\ V2t3)))))) \quad (7)$$

Assume the following.

$$(\forall V0x \in 2. (\forall V1x\_27 \in 2. (\forall V2y \in 2. (\forall V3y\_27 \in 2. (((((p\ V0x) \Leftrightarrow (p\ V1x\_27)) \wedge ((p\ V1x\_27) \Rightarrow ((p\ V2y) \Leftrightarrow (p\ V3y\_27)))))) \Rightarrow (((p\ V0x) \Rightarrow (p\ V2y)) \Leftrightarrow ((p\ V1x\_27) \Rightarrow (p\ V3y\_27)))))) \quad (8)$$

**Theorem 1**

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow (\forall V0v \in A\_27a. (\forall V1i \in A\_27a. (\forall V2P1 \in 2. (\forall V3P2 \in 2. ((p\ (ap\ (ap\ (ap\ (c\_2EquantHeuristics\_2ESIMPLE\_GUESS\_EXISTS\ A\_27a)\ V0v)\ V1i)\ V2P1)) \Rightarrow (p\ (ap\ (ap\ (ap\ (c\_2EquantHeuristics\_2ESIMPLE\_GUESS\_EXISTS\ A\_27a)\ V0v)\ V1i)\ (ap\ (ap\ c\_2Ebool\_2E\_2F\_5C\ V2P1)\ V3P2))))))))$$