

thm_2EquantHeuristics_2ESIMPLE_GUESS_FORALL_IMP_2 (TMKVRv5XnqcA2MEfqgrDwsUz4Fa318QTeZV)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define $c_2Ebool_2E_21$ to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 3 We define $c_2Ebool_2E_21$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap (ap (c_2Emin_2E_3D (2^{A_27a}))$

Definition 4 We define $c_2Ebool_2E_21$ to be $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$.

Definition 5 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow p Q)$ of type ι .

Definition 6 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t))$

Definition 7 We define $c_2Ebool_2E_7E$ to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2E_2F_5C$

Definition 8 We define $c_2EquantHeuristics_2ESIMPLE_GUESS_FORALL$ to be $\lambda A_27a : \iota.\lambda V0v \in A_27a.\lambda V1i \in A_27a.\lambda V2P \in 2.(ap (ap c_2Emin_2E_3D_3D_3E (ap c_2Ebool_2E_7E V2P$

Assume the following.

$$True \tag{1}$$

Assume the following.

$$(\forall V0t1 \in 2.(\forall V1t2 \in 2.(((p V0t1) \Rightarrow (p V1t2)) \Rightarrow (((p V1t2) \Rightarrow (p V0t1)) \Rightarrow ((p V0t1) \Leftrightarrow (p V1t2)))))) \tag{2}$$

Assume the following.

$$(\forall V0t \in 2.(False \Rightarrow (p V0t))) \tag{3}$$

Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow (\forall V0t \in 2.((\forall V1x \in A_27a.(p V0t)) \Leftrightarrow (p V0t))) \tag{4}$$

Assume the following.

$$(\forall V0t \in 2.(((True \Rightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Rightarrow True) \Leftrightarrow True) \wedge (((False \Rightarrow (p V0t)) \Leftrightarrow True) \wedge (((p V0t) \Rightarrow (p V0t)) \Leftrightarrow True) \wedge ((p V0t) \Rightarrow False) \Leftrightarrow (\neg(p V0t)))))) \quad (5)$$

Assume the following.

$$((\forall V0t \in 2.((\neg(\neg(p V0t))) \Leftrightarrow (p V0t)) \wedge (((\neg True) \Leftrightarrow False) \wedge ((\neg False) \Leftrightarrow True))) \quad (6)$$

Assume the following.

$$\forall A.27a.nonempty \ A.27a \Rightarrow (\forall V0x \in A.27a.((V0x = V0x) \Leftrightarrow True)) \quad (7)$$

Assume the following.

$$\forall A.27a.nonempty \ A.27a \Rightarrow (\forall V0x \in A.27a.(\forall V1y \in A.27a.((V0x = V1y) \Leftrightarrow (V1y = V0x)))) \quad (8)$$

Assume the following.

$$(\forall V0A \in 2.(\forall V1B \in 2.((\neg((p V0A) \Rightarrow (p V1B))) \Leftrightarrow ((p V0A) \wedge (\neg(p V1B))))) \quad (9)$$

Assume the following.

$$(\forall V0t1 \in 2.(\forall V1t2 \in 2.(\forall V2t3 \in 2.(((p V0t1) \Rightarrow ((p V1t2) \Rightarrow (p V2t3))) \Leftrightarrow (((p V0t1) \wedge (p V1t2)) \Rightarrow (p V2t3)))))) \quad (10)$$

Assume the following.

$$(\forall V0x \in 2.(\forall V1x.27 \in 2.(\forall V2y \in 2.(\forall V3y.27 \in 2.(((p V0x) \Leftrightarrow (p V1x.27)) \wedge ((p V1x.27) \Rightarrow ((p V2y) \Leftrightarrow (p V3y.27)))))) \Rightarrow (((p V0x) \Rightarrow (p V2y)) \Leftrightarrow ((p V1x.27) \Rightarrow (p V3y.27)))))) \quad (11)$$

Theorem 1

$$\forall A.27a.nonempty \ A.27a \Rightarrow (\forall V0v \in A.27a.(\forall V1i \in A.27a.(\forall V2P1 \in 2.(\forall V3P2 \in 2.((p (ap (ap (ap (c.2EquantHeuristics.2ESIMPLE_GUESS_FORALL A.27a) V0v) V1i) V3P2)) \Rightarrow (p (ap (ap (ap (c.2EquantHeuristics.2ESIMPLE_GUESS_FORALL A.27a) V0v) V1i) (ap (ap c.2Emin.2E.3D.3D.3E V2P1) V3P2))))))))))$$