

thm\_2EquantHeuristics\_2ESIMPLE\_\_GUESS\_\_SELECT\_\_THM  
 (TMXBDjP-  
 swBN17pCKPzBMLuwW7ceXSn3X5J3)

October 26, 2020

**Definition 1** We define `c_2Emin_2E_3D` to be  $\lambda A. \lambda x \in A. \lambda y \in A. \text{inj\_o } (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define `c_2Ebool_2ET` to be  $(\text{ap } (\text{ap } (\text{c\_2Emin\_2E\_3D } (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

**Definition 3** We define `c_2Ebool_2E_21` to be  $\lambda A. 27a : \iota. (\lambda V0P \in (2^{A-27a}). (\text{ap } (\text{ap } (\text{c\_2Emin\_2E\_3D } (2^{A-27a}))))$

**Definition 4** We define `c_2Ebool_2EF` to be  $(\text{ap } (\text{c\_2Ebool\_2E\_21 } 2) (\lambda V0t \in 2.V0t))$ .

**Definition 5** We define `c_2Emin_2E_3D_3D_3E` to be  $\lambda P \in 2. \lambda Q \in 2. \text{inj\_o } (p P \Rightarrow p Q)$  of type  $\iota$ .

**Definition 6** We define `c_2Ebool_2E_2F_5C` to be  $(\lambda V0t1 \in 2. (\lambda V1t2 \in 2. (\text{ap } (\text{c\_2Ebool\_2E\_21 } 2) (\lambda V2t \in 2.V2t))))$

**Definition 7** We define `c_2Emin_2E_40` to be  $\lambda A. \lambda P \in 2^A. \text{if } (\exists x \in A. p (ap P x)) \text{ then } (the (\lambda x. x \in A \wedge p x))$  of type  $\iota \Rightarrow \iota$ .

**Definition 8** We define `c_2Ebool_2ECOND` to be  $\lambda A. 27a : \iota. (\lambda V0t \in 2. (\lambda V1t1 \in A. 27a. (\lambda V2t2 \in A. 27a. (ap P (c_2Emin_2E_40 A))))$

**Definition 9** We define `c_2Ebool_2E_3F` to be  $\lambda A. 27a : \iota. (\lambda V0P \in (2^{A-27a}). (\text{ap } V0P (\text{ap } (\text{c\_2Emin\_2E\_40 } A))))$

**Definition 10** We define `c_2EquantHeuristics_2ESIMPLE__GUESS__EXISTS` to be  $\lambda A. 27a : \iota. \lambda V0v \in A. 27a. \lambda V1i \in A. 27a. \lambda V2P \in 2. (\text{ap } (\text{ap } (\text{c\_2Emin\_2E\_3D\_3D\_3E } V2P) (\text{ap } (\text{ap } (\text{c\_2Emin\_2E\_3D\_3D\_3E } V2P))))$

**Definition 11** We define `c_2Ebool_2E_5C_2F` to be  $(\lambda V0t1 \in 2. (\lambda V1t2 \in 2. (\text{ap } (\text{c\_2Ebool\_2E\_21 } 2) (\lambda V2t \in 2.V2t))))$

**Definition 12** We define `c_2Ebool_2E_7E` to be  $(\lambda V0t \in 2. (\text{ap } (\text{ap } (\text{c\_2Emin\_2E\_3D\_3D\_3E } V0t) (\text{c\_2Ebool\_2E\_5C\_2F } V0t))))$

Assume the following.

$$\text{True} \tag{1}$$

Assume the following.

$$(\forall V0t1 \in 2. (\forall V1t2 \in 2. (((p V0t1) \Rightarrow (p V1t2)) \Rightarrow (((p V1t2) \Rightarrow (p V0t1)) \Rightarrow ((p V0t1) \Leftrightarrow (p V1t2)))))) \quad (2)$$

Assume the following.

$$(\forall V0t \in 2. (False \Rightarrow (p V0t))) \quad (3)$$

Assume the following.

$$(\forall V0t \in 2. ((p V0t) \vee (\neg(p V0t)))) \quad (4)$$

Assume the following.

$$(\forall V0t \in 2. (((True \Rightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Rightarrow True) \Leftrightarrow True) \wedge (((False \Rightarrow (p V0t)) \Leftrightarrow True) \wedge (((p V0t) \Rightarrow (p V0t)) \Leftrightarrow True) \wedge ((p V0t) \Rightarrow False) \Leftrightarrow (\neg(p V0t)))))) \quad (5)$$

Assume the following.

$$((\forall V0t \in 2. ((\neg(\neg(p V0t))) \Leftrightarrow (p V0t)) \wedge ((\neg True) \Leftrightarrow False) \wedge ((\neg False) \Leftrightarrow True))) \quad (6)$$

Assume the following.

$$\forall A\_27a.nonempty A\_27a \Rightarrow (\forall V0x \in A\_27a. ((V0x = V0x) \Leftrightarrow True)) \quad (7)$$

Assume the following.

$$\forall A\_27a.nonempty A\_27a \Rightarrow (\forall V0x \in A\_27a. (\forall V1y \in A\_27a. ((V0x = V1y) \Leftrightarrow (V1y = V0x)))) \quad (8)$$

Assume the following.

$$(\forall V0t \in 2. (((True \Leftrightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Leftrightarrow True) \Leftrightarrow (p V0t)) \wedge (((False \Leftrightarrow (p V0t)) \Leftrightarrow (\neg(p V0t))) \wedge (((p V0t) \Leftrightarrow False) \Leftrightarrow (\neg(p V0t)))))) \quad (9)$$

Assume the following.

$$\forall A\_27a.nonempty A\_27a \Rightarrow (\forall V0t1 \in A\_27a. (\forall V1t2 \in A\_27a. (((ap (ap (ap (c\_2Ebool\_2ECOND A\_27a) c\_2Ebool\_2ET) V0t1) V1t2) = V0t1) \wedge ((ap (ap (ap (c\_2Ebool\_2ECOND A\_27a) c\_2Ebool\_2EF) V0t1) V1t2) = V1t2)))))) \quad (10)$$

Assume the following.

$$\forall A\_27a.nonempty A\_27a \Rightarrow (\forall V0P \in (2^{A\_27a}). (\forall V1Q \in (2^{A\_27a}). (((\exists V2x \in A\_27a. (p (ap V0P V2x))) \wedge (\forall V3x \in A\_27a. ((p (ap V0P V3x)) \Rightarrow (p (ap V1Q V3x)))))) \Rightarrow (p (ap V1Q (ap (c\_2Emin\_2E\_40 A\_27a) V0P)))))) \quad (11)$$

Assume the following.

$$(\forall V0t1 \in 2.(\forall V1t2 \in 2.(\forall V2t3 \in 2.(((p V0t1) \Rightarrow ((p V1t2) \Rightarrow (p V2t3))) \Leftrightarrow (((p V0t1) \wedge (p V1t2)) \Rightarrow (p V2t3)))))) \quad (12)$$

Assume the following.

$$2.(((p V0x) \Leftrightarrow (p V1x_{.27})) \wedge ((p V1x_{.27}) \Rightarrow ((p V2y) \Leftrightarrow (p V3y_{.27})))) \Rightarrow ((p V0x) \Rightarrow (p V2y)) \Leftrightarrow ((p V1x_{.27}) \Rightarrow (p V3y_{.27})) \quad (13)$$

Assume the following.

$$(\forall V0t \in 2.((\neg(\neg(p V0t))) \Leftrightarrow (p V0t))) \quad (14)$$

Assume the following.

$$(\forall V0A \in 2.((p V0A) \Rightarrow ((\neg(p V0A)) \Rightarrow \text{False}))) \quad (15)$$

Assume the following.

$$(\forall V0A \in 2.(\forall V1B \in 2.(((\neg(\neg(p V0A) \vee (p V1B))) \Rightarrow \text{False}) \Leftrightarrow (((p V0A) \Rightarrow \text{False}) \Rightarrow ((\neg(p V1B)) \Rightarrow \text{False})))) \quad (16)$$

Assume the following.

$$(\forall V0A \in 2.(\forall V1B \in 2.(((\neg(\neg(p V0A) \vee (p V1B))) \Rightarrow \text{False}) \Leftrightarrow ((p V0A) \Rightarrow ((\neg(p V1B)) \Rightarrow \text{False})))) \quad (17)$$

Assume the following.

$$(\forall V0A \in 2.(((\neg(p V0A)) \Rightarrow \text{False}) \Rightarrow (((p V0A) \Rightarrow \text{False}) \Rightarrow \text{False}))) \quad (18)$$

Assume the following.

$$(\forall V0p \in 2.(\forall V1q \in 2.(\forall V2r \in 2.(((p V0p) \Leftrightarrow ((p V1q) \Leftrightarrow (p V2r))) \Leftrightarrow (((p V0p) \vee ((p V1q) \vee (p V2r))) \wedge (((p V0p) \vee ((\neg(p V2r)) \vee (\neg(p V1q)))) \wedge (((p V1q) \vee ((\neg(p V2r)) \vee (\neg(p V0p)))) \wedge ((p V2r) \vee ((\neg(p V1q)) \vee (\neg(p V0p)))))))))) \quad (19)$$

Assume the following.

$$(\forall V0p \in 2.(\forall V1q \in 2.(\forall V2r \in 2.(((p V0p) \Leftrightarrow ((p V1q) \vee (p V2r))) \Leftrightarrow (((p V0p) \vee (\neg(p V1q))) \wedge (((p V0p) \vee (\neg(p V2r))) \wedge ((p V1q) \vee ((p V2r) \vee (\neg(p V0p)))))))))) \quad (20)$$

Assume the following.

$$(\forall V0p \in 2.(\forall V1q \in 2.(\forall V2r \in 2.(((p V0p) \Leftrightarrow ((p V1q) \Rightarrow (p V2r))) \Leftrightarrow (((p V0p) \vee (p V1q)) \wedge (((p V0p) \vee (\neg(p V2r))) \wedge ((\neg(p V1q)) \vee ((p V2r) \vee (\neg(p V0p)))))))))) \quad (21)$$

Assume the following.

$$(\forall V0p \in 2. (\forall V1q \in 2. (((p V0p) \Leftrightarrow \neg(p V1q)) \Leftrightarrow (((p V0p) \vee (p V1q)) \wedge (\neg(p V1q) \vee \neg(p V0p))))) \quad (22)$$

Assume the following.

$$(\forall V0p \in 2. (\forall V1q \in 2. (\neg((p V0p) \Rightarrow (p V1q))) \Rightarrow (p V0p))) \quad (23)$$

Assume the following.

$$(\forall V0p \in 2. (\forall V1q \in 2. (\neg((p V0p) \Rightarrow (p V1q))) \Rightarrow \neg(p V1q))) \quad (24)$$

Assume the following.

$$(\forall V0p \in 2. (\forall V1q \in 2. (\neg((p V0p) \vee (p V1q))) \Rightarrow \neg(p V0p))) \quad (25)$$

Assume the following.

$$(\forall V0p \in 2. (\forall V1q \in 2. (\neg((p V0p) \vee (p V1q))) \Rightarrow \neg(p V1q))) \quad (26)$$

Assume the following.

$$(\forall V0p \in 2. (\neg(\neg(p V0p))) \Rightarrow (p V0p)) \quad (27)$$

**Theorem 1**

$$\forall A.27a.nonempty \ A.27a \Rightarrow (\forall V0i \in A.27a. (\forall V1P \in (2^{A.27a}). ((\forall V2v \in A.27a. (p (ap (ap (ap (c.2EquantHeuristics.2ESIMPLE\_GUESS\_EXISTS A.27a) V2v) V0i) (ap V1P V2v)))) \Rightarrow ((ap (c.2Emin.2E.40 A.27a) (\lambda V3v \in A.27a. (ap V1P V3v))) = (ap (ap (ap (c.2Ebool.2ECOND A.27a) (ap V1P V0i)) V0i) (ap (c.2Emin.2E.40 A.27a) (\lambda V4v \in A.27a.c.2Ebool.2EF))))))))$$