

# thm\_2EquantHeuristics\_2ESOME\_\_THE\_\_EQ (TMFS9MQMu2rvWn64rwcTjs9nUyDLw53NeVx)

October 26, 2020

**Definition 1** We define `c_2Emin_2E_3D_3D_3E` to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o (p P \Rightarrow p Q)$  of type  $\iota$ .

**Definition 2** We define `c_2Emin_2E_40` to be  $\lambda A.\lambda P \in 2^A.if (\exists x \in A.p (ap P x))$  then (the  $(\lambda x.x \in A \wedge p x)$ ) of type  $\iota \Rightarrow \iota$ .

**Definition 3** We define `c_2Emin_2E_3D` to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 4** We define `c_2Ebool_2E_3F` to be  $\lambda A.27a : \iota.(\lambda V0P \in (2^{A-27a}).(ap V0P (ap (c_2Emin_2E_40 A P))))$

**Definition 5** We define `c_2Ebool_2E_2T` to be  $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

**Definition 6** We define `c_2Ebool_2E_21` to be  $\lambda A.27a : \iota.(\lambda V0P \in (2^{A-27a}).(ap (ap (c_2Emin_2E_3D (2^{A-27a})) P)))$

**Definition 7** We define `c_2Ebool_2E_5C_2F` to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t)))$

Let `ty_2Eoption_2Eoption` :  $\iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A0.nonempty A0 \Rightarrow nonempty (ty_2Eoption_2Eoption A0) \quad (1)$$

**Definition 8** We define `c_2Ebool_2E_2F` to be  $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$ .

**Definition 9** We define `c_2Ebool_2E_7E` to be  $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2E_2F))$

Let `ty_2Eone_2Eone` :  $\iota$  be given. Assume the following.

$$nonempty ty_2Eone_2Eone \quad (2)$$

**Definition 10** We define `c_2Eone_2Eone` to be  $(ap (c_2Emin_2E_40 ty_2Eone_2Eone) (\lambda V0x \in ty_2Eone_2Eone))$

**Definition 11** We define `c_2Ebool_2E_2F_5C` to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t)))$

Let  $ty\_2Esum\_2Esum : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty\_2Esum\_2Esum\ A0\ A1) \quad (3)$$

Let  $c\_2Esum\_2EABS\_sum : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow \forall A\_27b.nonempty\ A\_27b \Rightarrow c\_2Esum\_2EABS\_sum\ A\_27a\ A\_27b \in ((ty\_2Esum\_2Esum\ A\_27a\ A\_27b)^{((2^{A\_27b})^{A\_27a})^2}) \quad (4)$$

**Definition 12** We define  $c\_2Esum\_2EINR$  to be  $\lambda A\_27a : \iota.\lambda A\_27b : \iota.\lambda V0e \in A\_27b.(ap\ (c\_2Esum\_2EABS\_sum\ A\_27a\ A\_27b)\ V0e)$

Let  $c\_2Eoption\_2Eoption\_ABS : \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow c\_2Eoption\_2Eoption\_ABS\ A\_27a \in ((ty\_2Eoption\_2Eoption\ A\_27a)^{(ty\_2Esum\_2Esum\ A\_27a\ ty\_2Eone\_2Eone)}) \quad (5)$$

**Definition 13** We define  $c\_2Eoption\_2ENONE$  to be  $\lambda A\_27a : \iota.(ap\ (c\_2Eoption\_2Eoption\_ABS\ A\_27a)\ 0)$

Let  $c\_2Eoption\_2EIS\_SOME : \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow c\_2Eoption\_2EIS\_SOME\ A\_27a \in (2^{(ty\_2Eoption\_2Eoption\ A\_27a)}) \quad (6)$$

**Definition 14** We define  $c\_2Esum\_2EINL$  to be  $\lambda A\_27a : \iota.\lambda A\_27b : \iota.\lambda V0e \in A\_27a.(ap\ (c\_2Esum\_2EABS\_sum\ A\_27a\ A\_27b)\ V0e)$

**Definition 15** We define  $c\_2Eoption\_2ESOME$  to be  $\lambda A\_27a : \iota.\lambda V0x \in A\_27a.(ap\ (c\_2Eoption\_2Eoption\_ABS\ A\_27a)\ V0x)$

Let  $c\_2Eoption\_2ETHE : \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow c\_2Eoption\_2ETHE\ A\_27a \in (A\_27a^{(ty\_2Eoption\_2Eoption\ A\_27a)}) \quad (7)$$

Assume the following.

$$True \quad (8)$$

Assume the following.

$$(\forall V0t1 \in 2.(\forall V1t2 \in 2.(((p\ V0t1) \Rightarrow (p\ V1t2)) \Rightarrow (((p\ V1t2) \Rightarrow (p\ V0t1)) \Rightarrow ((p\ V0t1) \Leftrightarrow (p\ V1t2)))))) \quad (9)$$

Assume the following.

$$(\forall V0t \in 2.(False \Rightarrow (p\ V0t))) \quad (10)$$

Assume the following.

$$((\forall V0t \in 2.((\neg(\neg(p\ V0t))) \Leftrightarrow (p\ V0t))) \wedge (((\neg True) \Leftrightarrow False) \wedge ((\neg False) \Leftrightarrow True))) \quad (11)$$

Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow (\forall V0x \in A\_27a. ((V0x = V0x) \Leftrightarrow True)) \quad (12)$$

Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow (\forall V0x \in A\_27a. (\forall V1y \in A\_27a. ((V0x = V1y) \Leftrightarrow (V1y = V0x)))) \quad (13)$$

Assume the following.

$$(\forall V0t \in 2. (((True \Leftrightarrow (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \Leftrightarrow True) \Leftrightarrow (p\ V0t)) \wedge (((False \Leftrightarrow (p\ V0t)) \Leftrightarrow \neg(p\ V0t)) \wedge (((p\ V0t) \Leftrightarrow False) \Leftrightarrow \neg(p\ V0t)))))) \quad (14)$$

Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow (\forall V0opt \in (ty\_2Eoption\_2Eoption\ A\_27a). ((V0opt = (c\_2Eoption\_2ENONE\ A\_27a)) \vee (\exists V1x \in A\_27a. (V0opt = (ap\ (c\_2Eoption\_2ESOME\ A\_27a)\ V1x)))))) \quad (15)$$

Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow (\forall V0x \in A\_27a. (\forall V1y \in A\_27a. (((ap\ (c\_2Eoption\_2ESOME\ A\_27a)\ V0x) = (ap\ (c\_2Eoption\_2ESOME\ A\_27a)\ V1y)) \Leftrightarrow (V0x = V1y)))) \quad (16)$$

Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow (\forall V0x \in A\_27a. (\neg((c\_2Eoption\_2ENONE\ A\_27a) = (ap\ (c\_2Eoption\_2ESOME\ A\_27a)\ V0x)))) \quad (17)$$

Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow ((\forall V0x \in A\_27a. ((p\ (ap\ (c\_2Eoption\_2EIS\_SOME\ A\_27a)\ (ap\ (c\_2Eoption\_2ESOME\ A\_27a)\ V0x))) \Leftrightarrow True)) \wedge ((p\ (ap\ (c\_2Eoption\_2EIS\_SOME\ A\_27a)\ (c\_2Eoption\_2ENONE\ A\_27a))) \Leftrightarrow False)) \quad (18)$$

Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow (\forall V0x \in A\_27a. ((ap\ (c\_2Eoption\_2ETHE\ A\_27a)\ (ap\ (c\_2Eoption\_2ESOME\ A\_27a)\ V0x)) = V0x)) \quad (19)$$

### Theorem 1

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow (\forall V0opt \in (ty\_2Eoption\_2Eoption\ A\_27a). (((ap\ (c\_2Eoption\_2ESOME\ A\_27a)\ (ap\ (c\_2Eoption\_2ETHE\ A\_27a)\ V0opt)) = V0opt) \Leftrightarrow (p\ (ap\ (c\_2Eoption\_2EIS\_SOME\ A\_27a)\ V0opt))))$$