

# thm\_2Equotient\_2ECOND\_PRS (TMWAu- FyMeZvmmmbPR6rHcpn98XSsDvASqsrJ)

October 26, 2020

**Definition 1** We define `c_2Emin_2E_3D` to be  $\lambda A. \lambda x \in A. \lambda y \in A. \text{inj\_o } (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define `c_2Ebool_2ET` to be  $(\text{ap } (\text{ap } (\text{c\_2Emin\_2E\_3D } (2^2))) (\lambda V0x \in 2. V0x)) (\lambda V1x \in 2. V1x)$

**Definition 3** We define `c_2Ebool_2E_21` to be  $\lambda A\_27a : \iota. (\lambda V0P \in (2^{A\_27a}). (\text{ap } (\text{ap } (\text{c\_2Emin\_2E\_3D } (2^{A\_27a}))))$

**Definition 4** We define `c_2Ebool_2EF` to be  $(\text{ap } (\text{c\_2Ebool\_2E\_21 } 2) (\lambda V0t \in 2. V0t))$ .

**Definition 5** We define `c_2Emin_2E_3D_3D_3E` to be  $\lambda P \in 2. \lambda Q \in 2. \text{inj\_o } (p \Rightarrow q)$  of type  $\iota$ .

**Definition 6** We define `c_2Ebool_2E_2F_5C` to be  $(\lambda V0t1 \in 2. (\lambda V1t2 \in 2. (\text{ap } (\text{c\_2Ebool\_2E\_21 } 2) (\lambda V2t \in 2. V2t))))$

**Definition 7** We define `c_2Emin_2E_40` to be  $\lambda A. \lambda P \in 2^A. \text{if } (\exists x \in A. p (\text{ap } P x)) \text{ then } (\text{the } (\lambda x. x \in A \wedge p (\text{ap } P x)))$  of type  $\iota \Rightarrow \iota$ .

**Definition 8** We define `c_2Ebool_2ECOND` to be  $\lambda A\_27a : \iota. (\lambda V0t \in 2. (\lambda V1t1 \in A\_27a. (\lambda V2t2 \in A\_27a. (\text{ap } (\text{c\_2Emin\_2E\_40 } A\_27a) (V2t2))))$

**Definition 9** We define `c_2Equotient_2EQUOTIENT` to be  $\lambda A\_27a : \iota. \lambda A\_27b : \iota. \lambda V0R \in ((2^{A\_27a})^{A\_27b}). \lambda V1R \in ((2^{A\_27b})^{A\_27a}). \text{inj\_o } ((\lambda V0x \in A\_27a. (\lambda V1y \in A\_27b. (\text{ap } (\text{c\_2Emin\_2E\_40 } A\_27a) (V1y)))) (\lambda V0x \in A\_27a. (\lambda V1y \in A\_27b. (\text{ap } (\text{c\_2Emin\_2E\_40 } A\_27b) (V0x))))))$

Assume the following.

$$\text{True} \tag{1}$$

Assume the following.

$$\forall A\_27a. \text{nonempty } A\_27a \Rightarrow (\forall V0x \in A\_27a. ((V0x = V0x) \Leftrightarrow \text{True})) \tag{2}$$

Assume the following.

$$\forall A\_27a. \text{nonempty } A\_27a \Rightarrow (\forall V0x \in A\_27a. (\forall V1y \in A\_27a. ((V0x = V1y) \Leftrightarrow (V1y = V0x)))) \tag{3}$$

Assume the following.

$$\begin{aligned}
& \forall A\_27a.nonempty\ A\_27a \Rightarrow \forall A\_27b.nonempty\ A\_27b \Rightarrow ( \\
& \quad \forall V0f \in (A\_27b^{A\_27a}). (\forall V1b \in 2. (\forall V2x \in A\_27a. \\
& \quad (\forall V3y \in A\_27a. ((ap\ V0f\ (ap\ (ap\ (ap\ (c\_2Ebool\_2ECOND\ A\_27a) \\
& \quad V1b)\ V2x)\ V3y)) = (ap\ (ap\ (ap\ (c\_2Ebool\_2ECOND\ A\_27b)\ V1b)\ (ap\ V0f \\
& \quad V2x))\ (ap\ V0f\ V3y))))))))) \\
\end{aligned} \tag{4}$$

Assume the following.

$$\begin{aligned}
& \forall A\_27a.nonempty\ A\_27a \Rightarrow \forall A\_27b.nonempty\ A\_27b \Rightarrow ( \\
& \quad \forall V0R \in ((2^{A\_27a})^{A\_27a}). (\forall V1abs \in (A\_27b^{A\_27a}). \\
& \quad (\forall V2rep \in (A\_27a^{A\_27b}). ((p\ (ap\ (ap\ (ap\ (c\_2Equotient\_2EQUOTIENT \\
& \quad A\_27a\ A\_27b)\ V0R)\ V1abs)\ V2rep)) \Rightarrow (\forall V3a \in A\_27b. ((ap\ V1abs \\
& \quad (ap\ V2rep\ V3a)) = V3a))))))))) \\
\end{aligned} \tag{5}$$

**Theorem 1**

$$\begin{aligned}
& \forall A\_27a.nonempty\ A\_27a \Rightarrow \forall A\_27b.nonempty\ A\_27b \Rightarrow ( \\
& \quad \forall V0R \in ((2^{A\_27a})^{A\_27a}). (\forall V1abs \in (A\_27b^{A\_27a}). \\
& \quad (\forall V2rep \in (A\_27a^{A\_27b}). ((p\ (ap\ (ap\ (ap\ (c\_2Equotient\_2EQUOTIENT \\
& \quad A\_27a\ A\_27b)\ V0R)\ V1abs)\ V2rep)) \Rightarrow (\forall V3a \in 2. (\forall V4b \in \\
& \quad A\_27b. (\forall V5c \in A\_27b. ((ap\ (ap\ (ap\ (c\_2Ebool\_2ECOND\ A\_27b) \\
& \quad V3a)\ V4b)\ V5c) = (ap\ V1abs\ (ap\ (ap\ (ap\ (c\_2Ebool\_2ECOND\ A\_27a)\ V3a) \\
& \quad (ap\ V2rep\ V4b))\ (ap\ V2rep\ V5c))))))))))))) \\
\end{aligned}$$