

thm_2Equotient_2EEQUALS__EQUIV__IMPLIES (TMbC7hhGBxnwEvpZuis7xkEkW9ktuuhQFGz)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define c_2Ebool_2ET to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 3 We define $c_2Ebool_2E_21$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap (ap (c_2Emin_2E_3D (2^{A_27a})$

Definition 4 We define $c_2Equotient_2EEQUIV$ to be $\lambda A_27a : \iota.(\lambda V0E \in ((2^{A_27a})^{A_27a}).(ap (c_2Ebool_2E_21$

Definition 5 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow p Q)$ of type ι .

Definition 6 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2$

Assume the following.

$$\begin{aligned}
 & \forall A_27a.nonempty A_27a \Rightarrow (\forall V0R \in ((2^{A_27a})^{A_27a}). \\
 & ((\forall V1x \in A_27a.(\forall V2y \in A_27a.((p (ap (ap V0R V1x) V2y)) \Leftrightarrow \\
 & ((ap V0R V1x) = (ap V0R V2y)))))) \Leftrightarrow ((\forall V3x \in A_27a.(p (ap (ap V0R \\
 & V3x) V3x))) \wedge ((\forall V4x \in A_27a.(\forall V5y \in A_27a.((p (ap (\\
 & ap V0R V4x) V5y)) \Rightarrow (p (ap (ap V0R V5y) V4x)))))) \wedge (\forall V6x \in A_27a. \\
 & (\forall V7y \in A_27a.(\forall V8z \in A_27a.(((p (ap (ap V0R V6x) V7y)) \wedge \\
 & (p (ap (ap V0R V7y) V8z))) \Rightarrow (p (ap (ap V0R V6x) V8z))))))))))
 \end{aligned} \tag{1}$$

Theorem 1

$$\begin{aligned}
 & \forall A_27a.nonempty A_27a \Rightarrow (\forall V0a1 \in A_27a.(\forall V1a2 \in \\
 & A_27a.(\forall V2b1 \in A_27a.(\forall V3b2 \in A_27a.(\forall V4R \in \\
 & ((2^{A_27a})^{A_27a}).((p (ap (c_2Equotient_2EEQUIV A_27a) V4R)) \Rightarrow \\
 & (((p (ap (ap V4R V0a1) V1a2)) \wedge (p (ap (ap V4R V2b1) V3b2))) \Rightarrow ((V0a1 = \\
 & V2b1) \Rightarrow (p (ap (ap V4R V1a2) V3b2))))))))))
 \end{aligned}$$