

thm_2Equotient_2EEQUIV__RES__FORALL
 (TM-
 MGfM5A53xoPvogJm7BFjg6LPQFGB2doYV)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define $c_2Ebool_2E_2T$ to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 3 We define $c_2Ebool_2E_21$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap (ap (c_2Emin_2E_3D (2^{A_27a}))$

Definition 4 We define $c_2Ebool_2E_2F$ to be $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$.

Definition 5 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow p Q)$ of type ι .

Definition 6 We define $c_2Ebool_2E_27E$ to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2E_2F$

Definition 7 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t))$

Definition 8 We define $c_2Equotient_2EEQUIV$ to be $\lambda A_27a : \iota.\lambda V0E \in ((2^{A_27a})^{A_27a}).(ap (c_2Ebool_2E_2F$

Definition 9 We define $c_2Ecombin_2EW$ to be $\lambda A_27a : \iota.\lambda A_27b : \iota.(\lambda V0f \in ((A_27b^{A_27a})^{A_27a}).(\lambda V1x \in 2.V1x$

Definition 10 We define $c_2Equotient_2Erespects$ to be $\lambda A_27a : \iota.\lambda A_27b : \iota.(c_2Ecombin_2EW A_27a A_27b$

Definition 11 We define $c_2Ebool_2E_2IN$ to be $\lambda A_27a : \iota.(\lambda V0x \in A_27a.(\lambda V1f \in (2^{A_27a}).(ap V1f V0x))$

Definition 12 We define $c_2Ebool_2ERES_FORALL$ to be $\lambda A_27a : \iota.(\lambda V0p \in (2^{A_27a}).(\lambda V1m \in (2^{A_27a}).$

Assume the following.

$$True \tag{1}$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2.(((True \Rightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Rightarrow True) \Leftrightarrow \\ & True) \wedge (((False \Rightarrow (p V0t)) \Leftrightarrow True) \wedge (((p V0t) \Rightarrow (p V0t)) \Leftrightarrow True) \wedge ((\\ & (p V0t) \Rightarrow False) \Leftrightarrow (\neg (p V0t)))))) \end{aligned} \tag{2}$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0x \in A_27a. ((V0x = V0x) \Leftrightarrow True)) \quad (3)$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0x \in A_27a. (\forall V1y \in A_27a. ((V0x = V1y) \Leftrightarrow (V1y = V0x)))) \quad (4)$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0P \in (2^{A_27a}). (\forall V1x \in A_27a. ((p\ (ap\ (ap\ (c_2Ebool_2EIN\ A_27a)\ V1x)\ V0P)) \Leftrightarrow (p\ (ap\ V0P\ V1x)))))) \quad (5)$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0R \in ((2^{A_27a})^{A_27a}). (\forall V1x \in A_27a. ((p\ (ap\ (ap\ (c_2Equotient_2Erespects\ A_27a\ 2)\ V0R)\ V1x)) \Leftrightarrow (p\ (ap\ (ap\ V0R\ V1x)\ V1x)))))) \quad (6)$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0P \in (2^{A_27a}). (\forall V1f \in (2^{A_27a}). ((p\ (ap\ (ap\ (c_2Ebool_2ERES_FORALL\ A_27a)\ V0P)\ V1f)) \Leftrightarrow (\forall V2x \in A_27a. ((p\ (ap\ (ap\ (c_2Ebool_2EIN\ A_27a)\ V2x)\ V0P)) \Rightarrow (p\ (ap\ V1f\ V2x)))))))))) \quad (7)$$

Theorem 1

$$\forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0E \in ((2^{A_27a})^{A_27a}). (\forall V1P \in (2^{A_27a}). ((p\ (ap\ (c_2Equotient_2EEQUIV\ A_27a)\ V0E)) \Rightarrow ((p\ (ap\ (ap\ (c_2Ebool_2ERES_FORALL\ A_27a)\ (ap\ (c_2Equotient_2Erespects\ A_27a\ 2)\ V0E))\ V1P)) \Leftrightarrow (p\ (ap\ (c_2Ebool_2E_21\ A_27a)\ V1P))))))))))$$