

thm_2Equotient_2EFORALL_REGULAR (TM- NCFXzX7YjHZjtd5KkF5dwyuCMndawaN6v)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow p Q)$ of type ι .

Definition 2 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 3 We define $c_2Ebool_2E_21$ to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 4 We define $c_2Ebool_2E_21$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap (ap (c_2Emin_2E_3D (2^{A_27a}))$

Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow (\forall V0x \in A_27a.(\forall V1y \in A_27a.((V0x = V1y) \Leftrightarrow (V1y = V0x)))) \quad (1)$$

Theorem 1

$$\forall A_27a.nonempty A_27a \Rightarrow (\forall V0P \in (2^{A_27a}).(\forall V1Q \in (2^{A_27a}).((\forall V2x \in A_27a.((p (ap V0P V2x)) \Rightarrow (p (ap V1Q V2x)))) \Rightarrow ((p (ap (c_2Ebool_2E_21 A_27a) V0P)) \Rightarrow (p (ap (c_2Ebool_2E_21 A_27a) V1Q)))))))$$