

thm_2Equotient_2EFUN__MAP__THM
(TMTvPn3R6xEofkZydV9htnuY3EQL9mU6rNS)

October 26, 2020

Definition 1 We define `c_2Emin_2E_3D` to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define `c_2Ebool_2ET` to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 3 We define `c_2Ebool_2E_21` to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap (ap (c_2Emin_2E_3D (2^{A_27a}))$

Definition 4 We define `c_2Equotient_2E_2D_2D_3E` to be $\lambda A_27a : \iota.\lambda A_27b : \iota.\lambda A_27c : \iota.\lambda A_27d : \iota.\lambda V0f$

Theorem 1

$\forall A_27a.nonempty A_27a \Rightarrow \forall A_27b.nonempty A_27b \Rightarrow \forall A_27c.nonempty A_27c \Rightarrow \forall A_27d.nonempty A_27d \Rightarrow (\forall V0f \in (A_27c^{A_27a}).$
 $(\forall V1g \in (A_27d^{A_27b}).(\forall V2h \in (A_27b^{A_27c}).(\forall V3x \in$
 $A_27a.((ap (ap (ap (ap (c_2Equotient_2E_2D_2D_3E A_27a A_27b A_27c$
 $A_27d) V0f) V1g) V2h) V3x) = (ap V1g (ap V2h (ap V0f V3x))))))))))$