

thm_2Equotient_2EIDENTITY__QUOTIENT (TMb4bjPcMqV8kKVGezAnqMH1wPBHHfgkUR2)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define $c_2Ebool_2E_ET$ to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 3 We define $c_2Ebool_2E_21$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap (ap (c_2Emin_2E_3D (2^{A_27a})$

Definition 4 We define $c_2Ebool_2E_EF$ to be $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$.

Definition 5 We define $c_2Ecombin_2E_EK$ to be $\lambda A_27a : \iota.\lambda A_27b : \iota.(\lambda V0x \in A_27a.(\lambda V1y \in A_27b.V0x))$

Definition 6 We define $c_2Ecombin_2E_ES$ to be $\lambda A_27a : \iota.\lambda A_27b : \iota.\lambda A_27c : \iota.(\lambda V0f \in ((A_27c^{A_27b})^{A_27a})$

Definition 7 We define $c_2Ecombin_2E_EI$ to be $\lambda A_27a : \iota.(ap (ap (c_2Ecombin_2E_ES A_27a (A_27a^{A_27a}) A_27a$

Definition 8 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow p Q)$ of type ι .

Definition 9 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t$

Definition 10 We define $c_2Equotient_2EQUOTIENT$ to be $\lambda A_27a : \iota.\lambda A_27b : \iota.\lambda V0R \in ((2^{A_27a})^{A_27b}).\lambda$

Assume the following.

$$True \tag{1}$$

Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow (\forall V0t \in 2.((\forall V1x \in A_27a.(p V0t) \Leftrightarrow (p V1x))) \tag{2}$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2.(((True \wedge (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \wedge True) \Leftrightarrow \\ & (p V0t)) \wedge (((False \wedge (p V0t)) \Leftrightarrow False) \wedge (((p V0t) \wedge False) \Leftrightarrow False) \wedge \\ & (((p V0t) \wedge (p V0t)) \Leftrightarrow (p V0t)))))) \end{aligned} \tag{3}$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0x \in A_27a. ((V0x = V0x) \Leftrightarrow True)) \quad (4)$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0x \in A_27a. ((ap\ (c_2Ecombin_2EI\ A_27a)\ V0x) = V0x)) \quad (5)$$

Theorem 1

$$\forall A_27a.nonempty\ A_27a \Rightarrow (p\ (ap\ (ap\ (ap\ (c_2Equotient_2EQUOTIENT\ A_27a\ A_27a)\ (c_2Emin_2E_3D\ A_27a))\ (c_2Ecombin_2EI\ A_27a))\ (c_2Ecombin_2EI\ A_27a)))$$