

thm_2Equotient_2ELAMBDA_PRS (TM- MvPUWr3uysFQGUH83kMtzrUVpxxAMMLbu)

October 26, 2020

Definition 1 We define `c_2Emin_2E_3D` to be $\lambda A. \lambda x \in A. \lambda y \in A. \text{inj_o } (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define `c_2Ebool_2ET` to be $(\text{ap } (\text{ap } (\text{c_2Emin_2E_3D } (2^2)) (\lambda V 0x \in 2. V 0x)) (\lambda V 1x \in 2. V 1x))$

Definition 3 We define `c_2Emin_2E_3D_3D_3E` to be $\lambda P \in 2. \lambda Q \in 2. \text{inj_o } (p P \Rightarrow p Q)$ of type ι .

Definition 4 We define `c_2Ebool_2E_21` to be $\lambda A. 27a : \iota. (\lambda V 0P \in (2^{A-27a}). (\text{ap } (\text{ap } (\text{c_2Emin_2E_3D } (2^{A-27a}))))$

Definition 5 We define `c_2Ebool_2E_2F_5C` to be $(\lambda V 0t1 \in 2. (\lambda V 1t2 \in 2. (\text{ap } (\text{c_2Ebool_2E_21 } 2) (\lambda V 2t \in 2. V 2t))))$

Definition 6 We define `c_2Equotient_2EQUOTIENT` to be $\lambda A. 27a : \iota. \lambda A. 27b : \iota. \lambda V 0R \in ((2^{A-27a})^{A-27a}). \lambda V 0f$

Definition 7 We define `c_2Equotient_2E_2D_2D_3E` to be $\lambda A. 27a : \iota. \lambda A. 27b : \iota. \lambda A. 27c : \iota. \lambda A. 27d : \iota. \lambda V 0f$

Assume the following.

$$\text{True} \tag{1}$$

Assume the following.

$$(\forall V 0t1 \in 2. (\forall V 1t2 \in 2. (((p V 0t1) \Rightarrow (p V 1t2)) \Rightarrow (((p V 1t2) \Rightarrow (p V 0t1)) \Rightarrow ((p V 0t1) \Leftrightarrow (p V 1t2)))))) \tag{2}$$

Assume the following.

$$\forall A. 27a. \text{nonempty } A. 27a \Rightarrow (\forall V 0x \in A. 27a. ((V 0x = V 0x) \Leftrightarrow \text{True})) \tag{3}$$

Assume the following.

$$\begin{aligned} & \forall A. 27a. \text{nonempty } A. 27a \Rightarrow \forall A. 27b. \text{nonempty } A. 27b \Rightarrow (\\ & \quad \forall V 0R \in ((2^{A-27a})^{A-27a}). (\forall V 1\text{abs} \in (A. 27b)^{A-27a}). \\ & (\forall V 2\text{rep} \in (A. 27a)^{A-27b}). ((p (\text{ap } (\text{ap } (\text{ap } (\text{c_2Equotient_2EQUOTIENT } \\ & \quad A. 27a \ A. 27b) \ V 0R) \ V 1\text{abs}) \ V 2\text{rep})) \Rightarrow (\forall V 3a \in A. 27b. ((\text{ap } \ V 1\text{abs} \\ & \quad (\text{ap } \ V 2\text{rep } \ V 3a)) = V 3a)))))) \end{aligned} \tag{4}$$

Assume the following.

$$\begin{aligned}
& \forall A_27a.nonempty\ A_27a \Rightarrow \forall A_27b.nonempty\ A_27b \Rightarrow \forall A_27c. \\
& nonempty\ A_27c \Rightarrow \forall A_27d.nonempty\ A_27d \Rightarrow (\forall V0f \in (A_27c^{A_27a}). \\
& (\forall V1g \in (A_27d^{A_27b}). (\forall V2h \in (A_27b^{A_27c}). (\forall V3x \in \\
& A_27a. ((ap\ (ap\ (ap\ (ap\ (c_2Equotient_2E_2D_2D_3E\ A_27a\ A_27b\ A_27c \\
& A_27d)\ V0f)\ V1g)\ V2h)\ V3x) = (ap\ V1g\ (ap\ V2h\ (ap\ V0f\ V3x)))))))))
\end{aligned} \tag{5}$$

Theorem 1

$$\begin{aligned}
& \forall A_27a.nonempty\ A_27a \Rightarrow \forall A_27b.nonempty\ A_27b \Rightarrow \forall A_27c. \\
& nonempty\ A_27c \Rightarrow \forall A_27d.nonempty\ A_27d \Rightarrow (\forall V0R1 \in (\\
& (2^{A_27a})^{A_27a}). (\forall V1abs1 \in (A_27c^{A_27a}). (\forall V2rep1 \in \\
& (A_27a^{A_27c}). ((p\ (ap\ (ap\ (ap\ (c_2Equotient_2EQUOTIENT\ A_27a\ A_27c) \\
& V0R1)\ V1abs1)\ V2rep1)) \Rightarrow (\forall V3R2 \in ((2^{A_27b})^{A_27b}). (\forall V4abs2 \in \\
& (A_27d^{A_27b}). (\forall V5rep2 \in (A_27b^{A_27d}). ((p\ (ap\ (ap\ (ap\ (c_2Equotient_2EQUOTIENT \\
& A_27b\ A_27d)\ V3R2)\ V4abs2)\ V5rep2)) \Rightarrow (\forall V6f \in (A_27d^{A_27c}). \\
& ((\lambda V7x \in A_27c. (ap\ V6f\ V7x)) = (ap\ (ap\ (ap\ (c_2Equotient_2E_2D_2D_3E \\
& A_27c\ A_27b\ A_27a\ A_27d)\ V2rep1)\ V4abs2)\ (\lambda V8x \in A_27a. (ap\ V5rep2 \\
& (ap\ V6f\ (ap\ V1abs1\ V8x)))))))))))))
\end{aligned}$$