

# thm\_2Quotient\_2LEFT\_RES\_FORALL\_REGULAR (TMLq3gwTJ6BY7U4ob7Bico1dUV9PWrR7mqn)

October 26, 2020

**Definition 1** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$   
of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define  $c\_2Ebool\_2ET$  to be  $(ap (ap (c\_2Emin\_2E\_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

**Definition 3** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A\_27a : \iota.(\lambda V0P \in (2^{A\_27a}).(ap (ap (c\_2Emin\_2E\_3D (2^{A\_27a}))$

**Definition 4** We define  $c\_2Ebool\_2EF$  to be  $(ap (c\_2Ebool\_2E\_21 2) (\lambda V0t \in 2.V0t))$ .

**Definition 5** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o (p P \Rightarrow p Q)$   
of type  $\iota$ .

**Definition 6** We define  $c\_2Ebool\_2E\_7E$  to be  $(\lambda V0t \in 2.(ap (ap c\_2Emin\_2E\_3D\_3D\_3E V0t) c\_2Ebool\_2EF$

**Definition 7** We define  $c\_2Ebool\_2E\_2F\_5C$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_2Ebool\_2E\_21 2) (\lambda V2t \in 2.V2t$

**Definition 8** We define  $c\_2Ebool\_2EIN$  to be  $\lambda A\_27a : \iota.(\lambda V0x \in A\_27a.(\lambda V1f \in (2^{A\_27a}).(ap V1f V0x)))$

**Definition 9** We define  $c\_2Ebool\_2ERES\_FORALL$  to be  $\lambda A\_27a : \iota.(\lambda V0p \in (2^{A\_27a}).(\lambda V1m \in (2^{A\_27a}).(a$

Assume the following.

$$True \tag{1}$$

Assume the following.

$$(\forall V0t1 \in 2.(\forall V1t2 \in 2.(((p V0t1) \Rightarrow (p V1t2)) \Rightarrow (((p V1t2) \Rightarrow (p V0t1)) \Rightarrow ((p V0t1) \Leftrightarrow (p V1t2)))))) \tag{2}$$

Assume the following.

$$\forall A\_27a.nonempty A\_27a \Rightarrow (\forall V0x \in A\_27a.(\forall V1y \in A\_27a.((V0x = V1y) \Leftrightarrow (V1y = V0x)))) \tag{3}$$

Assume the following.

$$(\forall V0t \in 2.(((True \Leftrightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Leftrightarrow True) \Leftrightarrow (p V0t)) \wedge (((False \Leftrightarrow (p V0t)) \Leftrightarrow \neg(p V0t)) \wedge (((p V0t) \Leftrightarrow False) \Leftrightarrow \neg(p V0t)))))) \quad (4)$$

Assume the following.

$$\forall A.27a.nonempty A.27a \Rightarrow (\forall V0P \in (2^{A.27a}).(\forall V1x \in A.27a.((p (ap (ap (c.2Ebool.2EIN A.27a) V1x) V0P)) \Leftrightarrow (p (ap V0P V1x)))))) \quad (5)$$

Assume the following.

$$\forall A.27a.nonempty A.27a \Rightarrow (\forall V0P \in (2^{A.27a}).(\forall V1f \in (2^{A.27a}).((p (ap (ap (c.2Ebool.2ERES\_FORALL A.27a) V0P) V1f)) \Leftrightarrow (\forall V2x \in A.27a.((p (ap (ap (c.2Ebool.2EIN A.27a) V2x) V0P)) \Rightarrow (p (ap V1f V2x)))))))))) \quad (6)$$

**Theorem 1**

$$\forall A.27a.nonempty A.27a \Rightarrow (\forall V0P \in (2^{A.27a}).(\forall V1R \in (2^{A.27a}).(\forall V2Q \in (2^{A.27a}).((\forall V3x \in A.27a.((p (ap V1R V3x)) \wedge ((p (ap V2Q V3x)) \Rightarrow (p (ap V0P V3x)))))) \Rightarrow ((p (ap (ap (c.2Ebool.2ERES\_FORALL A.27a) V1R) V2Q)) \Rightarrow (p (ap (c.2Ebool.2E.21 A.27a) V0P))))))))))$$