

# thm\_2Equotient\_2ENOT\_\_IMPLIES (TMdBaoczBWRpFMfjQXkZmzejgbkshwN9z5X)

October 26, 2020

**Definition 1** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define  $c\_2Ebool\_2E\_T$  to be  $(ap (ap (c\_2Emin\_2E\_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

**Definition 3** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A\_27a : \iota.(\lambda V0P \in (2^{A\_27a}).(ap (ap (c\_2Emin\_2E\_3D (2^{A\_27a}))$

**Definition 4** We define  $c\_2Ebool\_2E\_EF$  to be  $(ap (c\_2Ebool\_2E\_21 2) (\lambda V0t \in 2.V0t))$ .

**Definition 5** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o (p \Rightarrow q)$  of type  $\iota$ .

**Definition 6** We define  $c\_2Ebool\_2E\_7E$  to be  $(\lambda V0t \in 2.(ap (ap c\_2Emin\_2E\_3D\_3D\_3E V0t) c\_2Ebool\_2E\_EF$

Assume the following.

$$(\forall V0t \in 2.(False \Rightarrow (p V0t))) \tag{1}$$

Assume the following.

$$(\forall V0t \in 2.((\neg(p V0t)) \Rightarrow ((p V0t) \Rightarrow False))) \tag{2}$$

**Theorem 1**

$$(\forall V0P \in 2.(\forall V1Q \in 2.(((p V1Q) \Rightarrow (p V0P)) \Rightarrow ((\neg(p V0P)) \Rightarrow (\neg(p V1Q))))))$$