

thm_2Equotient_2EQUOTIENT_REL_ABS (TMRc5oLdvdurMwzWWZHFFbaukphC9VmwHyz)

October 26, 2020

Definition 1 We define `c_2Emin_2E_3D_3D_3E` to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow p Q)$ of type ι .

Definition 2 We define `c_2Emin_2E_3D` to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 3 We define `c_2Ebool_2E_2T` to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 4 We define `c_2Ebool_2E_21` to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap (ap (c_2Emin_2E_3D (2^{A_27a}))$

Definition 5 We define `c_2Ebool_2E_2F_5C` to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2$

Definition 6 We define `c_2Equotient_2EQUOTIENT` to be $\lambda A_27a : \iota.\lambda A_27b : \iota.\lambda V0R \in ((2^{A_27a})^{A_27a}).\lambda V$

Theorem 1

$$\begin{aligned} & \forall A_27a.nonempty A_27a \Rightarrow \forall A_27b.nonempty A_27b \Rightarrow (\\ & \quad \forall V0R \in ((2^{A_27a})^{A_27a}).(\forall V1abs \in (A_27b)^{A_27a}). \\ & (\forall V2rep \in (A_27a)^{A_27b}).((p (ap (ap (c_2Equotient_2EQUOTIENT \\ & \quad A_27a A_27b) V0R) V1abs) V2rep)) \Rightarrow (\forall V3r \in A_27a.(\forall V4s \in \\ & A_27a.((p (ap (ap V0R V3r) V4s)) \Rightarrow ((ap V1abs V3r) = (ap V1abs V4s)))))))) \end{aligned}$$