

thm_2Equotient_2ERES_EXISTS_REGULAR
(TMZ-
JADHe4m2gCkUNG9B7CQ2RCLzj9xsnGiE)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow p Q)$ of type ι .

Definition 2 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 3 We define $c_2Ebool_2E_2T$ to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 4 We define $c_2Ebool_2E_21$ to be $\lambda A.\lambda a : \iota.(\lambda V0P \in (2^{A-27a}).(ap (ap (c_2Emin_2E_3D (2^{A-27a}))$

Definition 5 We define $c_2Ebool_2E_2F$ to be $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$.

Definition 6 We define $c_2Ebool_2E_27E$ to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2E_2F$

Definition 7 We define $c_2Ebool_2E_2IN$ to be $\lambda A.\lambda a : \iota.(\lambda V0x \in A.\lambda a.(\lambda V1f \in (2^{A-27a}).(ap V1f V0x)))$

Definition 8 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t))$

Definition 9 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A.if (\exists x \in A.p (ap P x)) \mathbf{then} (the (\lambda x.x \in A \wedge p$ of type $\iota \Rightarrow \iota$.

Definition 10 We define $c_2Ebool_2E_3F$ to be $\lambda A.\lambda a : \iota.(\lambda V0P \in (2^{A-27a}).(ap V0P (ap (c_2Emin_2E_40$

Definition 11 We define $c_2Ebool_2ERES_EXISTS$ to be $\lambda A.\lambda a : \iota.(\lambda V0p \in (2^{A-27a}).(\lambda V1m \in (2^{A-27a}).($

Assume the following.

$$True \tag{1}$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2.(((True \wedge (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \wedge True) \Leftrightarrow \\ & (p V0t)) \wedge (((False \wedge (p V0t)) \Leftrightarrow False) \wedge (((p V0t) \wedge False) \Leftrightarrow False) \wedge \\ & (((p V0t) \wedge (p V0t)) \Leftrightarrow (p V0t)))))) \end{aligned} \tag{2}$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0x \in A_27a. (\forall V1y \in A_27a. ((V0x = V1y) \Leftrightarrow (V1y = V0x)))) \quad (3)$$

Assume the following.

$$(\forall V0t \in 2. (((True \Leftrightarrow (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \Leftrightarrow True) \Leftrightarrow (p\ V0t)) \wedge (((False \Leftrightarrow (p\ V0t)) \Leftrightarrow (\neg(p\ V0t))) \wedge (((p\ V0t) \Leftrightarrow False) \Leftrightarrow (\neg(p\ V0t))))) \quad (4)$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0P \in (2^{A_27a}). (\forall V1x \in A_27a. ((p\ (ap\ (ap\ (c_2Ebool_2EIN\ A_27a)\ V1x)\ V0P)) \Leftrightarrow (p\ (ap\ V0P\ V1x)))) \quad (5)$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0P \in (2^{A_27a}). (\forall V1f \in (2^{A_27a}). ((p\ (ap\ (ap\ (c_2Ebool_2ERES_EXISTS\ A_27a)\ V0P)\ V1f)) \Leftrightarrow (\exists V2x \in A_27a. ((p\ (ap\ (ap\ (c_2Ebool_2EIN\ A_27a)\ V2x)\ V0P)) \wedge (p\ (ap\ V1f\ V2x)))))) \quad (6)$$

Theorem 1

$$\forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0P \in (2^{A_27a}). (\forall V1Q \in (2^{A_27a}). (\forall V2R \in (2^{A_27a}). ((\forall V3x \in A_27a. ((p\ (ap\ V2R\ V3x)) \Rightarrow ((p\ (ap\ V0P\ V3x)) \Rightarrow (p\ (ap\ V1Q\ V3x)))))) \Rightarrow ((p\ (ap\ (ap\ (c_2Ebool_2ERES_EXISTS\ A_27a)\ V2R)\ V0P)) \Rightarrow (p\ (ap\ (ap\ (c_2Ebool_2ERES_EXISTS\ A_27a)\ V2R)\ V1Q))))))$$