

thm_2Equotient_list_2EREVERSE_PRS (TMKuDG1ERp2fyQNtkWyatw1iBQvBfgK8Krv)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define c_2Ebool_2ET to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Let $ty_2Elist_2Elist : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty A0 \Rightarrow nonempty (ty_2Elist_2Elist A0) \quad (1)$$

Let $c_2Elist_2EMAP : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow \forall A_27b.nonempty A_27b \Rightarrow c_2Elist_2EMAP A_27a A_27b \in (((ty_2Elist_2Elist A_27b)^{(ty_2Elist_2Elist A_27a)})^{(A_27b^{A_27a})}) \quad (2)$$

Let $c_2Elist_2EAPPEND : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow c_2Elist_2EAPPEND A_27a \in (((ty_2Elist_2Elist A_27a)^{(ty_2Elist_2Elist A_27a)})^{(ty_2Elist_2Elist A_27a)}) \quad (3)$$

Let $c_2Elist_2ECONS : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow c_2Elist_2ECONS A_27a \in (((ty_2Elist_2Elist A_27a)^{(ty_2Elist_2Elist A_27a)})^{A_27a}) \quad (4)$$

Let $c_2Elist_2ENIL : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow c_2Elist_2ENIL A_27a \in (ty_2Elist_2Elist A_27a) \quad (5)$$

Let $c_2Elist_2EREVERSE : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow c_2Elist_2EREVERSE A_27a \in ((ty_2Elist_2Elist A_27a)^{(ty_2Elist_2Elist A_27a)}) \quad (6)$$

Definition 3 We define `c_2Emin_2E_3D_3D_3E` to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow p Q)$ of type ι .

Definition 4 We define `c_2Ebool_2E_21` to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap (ap (c_2Emin_2E_3D (2^{A_27a})))$

Definition 5 We define `c_2Ebool_2E_2F_5C` to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2$

Definition 6 We define `c_2Equotient_2EQUOTIENT` to be $\lambda A_27a : \iota.\lambda A_27b : \iota.\lambda V0R \in ((2^{A_27a})^{A_27a}).\lambda V$

Assume the following.

$$True \tag{7}$$

Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow (\forall V0x \in A_27a.((V0x = V0x) \Leftrightarrow True)) \tag{8}$$

Assume the following.

$$\begin{aligned} & \forall A_27a.nonempty A_27a \Rightarrow \forall A_27b.nonempty A_27b \Rightarrow (\\ & (\forall V0f \in (A_27b^{A_27a}).((ap (ap (c_2Elist_2EMAP A_27a A_27b) \\ & V0f) (c_2Elist_2ENIL A_27a)) = (c_2Elist_2ENIL A_27b))) \wedge (\forall V1f \in \\ & (A_27b^{A_27a}).(\forall V2h \in A_27a.(\forall V3t \in (ty_2Elist_2Elist \\ & A_27a).((ap (ap (c_2Elist_2EMAP A_27a A_27b) V1f) (ap (ap (c_2Elist_2ECONS \\ & A_27a) V2h) V3t)) = (ap (ap (c_2Elist_2ECONS A_27b) (ap V1f V2h)) \\ & (ap (ap (c_2Elist_2EMAP A_27a A_27b) V1f) V3t))))))) \end{aligned} \tag{9}$$

Assume the following.

$$\begin{aligned} & \forall A_27a.nonempty A_27a \Rightarrow (\forall V0P \in (2^{(ty_2Elist_2Elist A_27a)}). \\ & (((p (ap V0P (c_2Elist_2ENIL A_27a))) \wedge (\forall V1t \in (ty_2Elist_2Elist \\ & A_27a).((p (ap V0P V1t)) \Rightarrow (\forall V2h \in A_27a.(p (ap V0P (ap (ap (\\ & c_2Elist_2ECONS A_27a) V2h) V1t))))))) \Rightarrow (\forall V3l \in (ty_2Elist_2Elist \\ & A_27a).(p (ap V0P V3l)))))) \end{aligned} \tag{10}$$

Assume the following.

$$\begin{aligned} & \forall A_27a.nonempty A_27a \Rightarrow \forall A_27b.nonempty A_27b \Rightarrow (\\ & \forall V0f \in (A_27b^{A_27a}).(\forall V1l1 \in (ty_2Elist_2Elist A_27a). \\ & (\forall V2l2 \in (ty_2Elist_2Elist A_27a).((ap (ap (c_2Elist_2EMAP \\ & A_27a A_27b) V0f) (ap (ap (c_2Elist_2EAPPEND A_27a) V1l1) V2l2)) = \\ & (ap (ap (c_2Elist_2EAPPEND A_27b) (ap (ap (c_2Elist_2EMAP A_27a \\ & A_27b) V0f) V1l1)) (ap (ap (c_2Elist_2EMAP A_27a A_27b) V0f) V2l2)))))) \end{aligned} \tag{11}$$

Assume the following.

$$\begin{aligned}
& \forall A.27a.nonempty\ A.27a \Rightarrow (((ap\ (c.2Elist.2EREVERSE\ A.27a) \\
& \quad (c.2Elist.2ENIL\ A.27a)) = (c.2Elist.2ENIL\ A.27a)) \wedge (\forall V0h \in \\
& \quad A.27a.(\forall V1t \in (ty.2Elist.2Elist\ A.27a).((ap\ (c.2Elist.2EREVERSE \\
& \quad A.27a)\ (ap\ (ap\ (c.2Elist.2ECONS\ A.27a)\ V0h)\ V1t)) = (ap\ (ap\ (c.2Elist.2EAPPEND \\
& \quad A.27a)\ (ap\ (c.2Elist.2EREVERSE\ A.27a)\ V1t))\ (ap\ (ap\ (c.2Elist.2ECONS \\
& \quad A.27a)\ V0h)\ (c.2Elist.2ENIL\ A.27a))))))
\end{aligned} \tag{12}$$

Assume the following.

$$\begin{aligned}
& \forall A.27a.nonempty\ A.27a \Rightarrow \forall A.27b.nonempty\ A.27b \Rightarrow (\\
& \quad \forall V0R \in ((2^{A.27a})^{A.27a}).(\forall V1abs \in (A.27b^{A.27a}). \\
& \quad (\forall V2rep \in (A.27a^{A.27b}).((p\ (ap\ (ap\ (ap\ (c.2Equotient.2EQUOTIENT \\
& \quad A.27a\ A.27b)\ V0R)\ V1abs)\ V2rep)) \Rightarrow (\forall V3a \in A.27b.((ap\ V1abs \\
& \quad (ap\ V2rep\ V3a)) = V3a))))))
\end{aligned} \tag{13}$$

Theorem 1

$$\begin{aligned}
& \forall A.27a.nonempty\ A.27a \Rightarrow \forall A.27b.nonempty\ A.27b \Rightarrow (\\
& \quad \forall V0R \in ((2^{A.27a})^{A.27a}).(\forall V1abs \in (A.27b^{A.27a}). \\
& \quad (\forall V2rep \in (A.27a^{A.27b}).((p\ (ap\ (ap\ (ap\ (c.2Equotient.2EQUOTIENT \\
& \quad A.27a\ A.27b)\ V0R)\ V1abs)\ V2rep)) \Rightarrow (\forall V3l \in (ty.2Elist.2Elist \\
& \quad A.27b).((ap\ (c.2Elist.2EREVERSE\ A.27b)\ V3l) = (ap\ (ap\ (c.2Elist.2EMAP \\
& \quad A.27a\ A.27b)\ V1abs)\ (ap\ (c.2Elist.2EREVERSE\ A.27a)\ (ap\ (ap\ (c.2Elist.2EMAP \\
& \quad A.27b\ A.27a)\ V2rep)\ V3l))))))
\end{aligned}$$