

# thm\_2Equotient\_\_option\_2ENONE\_\_PRS (TMPxCNsvhhaGa9S2btE3dSmesRZE2PV83e3)

October 26, 2020

**Definition 1** We define `c_2Emin_2E_3D` to be  $\lambda A. \lambda x \in A. \lambda y \in A. \text{inj\_o } (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define `c_2Emin_2E_3D_3D_3E` to be  $\lambda P \in 2. \lambda Q \in 2. \text{inj\_o } (p P \Rightarrow p Q)$  of type  $\iota$ .

**Definition 3** We define `c_2Ebool_2E_2T` to be  $(\text{ap } (\text{ap } (\text{c_2Emin_2E_3D } (2^2)) (\lambda V0x \in 2. V0x)) (\lambda V1x \in 2. V1x))$

**Definition 4** We define `c_2Ebool_2E_21` to be  $\lambda A_27a : \iota. (\lambda V0P \in (2^{A_27a}). (\text{ap } (\text{ap } (\text{c_2Emin_2E_3D } (2^{A_27a}))))$

**Definition 5** We define `c_2Ebool_2E_2F_5C` to be  $(\lambda V0t1 \in 2. (\lambda V1t2 \in 2. (\text{ap } (\text{c_2Ebool_2E_21 } 2)) (\lambda V2t \in 2. V2t)))$

**Definition 6** We define `c_2Equotient_2EQUOTIENT` to be  $\lambda A_27a : \iota. \lambda A_27b : \iota. \lambda V0R \in ((2^{A_27a})^{A_27a}). \lambda V1R \in ((2^{A_27b})^{A_27b}). \text{inj\_o } (V0R = V1R)$

Let `ty_2Eone_2Eone` :  $\iota$  be given. Assume the following.

$$\text{nonempty } \text{ty\_2Eone\_2Eone} \tag{1}$$

**Definition 7** We define `c_2Emin_2E_40` to be  $\lambda A. \lambda P \in 2^A. \text{if } (\exists x \in A. p (\text{ap } P x)) \text{ then } (\text{the } (\lambda x. x \in A \wedge p (\text{ap } P x)))$  of type  $\iota \Rightarrow \iota$ .

**Definition 8** We define `c_2Eone_2Eone` to be  $(\text{ap } (\text{c_2Emin_2E_40 } \text{ty\_2Eone\_2Eone}) (\lambda V0x \in \text{ty\_2Eone\_2Eone}. V0x))$

**Definition 9** We define `c_2Ebool_2E_2F` to be  $(\text{ap } (\text{c_2Ebool_2E_21 } 2)) (\lambda V0t \in 2. V0t)$ .

**Definition 10** We define `c_2Ebool_2E_7E` to be  $(\lambda V0t \in 2. (\text{ap } (\text{ap } (\text{c_2Emin_2E_3D_3D_3E } V0t)) (\text{c_2Ebool_2E_2F } 2)))$

Let `ty_2Esum_2Esum` :  $\iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A0. \text{nonempty } A0 \Rightarrow \forall A1. \text{nonempty } A1 \Rightarrow \text{nonempty } (\text{ty\_2Esum\_2Esum } A0 A1) \tag{2}$$

Let `c_2Esum_2EABS__sum` :  $\iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A_27a. \text{nonempty } A_27a \Rightarrow \forall A_27b. \text{nonempty } A_27b \Rightarrow \text{c\_2Esum\_2EABS\_sum } A_27a A_27b \in ((\text{ty\_2Esum\_2Esum } A_27a A_27b)^{((2^{A_27b})^{A_27a})^2}) \tag{3}$$

**Definition 11** We define  $c\_Esum\_2EINR$  to be  $\lambda A\_27a : \iota. \lambda A\_27b : \iota. \lambda V0e \in A\_27b. (ap (c\_Esum\_2EABS$   
Let  $ty\_2Eoption\_2Eoption : \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A0.nonempty A0 \Rightarrow nonempty (ty\_2Eoption\_2Eoption A0) \quad (4)$$

Let  $c\_2Eoption\_2Eoption\_ABS : \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A\_27a.nonempty A\_27a \Rightarrow c\_2Eoption\_2Eoption\_ABS A\_27a \in ((ty\_2Eoption\_2Eoption A\_27a)^{(ty\_2Esum\_2Esum A\_27a ty\_2Eone\_2Eone)}) \quad (5)$$

**Definition 12** We define  $c\_2Eoption\_2ENONE$  to be  $\lambda A\_27a : \iota. (ap (c\_2Eoption\_2Eoption\_ABS A\_27a) ($

**Definition 13** We define  $c\_2Esum\_2EINL$  to be  $\lambda A\_27a : \iota. \lambda A\_27b : \iota. \lambda V0e \in A\_27a. (ap (c\_2Esum\_2EABS$

**Definition 14** We define  $c\_2Eoption\_2ESOME$  to be  $\lambda A\_27a : \iota. \lambda V0x \in A\_27a. (ap (c\_2Eoption\_2Eoption\_MAP$

Let  $c\_2Eoption\_2EOPTION\_MAP : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A\_27a.nonempty A\_27a \Rightarrow \forall A\_27b.nonempty A\_27b \Rightarrow c\_2Eoption\_2EOPTION\_MAP A\_27a A\_27b \in (((ty\_2Eoption\_2Eoption A\_27b)^{(ty\_2Eoption\_2Eoption A\_27a)})^{(A\_27b^{A\_27a})}) \quad (6)$$

Assume the following.

$$True \quad (7)$$

Assume the following.

$$\forall A\_27a.nonempty A\_27a \Rightarrow (\forall V0x \in A\_27a. ((V0x = V0x) \Leftrightarrow True)) \quad (8)$$

Assume the following.

$$\forall A\_27a.nonempty A\_27a \Rightarrow \forall A\_27b.nonempty A\_27b \Rightarrow ( (\forall V0f \in (A\_27b^{A\_27a}). (\forall V1x \in A\_27a. ((ap (ap (c\_2Eoption\_2EOPTION\_MAP A\_27a A\_27b) V0f) (ap (c\_2Eoption\_2ESOME A\_27a) V1x)) = (ap (c\_2Eoption\_2ESOME A\_27b) (ap V0f V1x)))))) \wedge (\forall V2f \in (A\_27b^{A\_27a}). ((ap (ap (c\_2Eoption\_2EOPTION\_MAP A\_27a A\_27b) V2f) (c\_2Eoption\_2ENONE A\_27a)) = (c\_2Eoption\_2ENONE A\_27b)))))) \quad (9)$$

**Theorem 1**

$$\forall A\_27a.nonempty A\_27a \Rightarrow \forall A\_27b.nonempty A\_27b \Rightarrow ( (\forall V0R \in ((2^{A\_27a})^{A\_27a}). (\forall V1abs \in (A\_27b^{A\_27a}). (\forall V2rep \in (A\_27a^{A\_27b}). ((p (ap (ap (ap (c\_2Equotient\_2EQUOTIENT A\_27a A\_27b) V0R) V1abs) V2rep))) \Rightarrow ((c\_2Eoption\_2ENONE A\_27b) = (ap (ap (c\_2Eoption\_2EOPTION\_MAP A\_27a A\_27b) V1abs) (c\_2Eoption\_2ENONE A\_27a))))))))))$$