

thm_2Erat_2ERAT (TM- caB6FB2YpePFTd3AYaMQBn9MPQ7KS2Bbu)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow p Q)$ of type ι .

Definition 2 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 3 We define $c_2Ebool_2E_2T$ to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 4 We define $c_2Ebool_2E_21$ to be $\lambda A.\lambda a : \iota.(\lambda V0P \in (2^{A-27a}).(ap (ap (c_2Emin_2E_3D (2^{A-27a}))$

Definition 5 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t)))$

Let $ty_2Efrac_2Efrac : \iota$ be given. Assume the following.

$$nonempty\ ty_2Efrac_2Efrac \tag{1}$$

Let $ty_2Erat_2Erat : \iota$ be given. Assume the following.

$$nonempty\ ty_2Erat_2Erat \tag{2}$$

Let $c_2Erat_2Erep_rat_CLASS : \iota$ be given. Assume the following.

$$c_2Erat_2Erep_rat_CLASS \in ((2^{ty_2Efrac_2Efrac})^{ty_2Erat_2Erat}) \tag{3}$$

Definition 6 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A.if (\exists x \in A.p (ap P x)) \text{ then } (the (\lambda x.x \in A \wedge p x))$ of type $\iota \Rightarrow \iota$.

Definition 7 We define $c_2Erat_2Erep_rat$ to be $\lambda V0a \in ty_2Erat_2Erat.(ap (c_2Emin_2E_40 ty_2Efrac_2Efrac 2^a))$

Let $ty_2Einteger_2Eint : \iota$ be given. Assume the following.

$$nonempty\ ty_2Einteger_2Eint \tag{4}$$

Let $ty_2Epair_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty_2Epair_2Eprod\ A0\ A1) \tag{5}$$

Let $c_2Efrac_2Erep_frac : \iota$ be given. Assume the following.

$$c_2Efrac_2Erep_frac \in ((ty_2Epair_2Eprod\ ty_2Einteger_2Eint\ ty_2Einteger_2Eint)^{ty_2Efrac_2Efrac}) \quad (6)$$

Let $c_2Epair_2ESND : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow \forall A_27b.nonempty\ A_27b \Rightarrow c_2Epair_2ESND\ A_27a\ A_27b \in (A_27b)^{(ty_2Epair_2Eprod\ A_27a\ A_27b)} \quad (7)$$

Definition 8 We define $c_2Efrac_2Efrac_dnm$ to be $\lambda V0f \in ty_2Efrac_2Efrac.(ap\ (c_2Epair_2ESND\ ty_2Efrac_2Efrac))$

Let $c_2Epair_2EFST : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow \forall A_27b.nonempty\ A_27b \Rightarrow c_2Epair_2EFST\ A_27a\ A_27b \in (A_27a)^{(ty_2Epair_2Eprod\ A_27a\ A_27b)} \quad (8)$$

Definition 9 We define $c_2Efrac_2Efrac_nmr$ to be $\lambda V0f \in ty_2Efrac_2Efrac.(ap\ (c_2Epair_2EFST\ ty_2Efrac_2Efrac))$

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \quad (9)$$

Let $c_2Einteger_2Eint_REP_CLASS : \iota$ be given. Assume the following.

$$c_2Einteger_2Eint_REP_CLASS \in ((2^{(ty_2Epair_2Eprod\ ty_2Enum_2Enum\ ty_2Enum_2Enum)})^{ty_2Einteger_2Eint}) \quad (10)$$

Definition 10 We define $c_2Einteger_2Eint_REP$ to be $\lambda V0a \in ty_2Einteger_2Eint.(ap\ (c_2Emin_2E.40\ ty_2Einteger_2Eint_REP_CLASS\ a))$

Let $c_2Einteger_2Etint_mul : \iota$ be given. Assume the following.

$$c_2Einteger_2Etint_mul \in (((ty_2Epair_2Eprod\ ty_2Enum_2Enum\ ty_2Enum_2Enum)^{(ty_2Epair_2Eprod\ ty_2Enum_2Enum\ ty_2Enum_2Enum)})^{(ty_2Epair_2Eprod\ ty_2Enum_2Enum\ ty_2Enum_2Enum)}) \quad (11)$$

Let $c_2Einteger_2Etint_eq : \iota$ be given. Assume the following.

$$c_2Einteger_2Etint_eq \in ((2^{(ty_2Epair_2Eprod\ ty_2Enum_2Enum\ ty_2Enum_2Enum)})^{(ty_2Epair_2Eprod\ ty_2Enum_2Enum\ ty_2Enum_2Enum)}) \quad (12)$$

Let $c_2Einteger_2Eint_ABS_CLASS : \iota$ be given. Assume the following.

$$c_2Einteger_2Eint_ABS_CLASS \in (ty_2Einteger_2Eint)^{(2^{(ty_2Epair_2Eprod\ ty_2Enum_2Enum\ ty_2Enum_2Enum)})} \quad (13)$$

Definition 11 We define $c_2Einteger_2Eint_ABS$ to be $\lambda V0r \in (ty_2Epair_2Eprod\ ty_2Enum_2Enum\ ty_2Enum_2Enum).(c_2Einteger_2Eint_ABS_CLASS\ r)$

Definition 12 We define $c_2Einteger_2Eint_mul$ to be $\lambda V0T1 \in ty_2Einteger_2Eint.\lambda V1T2 \in ty_2Einteger_2Eint.(c_2Einteger_2Etint_mul\ T1\ T2)$

Definition 13 We define $c_2Erat_2Erat_equiv$ to be $\lambda V0f1 \in ty_2Efrac_2Efrac.\lambda V1f2 \in ty_2Efrac_2Efrac.(c_2Efrac_2Efrac_dnm\ f1\ f2)$

Let $c_2Erat_2Eabs_rat_CLASS : \iota$ be given. Assume the following.

$$c_2Erat_2Eabs_rat_CLASS \in (ty_2Erat_2Erat^{(2^{ty_2Efrac_2Efrac})})} \quad (14)$$

Definition 14 We define $c_2Erat_2Eabs_rat$ to be $\lambda V0r \in ty_2Efrac_2Efrac.(ap\ c_2Erat_2Eabs_rat_CLASS\ V0r)$

Definition 15 We define $c_2Equotient_2EQUOTIENT$ to be $\lambda A_27a : \iota.\lambda A_27b : \iota.\lambda V0R \in ((2^{A_27a})^{A_27a}).\lambda V0r \in ty_2Efrac_2Efrac.(ap\ c_2Equotient_2EQUOTIENT\ A_27a\ A_27b\ V0r)$

Assume the following.

$$(p\ (ap\ (ap\ (ap\ (c_2Equotient_2EQUOTIENT\ ty_2Efrac_2Efrac\ ty_2Erat_2Erat)\ c_2Erat_2Erat_equiv)\ c_2Erat_2Eabs_rat)\ c_2Erat_2Erep_rat)) \quad (15)$$

Theorem 1

$$(\forall V0r \in ty_2Erat_2Erat.((ap\ c_2Erat_2Eabs_rat\ (ap\ c_2Erat_2Erep_rat\ V0r)) = V0r))$$