

thm_2Erat_2ERAT__EQ__ALT (TM- TYHrgyLeoa3wVTLUVpeXbq9qFwJpEPJn5)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define $c_2Ebool_2E_21$ to be $(ap (ap (c_2Emin_2E_3D (2^2))) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x)$

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \tag{1}$$

Let $ty_2Epair_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty_2Epair_2Eprod\ A0\ A1) \tag{2}$$

Let $ty_2Einteger_2Eint : \iota$ be given. Assume the following.

$$nonempty\ ty_2Einteger_2Eint \tag{3}$$

Let $c_2Einteger_2Eint_REP_CLASS : \iota$ be given. Assume the following.

$$c_2Einteger_2Eint_REP_CLASS \in ((2^{(ty_2Epair_2Eprod\ ty_2Enum_2Enum\ ty_2Enum_2Enum)})^{ty_2Einteger_2Eint}) \tag{4}$$

Definition 3 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A.if (\exists x \in A.p (ap\ P\ x))$ then (the $(\lambda x.x \in A \wedge p)$ of type $\iota \Rightarrow \iota$).

Definition 4 We define $c_2Ebool_2E_21$ to be $\lambda A.27a : \iota.(\lambda V0P \in (2^{A-27a}).(ap (ap (c_2Emin_2E_3D (2^{A-27a})))$

Definition 5 We define $c_2Einteger_2Eint_REP$ to be $\lambda V0a \in ty_2Einteger_2Eint.(ap (c_2Emin_2E_40 (ty$

Let $c_2Einteger_2Eint_mul : \iota$ be given. Assume the following.

$$c_2Einteger_2Eint_mul \in (((ty_2Epair_2Eprod\ ty_2Enum_2Enum\ ty_2Enum_2Enum)^{ty_2Epair_2Eprod\ ty_2Enum_2Enum\ ty_2Enum_2Enum})^{ty_2Epair_2Eprod\ ty_2Enum_2Enum\ ty_2Enum_2Enum}) \tag{5}$$

Let $c_2Einteger_2Etint_eq : \iota$ be given. Assume the following.

$$c_2Einteger_2Etint_eq \in ((2^{(ty_2Epair_2Eprod\ ty_2Enum_2Enum\ ty_2Enum_2Enum)})^{(ty_2Epair_2Eprod\ ty_2Enum_2Enum)}) \quad (6)$$

Let $c_2Einteger_2Eint_ABS_CLASS : \iota$ be given. Assume the following.

$$c_2Einteger_2Eint_ABS_CLASS \in (ty_2Einteger_2Eint)^{(2^{(ty_2Epair_2Eprod\ ty_2Enum_2Enum\ ty_2Enum_2Enum)})} \quad (7)$$

Definition 6 We define $c_2Einteger_2Eint_ABS$ to be $\lambda V0r \in (ty_2Epair_2Eprod\ ty_2Enum_2Enum\ ty_2Enum_2Enum)$

Definition 7 We define $c_2Einteger_2Eint_mul$ to be $\lambda V0T1 \in ty_2Einteger_2Eint.\lambda V1T2 \in ty_2Einteger$

Let $ty_2Efrac_2Efrac : \iota$ be given. Assume the following.

$$nonempty\ ty_2Efrac_2Efrac \quad (8)$$

Let $c_2Efrac_2Erep_frac : \iota$ be given. Assume the following.

$$c_2Efrac_2Erep_frac \in ((ty_2Epair_2Eprod\ ty_2Einteger_2Eint\ ty_2Einteger_2Eint)^{ty_2Efrac_2Efrac}) \quad (9)$$

Let $c_2Epair_2ESND : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow \forall A_27b.nonempty\ A_27b \Rightarrow c_2Epair_2ESND\ A_27a\ A_27b \in (A_27b)^{(ty_2Epair_2Eprod\ A_27a\ A_27b)} \quad (10)$$

Definition 8 We define $c_2Efrac_2Efrac_dnm$ to be $\lambda V0f \in ty_2Efrac_2Efrac.(ap\ (c_2Epair_2ESND\ ty_2Efrac_2Efrac\ f))$

Let $c_2Epair_2EFST : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow \forall A_27b.nonempty\ A_27b \Rightarrow c_2Epair_2EFST\ A_27a\ A_27b \in (A_27a)^{(ty_2Epair_2Eprod\ A_27a\ A_27b)} \quad (11)$$

Definition 9 We define $c_2Efrac_2Efrac_nmr$ to be $\lambda V0f \in ty_2Efrac_2Efrac.(ap\ (c_2Epair_2EFST\ ty_2Efrac_2Efrac\ f))$

Definition 10 We define $c_2Erat_2Erat_equiv$ to be $\lambda V0f1 \in ty_2Efrac_2Efrac.\lambda V1f2 \in ty_2Efrac_2Efrac$

Let $ty_2Erat_2Erat : \iota$ be given. Assume the following.

$$nonempty\ ty_2Erat_2Erat \quad (12)$$

Let $c_2Erat_2Eabs_rat_CLASS : \iota$ be given. Assume the following.

$$c_2Erat_2Eabs_rat_CLASS \in (ty_2Erat_2Erat)^{(2^{ty_2Efrac_2Efrac})} \quad (13)$$

Definition 11 We define $c_2Erat_2Eabs_rat$ to be $\lambda V0r \in ty_2Efrac_2Efrac.(ap\ c_2Erat_2Eabs_rat_CLASS\ r)$

Definition 12 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o\ (p\ P \Rightarrow p\ Q)$ of type ι .

Definition 13 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21) 2) (\lambda V2t \in 2.))$

Let $c_2Erat_2Erep_rat_CLASS : \iota$ be given. Assume the following.

$$c_2Erat_2Erep_rat_CLASS \in ((2^{ty_2Efrac_2Efrac})^{ty_2Erat_2Erat}) \quad (14)$$

Definition 14 We define $c_2Erat_2Erep_rat$ to be $\lambda V0a \in ty_2Erat_2Erat.(ap (c_2Emin_2E_40) ty_2Efrac$

Definition 15 We define $c_2Erat_2Erat_nmr$ to be $\lambda V0r \in ty_2Erat_2Erat.(ap c_2Efrac_2Efrac_nmr (ap c$

Definition 16 We define $c_2Erat_2Erat_dnm$ to be $\lambda V0r \in ty_2Erat_2Erat.(ap c_2Efrac_2Efrac_dnm (ap c$

Assume the following.

$$True \quad (15)$$

Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow (\forall V0x \in A_27a.((V0x = V0x) \Leftrightarrow True)) \quad (16)$$

Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow (\forall V0x \in A_27a.(\forall V1y \in A_27a.((V0x = V1y) \Leftrightarrow (V1y = V0x)))) \quad (17)$$

Assume the following.

$$\begin{aligned} & ((\forall V0a \in ty_2Erat_2Erat.((ap c_2Erat_2Eabs_rat (ap c_2Erat_2Erep_rat \\ & V0a)) = V0a)) \wedge (\forall V1r \in ty_2Efrac_2Efrac.(\forall V2s \in ty_2Efrac_2Efrac. \\ & ((p (ap (ap c_2Erat_2Erat_equiv V1r) V2s)) \Leftrightarrow ((ap c_2Erat_2Eabs_rat \\ & V1r) = (ap c_2Erat_2Eabs_rat V2s)))))) \end{aligned} \quad (18)$$

Theorem 1

$$\begin{aligned} & (\forall V0r1 \in ty_2Erat_2Erat.(\forall V1r2 \in ty_2Erat_2Erat. \\ & ((V0r1 = V1r2) \Leftrightarrow ((ap (ap c_2Einteger_2Eint_mul (ap c_2Erat_2Erat_nmr \\ & V0r1)) (ap c_2Erat_2Erat_dnm V1r2)) = (ap (ap c_2Einteger_2Eint_mul \\ & (ap c_2Erat_2Erat_nmr V1r2)) (ap c_2Erat_2Erat_dnm V0r1)))))) \end{aligned}$$