

thm_2Erat_2ERAT__NMRGT0__CONG
 (TMVzDnArSp-
 wXqzaTyQ6XBCHbZQbW7P4Y3bv)

October 26, 2020

Definition 1 We define $c_2Enum_2E_3D$ to be $\lambda A. \lambda x \in A. \lambda y \in A. inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define c_2Ebool_2ET to be $(ap (ap (c_2Enum_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Let $c_2Enum_2EZERO_REP : \iota$ be given. Assume the following.

$$c_2Enum_2EZERO_REP \in \omega \quad (1)$$

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \quad (2)$$

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum^\omega) \quad (3)$$

Definition 3 We define c_2Enum_2E0 to be $(ap c_2Enum_2EABS_num c_2Enum_2EZERO_REP)$.

Let $ty_2Einteger_2Eint : \iota$ be given. Assume the following.

$$nonempty\ ty_2Einteger_2Eint \quad (4)$$

Let $c_2Einteger_2Eint_of_num : \iota$ be given. Assume the following.

$$c_2Einteger_2Eint_of_num \in (ty_2Einteger_2Eint^{ty_2Enum_2Enum}) \quad (5)$$

Let $ty_2Epair_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} & \forall A0. nonempty\ A0 \Rightarrow \forall A1. nonempty\ A1 \Rightarrow nonempty\ (ty_2Epair_2Eprod \\ & \quad A0\ A1) \end{aligned} \quad (6)$$

Let $ty_2Efrac_2Efrac : \iota$ be given. Assume the following.

$$nonempty\ ty_2Efrac_2Efrac \quad (7)$$

Let $c_2Efrac_2Erep_frac : \iota$ be given. Assume the following.

$$c_2Efrac_2Erep_frac \in ((ty_2Epair_2Eprod\ ty_2Einteger_2Eint\ ty_2Einteger_2Eint)^{ty_2Efrac_2Efrac}) \quad (8)$$

Let $c_2Epair_2EFST : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} \forall A_27a.\text{nonempty } A_27a \Rightarrow \forall A_27b.\text{nonempty } A_27b \Rightarrow c_2Epair_2EFST \\ A_27a\ A_27b \in (A_27a^{(ty_2Epair_2Eprod\ A_27a\ A_27b)}) \end{aligned} \quad (9)$$

Definition 4 We define $c_2Ebool_2E_21$ to be $\lambda A_27a : \iota. (\lambda V0P \in (2^{A_27a}). (ap\ (ap\ (c_2Emin_2E_3D\ (2^{A_27a}))\ P)\ V0))$

Definition 5 We define $c_2Efrac_2Efrac_nrm$ to be $\lambda V0f \in ty_2Efrac_2Efrac. (ap\ (c_2Epair_2EFST\ ty_2Efrac_2Efrac))$

Let $c_2Einteger_2Eint_REP_CLASS : \iota$ be given. Assume the following.

$$c_2Einteger_2Eint_REP_CLASS \in ((2^{(ty_2Epair_2Eprod\ ty_2Enum_2Enum\ ty_2Enum_2Enum)})^{ty_2Einteger_2Eint}) \quad (10)$$

Definition 6 We define $c_2Emin_2E_40$ to be $\lambda A. \lambda P \in 2^A. \text{if } (\exists x \in A. p\ (ap\ P\ x)) \text{ then } (\lambda x. x \in A \wedge p\ x) \text{ of type } \iota \Rightarrow \iota$.

Definition 7 We define $c_2Einteger_2Eint_REP$ to be $\lambda V0a \in ty_2Einteger_2Eint. (ap\ (c_2Emin_2E_40\ (ty_2Einteger_2Eint)))$

Let $c_2Einteger_2Etint_lt : \iota$ be given. Assume the following.

$$c_2Einteger_2Etint_lt \in ((2^{(ty_2Epair_2Eprod\ ty_2Enum_2Enum\ ty_2Enum_2Enum)})^{(ty_2Epair_2Eprod\ ty_2Enum_2Enum)}) \quad (11)$$

Definition 8 We define $c_2Einteger_2Eint_lt$ to be $\lambda V0T1 \in ty_2Einteger_2Eint. \lambda V1T2 \in ty_2Einteger_2Eint. (V0T1 \wedge V1T2)$

Definition 9 We define $c_2Einteger_2Eint_gt$ to be $\lambda V0x \in ty_2Einteger_2Eint. \lambda V1y \in ty_2Einteger_2Eint. (V0x \wedge V1y)$

Definition 10 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2. \lambda Q \in 2. \text{inj_o } (p\ P \Rightarrow p\ Q)$ of type ι .

Let $c_2Epair_2ESND : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} \forall A_27a.\text{nonempty } A_27a \Rightarrow \forall A_27b.\text{nonempty } A_27b \Rightarrow c_2Epair_2ESND \\ A_27a\ A_27b \in (A_27b^{(ty_2Epair_2Eprod\ A_27a\ A_27b)}) \end{aligned} \quad (12)$$

Definition 11 We define $c_2Efrac_2Efrac_dnm$ to be $\lambda V0f \in ty_2Efrac_2Efrac. (ap\ (c_2Epair_2ESND\ ty_2Efrac_2Efrac))$

Let $c_2Einteger_2Etint_mul : \iota$ be given. Assume the following.

$$c_2Einteger_2Etint_mul \in (((ty_2Epair_2Eprod\ ty_2Enum_2Enum\ ty_2Enum_2Enum)^{(ty_2Epair_2Eprod\ ty_2Enum_2Enum\ ty_2Enum_2Enum)})^{(ty_2Epair_2Eprod\ ty_2Enum_2Enum\ ty_2Enum_2Enum)}) \quad (13)$$

Let $c_2Einteger_2Etint_eq : \iota$ be given. Assume the following.

$$c_2Einteger_2Etint_eq \in ((2^{(ty_2Epair_2Eprod\ ty_2Enum_2Enum\ ty_2Enum_2Enum)})^{(ty_2Epair_2Eprod\ ty_2Enum)})^{(ty_2Epair_2Eprod\ ty_2Enum)} \quad (14)$$

Let $c_2Einteger_2Eint_ABS_CLASS : \iota$ be given. Assume the following.

$$c_2Einteger_2Eint_ABS_CLASS \in (ty_2Einteger_2Eint^{(2^{(ty_2Epair_2Eprod\ ty_2Enum_2Enum\ ty_2Enum_2Enum)})})$$

Definition 12 We define $c_2Einteger_2Eint_ABS$ to be $\lambda V0r \in (ty_2Epair_2Eprod\ ty_2Enum_2Enum\ ty_2Eint)$

Definition 13 We define $c_2Einteger_2Eint_mul$ to be $\lambda V0T1 \in ty_2Einteger_2Eint. \lambda V1T2 \in ty_2Einteger_2Eint.$

Let $ty_2Erat_2Erat : \iota$ be given. Assume the following.

nonempty *ty_2Erat_2Erat* (16)

Let $c_2Erat_2Erep_rat_CLASS : \iota$ be given. Assume the following.

$$c_2Erat_2Erep_rat_CLASS \in ((2^{ty_2Efrac_2Efrac})^{ty_2Erat_2Erat}) \quad (17)$$

Definition 15 We define $c_2Erat_2Erep_rat$ to be $\lambda V o \in ty_2Erat_2Erat.(ap\ (c_2Emin_2E_40\ ty_2Efraction)\ o)$

Let $c_2Erat_2Eabs_rat_CLASS : \iota$ be given. Assume the following.

$$c_2Erat_2Eabs_rat_CLASS \in (ty_2Erat_2Erat^{(2^{ty_2Efrac_2Efrac})}) \quad (18)$$

Definition 16 We define $c_{\text{2Erat_2Eabs_rat}}$ to be $\lambda V0r \in ty_2Efrac_2Efrac.(ap\ c_{\text{2Erat_2Eabs_rat_CI}}\ r)$

Definition 17 We define $c_{\text{Ebool}} : \mathcal{C}_F \rightarrow \mathcal{C}$ to be $(\lambda V0t1 \in 2. (\lambda V1t2 \in 2. (ap (c_{\text{Ebool}}_2) t1) t2)) (\lambda V2t \in 2. (ap (c_{\text{Ebool}}_1) t))$

Assume the following.

True (19)

Assume the following.

$$\forall A_27a.\text{nonempty } A_27a \Rightarrow (\forall V0t \in 2.((\forall V1x \in A_27a.(p \ V0t)) \Leftrightarrow (p \ V0t))) \quad (20)$$

Assume the following.

$$\forall A. \text{nonempty } A \Rightarrow (\forall V0x \in A. ((V0x = V0x) \Leftrightarrow \text{True})) \quad (21)$$

Assume the following.

$$(\forall V0a \in ty_2Efrac_2Efrac. (\forall V1b \in ty_2Efrac_2Efrac. ((p (ap (ap c_2Erat_2Erat_equiv V0a) V1b)) \Leftrightarrow (p (ap (ap c_2Erat_2Erat_equiv V1b) V0a)))))) \quad (22)$$

Assume the following.

$$\begin{aligned}
 & (\forall V0a \in ty_2Efrac_2Efrac. (\forall V1b \in ty_2Efrac_2Efrac. \\
 & ((p (ap (ap c_2Erat_2Erat_equiv V0a) V1b)) \Rightarrow ((p (ap (ap c_2Einteger_2Eint_gt \\
 & (ap c_2Efrac_2Efrac_nmr V0a)) (ap c_2Einteger_2Eint_of_num \\
 & c_2Enum_2E0))) \Leftrightarrow (p (ap (ap c_2Einteger_2Eint_gt (ap c_2Efrac_2Efrac_nmr \\
 & V1b)) (ap c_2Einteger_2Eint_of_num c_2Enum_2E0))))))) \\
 & (23)
 \end{aligned}$$

Assume the following.

$$\begin{aligned}
 & ((\forall V0a \in ty_2Erat_2Erat. ((ap c_2Erat_2Eabs_rat (ap c_2Erat_2Erep_rat \\
 & V0a)) = V0a)) \wedge (\forall V1r \in ty_2Efrac_2Efrac. (\forall V2s \in ty_2Efrac_2Efrac. \\
 & ((p (ap (ap c_2Erat_2Erat_equiv V1r) V2s)) \Leftrightarrow ((ap c_2Erat_2Eabs_rat \\
 & V1r) = (ap c_2Erat_2Eabs_rat V2s)))))) \\
 & (24)
 \end{aligned}$$

Theorem 1

$$\begin{aligned}
 & (\forall V0f1 \in ty_2Efrac_2Efrac. ((p (ap (ap c_2Einteger_2Eint_gt \\
 & (ap c_2Efrac_2Efrac_nmr (ap c_2Erat_2Erep_rat (ap c_2Erat_2Eabs_rat \\
 & V0f1)))) (ap c_2Einteger_2Eint_of_num c_2Enum_2E0))) \Leftrightarrow (p (\\
 & ap (ap c_2Einteger_2Eint_gt (ap c_2Efrac_2Efrac_nmr V0f1)) \\
 & (ap c_2Einteger_2Eint_of_num c_2Enum_2E0))))))
 \end{aligned}$$