

thm_2Erat_2ERAT__NMRLT0__CONG
(TMYnd2U9LR3iXZCRe4Z4zmuymcDdMSVVvc1)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define c_2Ebool_2ET to be $(ap (ap (c_2Emin_2E_3D (2^2))) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Let $c_2Enum_2EZERO_REP : \iota$ be given. Assume the following.

$$c_2Enum_2EZERO_REP \in \omega \tag{1}$$

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \tag{2}$$

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum^{\omega}) \tag{3}$$

Definition 3 We define c_2Enum_2E0 to be $(ap\ c_2Enum_2EABS_num\ c_2Enum_2EZERO_REP)$.

Let $ty_2Einteger_2Eint : \iota$ be given. Assume the following.

$$nonempty\ ty_2Einteger_2Eint \tag{4}$$

Let $c_2Einteger_2Eint_of_num : \iota$ be given. Assume the following.

$$c_2Einteger_2Eint_of_num \in (ty_2Einteger_2Eint^{ty_2Enum_2Enum}) \tag{5}$$

Let $ty_2Epair_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty_2Epair_2Eprod\ A0\ A1) \tag{6}$$

Let $ty_2Efrac_2Efrac : \iota$ be given. Assume the following.

$$nonempty\ ty_2Efrac_2Efrac \tag{7}$$

Let $c_2Efrac_2Erep_frac : \iota$ be given. Assume the following.

$$c_2Efrac_2Erep_frac \in ((ty_2Epair_2Eprod\ ty_2Einteger_2Eint\ ty_2Einteger_2Eint)^{ty_2Efrac_2Efrac}) \quad (8)$$

Let $c_2Epair_2EFST : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow \forall A_27b.nonempty\ A_27b \Rightarrow c_2Epair_2EFST\ A_27a\ A_27b \in (A_27a^{(ty_2Epair_2Eprod\ A_27a\ A_27b)}) \quad (9)$$

Definition 4 We define $c_2Ebool_2E_21$ to be $\lambda A_27a : \iota. (\lambda V0P \in (2^{A_27a}). (ap\ (ap\ (c_2Emin_2E_3D\ (2^{A_27a})))$

Definition 5 We define $c_2Efrac_2Efrac_nmr$ to be $\lambda V0f \in ty_2Efrac_2Efrac. (ap\ (c_2Epair_2EFST\ ty_2Efrac_2Efrac\ f))$

Let $c_2Einteger_2Eint_REP_CLASS : \iota$ be given. Assume the following.

$$c_2Einteger_2Eint_REP_CLASS \in ((2^{(ty_2Epair_2Eprod\ ty_2Enum_2Enum\ ty_2Enum_2Enum)})^{ty_2Einteger_2Eint}) \quad (10)$$

Definition 6 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A. \text{if } (\exists x \in A. p\ (ap\ P\ x)) \text{ then } (the\ (\lambda x. x \in A \wedge p\ x))$ of type ι .

Definition 7 We define $c_2Einteger_2Eint_REP$ to be $\lambda V0a \in ty_2Einteger_2Eint. (ap\ (c_2Emin_2E_40\ ty_2Einteger_2Eint\ a))$

Let $c_2Einteger_2Etint_lt : \iota$ be given. Assume the following.

$$c_2Einteger_2Etint_lt \in ((2^{(ty_2Epair_2Eprod\ ty_2Enum_2Enum\ ty_2Enum_2Enum)})^{(ty_2Epair_2Eprod\ ty_2Enum_2Enum)}) \quad (11)$$

Definition 8 We define $c_2Einteger_2Eint_lt$ to be $\lambda V0T1 \in ty_2Einteger_2Eint. \lambda V1T2 \in ty_2Einteger_2Eint. (c_2Einteger_2Etint_lt\ T1\ T2)$

Definition 9 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2. \lambda Q \in 2. inj_o\ (p\ P \Rightarrow p\ Q)$ of type ι .

Let $c_2Epair_2ESND : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow \forall A_27b.nonempty\ A_27b \Rightarrow c_2Epair_2ESND\ A_27a\ A_27b \in (A_27b^{(ty_2Epair_2Eprod\ A_27a\ A_27b)}) \quad (12)$$

Definition 10 We define $c_2Efrac_2Efrac_dnm$ to be $\lambda V0f \in ty_2Efrac_2Efrac. (ap\ (c_2Epair_2ESND\ ty_2Efrac_2Efrac\ f))$

Let $c_2Einteger_2Etint_mul : \iota$ be given. Assume the following.

$$c_2Einteger_2Etint_mul \in (((ty_2Epair_2Eprod\ ty_2Enum_2Enum\ ty_2Enum_2Enum)^{ty_2Epair_2Eprod\ ty_2Enum_2Enum})^{(ty_2Epair_2Eprod\ ty_2Enum_2Enum)}) \quad (13)$$

Let $c_2Einteger_2Etint_eq : \iota$ be given. Assume the following.

$$c_2Einteger_2Etint_eq \in ((2^{(ty_2Epair_2Eprod\ ty_2Enum_2Enum\ ty_2Enum_2Enum)})^{(ty_2Epair_2Eprod\ ty_2Enum_2Enum)}) \quad (14)$$

Let $c_2Einteger_2Eint_ABS_CLASS : \iota$ be given. Assume the following.

$$c_2Einteger_2Eint_ABS_CLASS \in (ty_2Einteger_2Eint)^{(2^{(ty_2Epair_2Eprod\ ty_2Enum_2Enum\ ty_2Enum_2Enum)})} \quad (15)$$

Definition 11 We define $c_Einteger_Eint_ABS$ to be $\lambda V0r \in (ty_Epair_Eprod\ ty_Eenum_Eenum\ ty_Eint_Eint)$

Definition 12 We define $c_Einteger_Eint_mul$ to be $\lambda V0T1 \in ty_Einteger_Eint.\lambda V1T2 \in ty_Einteger_Eint$

Definition 13 We define $c_Erat_Erat_equiv$ to be $\lambda V0f1 \in ty_Efrac_Efrac.\lambda V1f2 \in ty_Efrac_Efrac$

Let $ty_Erat_Erat : \iota$ be given. Assume the following.

$$nonempty\ ty_Erat_Erat \quad (16)$$

Let $c_Erat_Erep_rat_CLASS : \iota$ be given. Assume the following.

$$c_Erat_Erep_rat_CLASS \in ((2^{ty_Efrac_Efrac})^{ty_Erat_Erat}) \quad (17)$$

Definition 14 We define $c_Erat_Erep_rat$ to be $\lambda V0a \in ty_Erat_Erat.(ap\ (c_Emin_E40\ ty_Efrac_Efrac))$

Let $c_Erat_Eabs_rat_CLASS : \iota$ be given. Assume the following.

$$c_Erat_Eabs_rat_CLASS \in (ty_Erat_Erat^{(2^{ty_Efrac_Efrac})}) \quad (18)$$

Definition 15 We define $c_Erat_Eabs_rat$ to be $\lambda V0r \in ty_Efrac_Efrac.(ap\ c_Erat_Eabs_rat_CLASS)$

Definition 16 We define c_Ebool_E2F5C to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap\ (c_Ebool_E21\ 2))\ (\lambda V2t \in 2)))$

Assume the following.

$$True \quad (19)$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0t \in 2.((\forall V1x \in A_27a.(p\ V0t)) \Leftrightarrow (p\ V0t))) \quad (20)$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0x \in A_27a.((V0x = V0x) \Leftrightarrow True)) \quad (21)$$

Assume the following.

$$(\forall V0a \in ty_Efrac_Efrac.(\forall V1b \in ty_Efrac_Efrac.((p\ (ap\ (ap\ c_Erat_Erat_equiv\ V0a)\ V1b)) \Leftrightarrow (p\ (ap\ (ap\ c_Erat_Erat_equiv\ V1b)\ V0a)))))) \quad (22)$$

Assume the following.

$$(\forall V0a \in ty_Efrac_Efrac.(\forall V1b \in ty_Efrac_Efrac.((p\ (ap\ (ap\ c_Erat_Erat_equiv\ V0a)\ V1b)) \Rightarrow ((p\ (ap\ (ap\ c_Einteger_Eint_lt\ (ap\ c_Efrac_Efrac_nmr\ V0a))\ (ap\ c_Einteger_Eint_of_num\ c_Eenum_E0))) \Leftrightarrow (p\ (ap\ (ap\ c_Einteger_Eint_lt\ (ap\ c_Efrac_Efrac_nmr\ V1b))\ (ap\ c_Einteger_Eint_of_num\ c_Eenum_E0))))))) \quad (23)$$

Assume the following.

$$\begin{aligned}
& ((\forall V0a \in ty_2Erat_2Erat. ((ap\ c_2Erat_2Eabs_rat\ (ap\ c_2Erat_2Erep_rat \\
& V0a)) = V0a)) \wedge (\forall V1r \in ty_2Efrac_2Efrac. (\forall V2s \in ty_2Efrac_2Efrac. \\
& ((p\ (ap\ (ap\ c_2Erat_2Erat_equiv\ V1r)\ V2s)) \Leftrightarrow ((ap\ c_2Erat_2Eabs_rat \\
& V1r) = (ap\ c_2Erat_2Eabs_rat\ V2s))))))
\end{aligned} \tag{24}$$

Theorem 1

$$\begin{aligned}
& (\forall V0f1 \in ty_2Efrac_2Efrac. ((p\ (ap\ (ap\ c_2Einteger_2Eint_lt \\
& (ap\ c_2Efrac_2Efrac_nmr\ (ap\ c_2Erat_2Erep_rat\ (ap\ c_2Erat_2Eabs_rat \\
& V0f1))))\ (ap\ c_2Einteger_2Eint_of_num\ c_2Enum_2E0))) \Leftrightarrow (p\ (\\
& ap\ (ap\ c_2Einteger_2Eint_lt\ (ap\ c_2Efrac_2Efrac_nmr\ V0f1)) \\
& (ap\ c_2Einteger_2Eint_of_num\ c_2Enum_2E0))))))
\end{aligned}$$