

thm_2Erat_2ERAT__SGN__CONG (TM- cnX6FUm4eQfWKny5oqWAnrqTSygR5B6CN)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define c_2Ebool_2ET to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Let $ty_2Einteger_2Eint : \iota$ be given. Assume the following.

$$nonempty\ ty_2Einteger_2Eint \tag{1}$$

Let $ty_2Epair_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty_2Epair_2Eprod\ A0\ A1) \tag{2}$$

Let $ty_2Efrac_2Efrac : \iota$ be given. Assume the following.

$$nonempty\ ty_2Efrac_2Efrac \tag{3}$$

Let $c_2Efrac_2Erep_frac : \iota$ be given. Assume the following.

$$c_2Efrac_2Erep_frac \in ((ty_2Epair_2Eprod\ ty_2Einteger_2Eint\ ty_2Einteger_2Eint)^{ty_2Efrac_2Efrac}) \tag{4}$$

Let $c_2Epair_2EFST : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A.27a.nonempty\ A.27a \Rightarrow \forall A.27b.nonempty\ A.27b \Rightarrow c_2Epair_2EFST\ A.27a\ A.27b \in (A.27a^{(ty_2Epair_2Eprod\ A.27a\ A.27b)}) \tag{5}$$

Definition 3 We define $c_2Ebool_2E_21$ to be $\lambda A.27a : \iota.(\lambda V0P \in (2^{A.27a}).(ap (ap (c_2Emin_2E_3D (2^{A.27a}))$

Definition 4 We define $c_2Efrac_2Efrac_nmr$ to be $\lambda V0f \in ty_2Efrac_2Efrac.(ap (c_2Epair_2EFST\ ty_2Efrac_2Efrac$

Let $c_2Enum_2EZERO_REP : \iota$ be given. Assume the following.

$$c_2Enum_2EZERO_REP \in omega \tag{6}$$

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \quad (7)$$

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum^{omega}) \quad (8)$$

Definition 5 We define c_2Enum_2E0 to be $(ap\ c_2Enum_2EABS_num\ c_2Enum_2EZERO_REP)$.

Definition 6 We define $c_2Earithmic_2EZERO$ to be c_2Enum_2E0 .

Let $c_2Enum_2EREP_num : \iota$ be given. Assume the following.

$$c_2Enum_2EREP_num \in (omega^{ty_2Enum_2Enum}) \quad (9)$$

Let $c_2Enum_2ESUC_REP : \iota$ be given. Assume the following.

$$c_2Enum_2ESUC_REP \in (omega^{omega}) \quad (10)$$

Definition 7 We define c_2Enum_2ESUC to be $\lambda V0m \in ty_2Enum_2Enum.(ap\ c_2Enum_2EABS_num\ ($

Let $c_2Earithmic_2E_2B : \iota$ be given. Assume the following.

$$c_2Earithmic_2E_2B \in ((ty_2Enum_2Enum^{ty_2Enum_2Enum})^{ty_2Enum_2Enum}) \quad (11)$$

Definition 8 We define $c_2Earithmic_2EBIT1$ to be $\lambda V0n \in ty_2Enum_2Enum.(ap\ (ap\ c_2Earithmic_2E_2B\ ($

Definition 9 We define $c_2Earithmic_2ENUMERAL$ to be $\lambda V0x \in ty_2Enum_2Enum.V0x$.

Let $c_2Einteger_2Eint_of_num : \iota$ be given. Assume the following.

$$c_2Einteger_2Eint_of_num \in (ty_2Einteger_2Eint^{ty_2Enum_2Enum}) \quad (12)$$

Let $c_2Einteger_2Eint_REP_CLASS : \iota$ be given. Assume the following.

$$c_2Einteger_2Eint_REP_CLASS \in ((2^{(ty_2Epair_2Eprod\ ty_2Enum_2Enum\ ty_2Enum_2Enum)})^{ty_2Einteger_2Eint}) \quad (13)$$

Definition 10 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A.\text{if } (\exists x \in A.p\ (ap\ P\ x)) \text{ then } (the\ (\lambda x.x \in A \wedge P\ x)) \text{ of type } \iota \Rightarrow \iota$.

Definition 11 We define $c_2Einteger_2Eint_REP$ to be $\lambda V0a \in ty_2Einteger_2Eint.(ap\ (c_2Emin_2E_40\ ($

Let $c_2Einteger_2Etint_neg : \iota$ be given. Assume the following.

$$c_2Einteger_2Etint_neg \in ((ty_2Epair_2Eprod\ ty_2Enum_2Enum\ ty_2Enum_2Enum)^{(ty_2Epair_2Eprod\ ty_2Enum_2Enum\ ty_2Enum_2Enum)}) \quad (14)$$

Let $c_2Einteger_2Etint_eq : \iota$ be given. Assume the following.

$$c_2Einteger_2Etint_eq \in ((2^{(ty_2Epair_2Eprod\ ty_2Enum_2Enum\ ty_2Enum_2Enum)})^{(ty_2Epair_2Eprod\ ty_2Enum_2Enum\ ty_2Enum_2Enum)}) \quad (15)$$

Let $c_2Einteger_2Eint_ABS_CLASS : \iota$ be given. Assume the following.

$$c_2Einteger_2Eint_ABS_CLASS \in (ty_2Einteger_2Eint^{(2^{(ty_2Epair_2Eprod\ ty_2Enum_2Enum\ ty_2Enum_2Enum)})}) \quad (16)$$

Definition 12 We define $c_2Einteger_2Eint_ABS$ to be $\lambda V0r \in (ty_2Epair_2Eprod\ ty_2Enum_2Enum\ ty_2Enum_2Enum)$

Definition 13 We define $c_2Einteger_2Eint_neg$ to be $\lambda V0T1 \in ty_2Einteger_2Eint.(ap\ c_2Einteger_2Eint$

Let $c_2Einteger_2Eint_lt : \iota$ be given. Assume the following.

$$c_2Einteger_2Eint_lt \in ((2^{(ty_2Epair_2Eprod\ ty_2Enum_2Enum\ ty_2Enum_2Enum)})^{(ty_2Epair_2Eprod\ ty_2Enum_2Enum)}) \quad (17)$$

Definition 14 We define $c_2Einteger_2Eint_lt$ to be $\lambda V0T1 \in ty_2Einteger_2Eint.\lambda V1T2 \in ty_2Einteger$

Definition 15 We define c_2Ebool_2EF to be $(ap\ (c_2Ebool_2E_21\ 2)\ (\lambda V0t \in 2.V0t))$.

Definition 16 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o\ (p\ P \Rightarrow p\ Q)$ of type ι .

Definition 17 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap\ (c_2Ebool_2E_21\ 2)\ (\lambda V2t \in 2.V2t))))$

Definition 18 We define c_2Ebool_2ECOND to be $\lambda A_27a : \iota.(\lambda V0t \in 2.(\lambda V1t1 \in A_27a.(\lambda V2t2 \in A_27a.(ap\ (c_2Ebool_2E_21\ 2)\ (\lambda V3t3 \in 2.V3t3))))))$

Definition 19 We define $c_2EintExtension_2ESGN$ to be $\lambda V0x \in ty_2Einteger_2Eint.(ap\ (ap\ (ap\ (c_2Ebool_2E_21\ 2)\ (\lambda V1t1 \in 2.V1t1))))$

Definition 20 We define $c_2Efrac_2Efrac_sgn$ to be $\lambda V0f1 \in ty_2Efrac_2Efrac.(ap\ c_2EintExtension_2ESGN)$

Let $c_2Epair_2ESND : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow \forall A_27b.nonempty\ A_27b \Rightarrow c_2Epair_2ESND\ A_27a\ A_27b \in (A_27b^{(ty_2Epair_2Eprod\ A_27a\ A_27b)}) \quad (18)$$

Definition 21 We define $c_2Efrac_2Efrac_dnm$ to be $\lambda V0f \in ty_2Efrac_2Efrac.(ap\ (c_2Epair_2ESND\ t1\ t2))$

Let $c_2Einteger_2Eint_mul : \iota$ be given. Assume the following.

$$c_2Einteger_2Eint_mul \in (((ty_2Epair_2Eprod\ ty_2Enum_2Enum\ ty_2Enum_2Enum)^{(ty_2Epair_2Eprod\ ty_2Enum_2Enum\ ty_2Enum_2Enum)})^{(ty_2Epair_2Eprod\ ty_2Enum_2Enum\ ty_2Enum_2Enum)}) \quad (19)$$

Definition 22 We define $c_2Einteger_2Eint_mul$ to be $\lambda V0T1 \in ty_2Einteger_2Eint.\lambda V1T2 \in ty_2Einteger$

Definition 23 We define $c_2Erat_2Erat_equiv$ to be $\lambda V0f1 \in ty_2Efrac_2Efrac.\lambda V1f2 \in ty_2Efrac_2Efrac$

Let $ty_2Erat_2Erat : \iota$ be given. Assume the following.

$$nonempty\ ty_2Erat_2Erat \quad (20)$$

Let $c_2Erat_2Eabs_rat_CLASS : \iota$ be given. Assume the following.

$$c_2Erat_2Eabs_rat_CLASS \in (ty_2Erat_2Erat^{(2^{ty_2Efrac_2Efrac})}) \quad (21)$$

Definition 24 We define $c_2Erat_2Eabs_rat$ to be $\lambda V0r \in ty_2Efrac_2Efrac.(ap\ c_2Erat_2Eabs_rat_CLASS)$

Let $c_2Erat_2Erep_rat_CLASS : \iota$ be given. Assume the following.

$$c_2Erat_2Erep_rat_CLASS \in ((2^{ty_2Efrac_2Efrac})^{ty_2Erat_2Erat}) \quad (22)$$

Definition 25 We define $c_2Erat_2Erep_rat$ to be $\lambda V0a \in ty_2Erat_2Erat.(ap (c_2Emin_2E_40 ty_2Efrac$

Assume the following.

$$True \quad (23)$$

Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow (\forall V0x \in A_27a.((V0x = V0x) \Leftrightarrow True)) \quad (24)$$

Assume the following.

$$\begin{aligned} & (\forall V0f1 \in ty_2Efrac_2Efrac.(((ap c_2Efrac_2Efrac_nmr \\ & (ap c_2Erat_2Erep_rat (ap c_2Erat_2Eabs_rat V0f1))) = (ap c_2Einteger_2Eint_of_num \\ & c_2Enum_2E0)) \Leftrightarrow ((ap c_2Efrac_2Efrac_nmr V0f1) = (ap c_2Einteger_2Eint_of_num \\ & c_2Enum_2E0)))) \end{aligned} \quad (25)$$

Assume the following.

$$\begin{aligned} & (\forall V0f1 \in ty_2Efrac_2Efrac.((p (ap (ap c_2Einteger_2Eint_lt \\ & (ap c_2Efrac_2Efrac_nmr (ap c_2Erat_2Erep_rat (ap c_2Erat_2Eabs_rat \\ & V0f1)))) (ap c_2Einteger_2Eint_of_num c_2Enum_2E0))) \Leftrightarrow (p (\\ & ap (ap c_2Einteger_2Eint_lt (ap c_2Efrac_2Efrac_nmr V0f1)) \\ & (ap c_2Einteger_2Eint_of_num c_2Enum_2E0)))) \end{aligned} \quad (26)$$

Theorem 1

$$\begin{aligned} & (\forall V0f1 \in ty_2Efrac_2Efrac.((ap c_2Efrac_2Efrac_sgn (\\ & ap c_2Erat_2Erep_rat (ap c_2Erat_2Eabs_rat V0f1))) = (ap c_2Efrac_2Efrac_sgn \\ & V0f1))) \end{aligned}$$