

thm\_2Erat\_2Erat\_\_equiv\_\_reps  
(TMJ8mqwidi24nzkBdBXsEh9qRToX4eyCrZ4)

October 26, 2020

**Definition 1** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define  $c\_2Ebool\_2E\_21$  to be  $(ap (ap (c\_2Emin\_2E\_3D (2^2))) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x)$

Let  $ty\_2Einteger\_2Eint : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Einteger\_2Eint \tag{1}$$

Let  $ty\_2Epair\_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty\_2Epair\_2Eprod\ A0\ A1) \tag{2}$$

Let  $ty\_2Efrac\_2Efrac : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Efrac\_2Efrac \tag{3}$$

Let  $c\_2Efrac\_2Erep\_frac : \iota$  be given. Assume the following.

$$c\_2Efrac\_2Erep\_frac \in ((ty\_2Epair\_2Eprod\ ty\_2Einteger\_2Eint\ ty\_2Einteger\_2Eint)^{ty\_2Efrac\_2Efrac}) \tag{4}$$

Let  $c\_2Epair\_2ESND : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A.27a.nonempty\ A.27a \Rightarrow \forall A.27b.nonempty\ A.27b \Rightarrow c\_2Epair\_2ESND\ A.27a\ A.27b \in (A.27b)^{(ty\_2Epair\_2Eprod\ A.27a\ A.27b)} \tag{5}$$

**Definition 3** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A.27a : \iota.(\lambda V0P \in (2^{A.27a}).(ap (ap (c\_2Emin\_2E\_3D (2^{A.27a})) (\lambda V1x \in 2.V1x)) (\lambda V2x \in 2.V2x)) P))$

**Definition 4** We define  $c\_2Efrac\_2Efrac\_dnm$  to be  $\lambda V0f \in ty\_2Efrac\_2Efrac.(ap (c\_2Epair\_2ESND\ ty\_2Einteger\_2Eint\ ty\_2Einteger\_2Eint)^{ty\_2Efrac\_2Efrac}) f)$

Let  $c\_2Epair\_2EFST : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A.27a.nonempty\ A.27a \Rightarrow \forall A.27b.nonempty\ A.27b \Rightarrow c\_2Epair\_2EFST\ A.27a\ A.27b \in (A.27a)^{(ty\_2Epair\_2Eprod\ A.27a\ A.27b)} \tag{6}$$

**Definition 5** We define  $c\_2Efrac\_2Efrac\_nmr$  to be  $\lambda V0f \in ty\_2Efrac\_2Efrac.(ap (c\_2Epair\_2EFST ty\_2Enum\_2Enum))$ .  
Let  $ty\_2Enum\_2Enum : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Enum\_2Enum \quad (7)$$

Let  $c\_2Einteger\_2Eint\_REP\_CLASS : \iota$  be given. Assume the following.

$$c\_2Einteger\_2Eint\_REP\_CLASS \in ((2^{(ty\_2Epair\_2Eprod\ ty\_2Enum\_2Enum\ ty\_2Enum\_2Enum)}\ ty\_2Einteger\_2Eint)) \quad (8)$$

**Definition 6** We define  $c\_2Emin\_2E.40$  to be  $\lambda A.\lambda P \in 2^A$ . **if**  $(\exists x \in A.p (ap\ P\ x))$  **then** *(the  $(\lambda x.x \in A \wedge p\ x)$  of type  $\iota \Rightarrow \iota$ ).*

**Definition 7** We define  $c\_2Einteger\_2Eint\_REP$  to be  $\lambda V0a \in ty\_2Einteger\_2Eint.(ap (c\_2Emin\_2E.40\ ty\_2Enum\_2Enum))$ .

Let  $c\_2Einteger\_2Eint\_mul : \iota$  be given. Assume the following.

$$c\_2Einteger\_2Eint\_mul \in (((ty\_2Epair\_2Eprod\ ty\_2Enum\_2Enum\ ty\_2Enum\_2Enum)^{(ty\_2Epair\_2Eprod\ ty\_2Enum\_2Enum\ ty\_2Enum\_2Enum)})^{(ty\_2Epair\_2Eprod\ ty\_2Enum\_2Enum\ ty\_2Enum\_2Enum)}) \quad (9)$$

Let  $c\_2Einteger\_2Eint\_eq : \iota$  be given. Assume the following.

$$c\_2Einteger\_2Eint\_eq \in ((2^{(ty\_2Epair\_2Eprod\ ty\_2Enum\_2Enum\ ty\_2Enum\_2Enum)})^{(ty\_2Epair\_2Eprod\ ty\_2Enum\_2Enum\ ty\_2Enum\_2Enum)}) \quad (10)$$

Let  $c\_2Einteger\_2Eint\_ABS\_CLASS : \iota$  be given. Assume the following.

$$c\_2Einteger\_2Eint\_ABS\_CLASS \in (ty\_2Einteger\_2Eint)^{(2^{(ty\_2Epair\_2Eprod\ ty\_2Enum\_2Enum\ ty\_2Enum\_2Enum)})} \quad (11)$$

**Definition 8** We define  $c\_2Einteger\_2Eint\_ABS$  to be  $\lambda V0r \in (ty\_2Epair\_2Eprod\ ty\_2Enum\_2Enum\ ty\_2Enum\_2Enum)$ .

**Definition 9** We define  $c\_2Einteger\_2Eint\_mul$  to be  $\lambda V0T1 \in ty\_2Einteger\_2Eint.\lambda V1T2 \in ty\_2Einteger\_2Eint$ .

**Definition 10** We define  $c\_2Erat\_2Erat\_equiv$  to be  $\lambda V0f1 \in ty\_2Efrac\_2Efrac.\lambda V1f2 \in ty\_2Efrac\_2Efrac$ .

Let  $ty\_2Erat\_2Erat : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Erat\_2Erat \quad (12)$$

Let  $c\_2Erat\_2Erep\_rat\_CLASS : \iota$  be given. Assume the following.

$$c\_2Erat\_2Erep\_rat\_CLASS \in ((2^{ty\_2Efrac\_2Efrac})^{ty\_2Erat\_2Erat}) \quad (13)$$

**Definition 11** We define  $c\_2Erat\_2Erep\_rat$  to be  $\lambda V0a \in ty\_2Erat\_2Erat.(ap (c\_2Emin\_2E.40\ ty\_2Efrac\_2Efrac))$ .

Let  $c\_2Erat\_2Eabs\_rat\_CLASS : \iota$  be given. Assume the following.

$$c\_2Erat\_2Eabs\_rat\_CLASS \in (ty\_2Erat\_2Erat)^{(2^{ty\_2Efrac\_2Efrac})} \quad (14)$$

**Definition 12** We define  $c\_2Erat\_2Eabs\_rat$  to be  $\lambda V0r \in ty\_2Efrac\_2Efrac.(ap\ c\_2Erat\_2Eabs\_rat\_CLASS)$ .

**Definition 13** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o (p P \Rightarrow p Q)$  of type  $\iota$ .

**Definition 14** We define  $c\_2Ebool\_2E\_2F\_5C$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_2Ebool\_2E\_21 2) (\lambda V2t \in$

Assume the following.

$$True \tag{15}$$

Assume the following.

$$\forall A\_27a.nonempty A\_27a \Rightarrow (\forall V0x \in A\_27a.((V0x = V0x) \Leftrightarrow True)) \tag{16}$$

Assume the following.

$$\begin{aligned} & ((\forall V0a \in ty\_2Erat\_2Erat.((ap c\_2Erat\_2Eabs\_rat (ap c\_2Erat\_2Erep\_rat \\ & V0a)) = V0a)) \wedge (\forall V1r \in ty\_2Efrac\_2Efrac.(\forall V2s \in ty\_2Efrac\_2Efrac. \\ & ((p (ap (ap c\_2Erat\_2Erat\_equiv V1r) V2s)) \Leftrightarrow ((ap c\_2Erat\_2Eabs\_rat \\ & V1r) = (ap c\_2Erat\_2Eabs\_rat V2s)))))) \end{aligned} \tag{17}$$

**Theorem 1**

$$\begin{aligned} & (\forall V0r1 \in ty\_2Erat\_2Erat.(\forall V1r2 \in ty\_2Erat\_2Erat. \\ & ((p (ap (ap c\_2Erat\_2Erat\_equiv (ap c\_2Erat\_2Erep\_rat V0r1)) \\ & (ap c\_2Erat\_2Erep\_rat V1r2)))) \Leftrightarrow (V0r1 = V1r2)))) \end{aligned}$$