

thm_2Ereal_2EREAL__10
(TMc2PgBJ8A6PfxNqi8u9ctjGSRdoNNzbeeB)

October 26, 2020

Let $c_2Enum_2EZERO_REP : \iota$ be given. Assume the following.

$$c_2Enum_2EZERO_REP \in \omega \tag{1}$$

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \tag{2}$$

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum^{\omega}) \tag{3}$$

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define c_2Enum_2E0 to be $(ap\ c_2Enum_2EABS_num\ c_2Enum_2EZERO_REP)$.

Definition 3 We define $c_2Earithmetic_2EZERO$ to be c_2Enum_2E0 .

Let $c_2Enum_2EREP_num : \iota$ be given. Assume the following.

$$c_2Enum_2EREP_num \in (\omega^{ty_2Enum_2Enum}) \tag{4}$$

Let $c_2Enum_2ESUC_REP : \iota$ be given. Assume the following.

$$c_2Enum_2ESUC_REP \in (\omega^{\omega}) \tag{5}$$

Definition 4 We define c_2Ebool_2ET to be $(ap\ (ap\ (c_2Emin_2E_3D\ (2^2))\ (\lambda V0x \in 2.V0x))\ (\lambda V1x \in 2.V1x))$

Definition 5 We define $c_2Ebool_2E_21$ to be $\lambda A.\lambda 27a : \iota.(\lambda V0P \in (2^{A-27a}).(ap\ (ap\ (c_2Emin_2E_3D\ (2^{A-27a}))\ (\lambda V1x \in 2.V1x))\ (\lambda V2x \in 2.V2x)))$

Definition 6 We define c_2Enum_2ESUC to be $\lambda V0m \in ty_2Enum_2Enum.(ap\ c_2Enum_2EABS_num\ (c_2Enum_2ESUC_REP\ m))$

Let $c_2Earithmetic_2E_2B : \iota$ be given. Assume the following.

$$c_2Earithmetic_2E_2B \in ((ty_2Enum_2Enum^{ty_2Enum_2Enum})^{ty_2Enum_2Enum}) \tag{6}$$

Definition 7 We define $c_2Earithmic_2EBIT1$ to be $\lambda V0n \in ty_2Enum_2Enum.(ap (ap c_2Earithmic_2EBIT1))$

Definition 8 We define $c_2Earithmic_2ENUMERAL$ to be $\lambda V0x \in ty_2Enum_2Enum.V0x$.

Let $ty_2Erealax_2Ereal : \iota$ be given. Assume the following.

$$nonempty\ ty_2Erealax_2Ereal \quad (7)$$

Let $c_2Ereal_2Ereal_of_num : \iota$ be given. Assume the following.

$$c_2Ereal_2Ereal_of_num \in (ty_2Erealax_2Ereal^{ty_2Enum_2Enum}) \quad (8)$$

Definition 9 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow p Q)$ of type ι .

Definition 10 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2)) (\lambda V2t \in 2.t)))$

Let $ty_2Epair_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty_2Epair_2Eprod\ A0\ A1) \quad (9)$$

Let $c_2Epair_2EABS_prod : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow \forall A_27b.nonempty\ A_27b \Rightarrow c_2Epair_2EABS_prod\ A_27a\ A_27b \in ((ty_2Epair_2Eprod\ A_27a\ A_27b)^{(2^{A_27b})^{A_27a}}) \quad (10)$$

Definition 11 We define $c_2Epair_2E_2C$ to be $\lambda A_27a : \iota.\lambda A_27b : \iota.\lambda V0x \in A_27a.\lambda V1y \in A_27b.(ap (c_2Epair_2E_2C))$

Definition 12 We define $c_2Ehrat_2Etrat_1$ to be $(ap (ap (c_2Epair_2E_2C\ ty_2Enum_2Enum\ ty_2Enum_2Enum)))$

Let $c_2Ehrat_2Etrat_eq : \iota$ be given. Assume the following.

$$c_2Ehrat_2Etrat_eq \in ((2^{(ty_2Epair_2Eprod\ ty_2Enum_2Enum\ ty_2Enum_2Enum)})^{(ty_2Epair_2Eprod\ ty_2Enum_2Enum)}) \quad (11)$$

Let $ty_2Ehrat_2Ehrat : \iota$ be given. Assume the following.

$$nonempty\ ty_2Ehrat_2Ehrat \quad (12)$$

Let $c_2Ehrat_2Ehrat_ABS_CLASS : \iota$ be given. Assume the following.

$$c_2Ehrat_2Ehrat_ABS_CLASS \in (ty_2Ehrat_2Ehrat^{(2^{(ty_2Epair_2Eprod\ ty_2Enum_2Enum\ ty_2Enum_2Enum)})}) \quad (13)$$

Definition 13 We define $c_2Ehrat_2Ehrat_ABS$ to be $\lambda V0r \in (ty_2Epair_2Eprod\ ty_2Enum_2Enum\ ty_2Enum_2Enum)$

Definition 14 We define $c_2Ehrat_2Ehrat_1$ to be $(ap\ c_2Ehrat_2Ehrat_ABS\ c_2Ehrat_2Etrat_1)$.

Let $c_2Eh_rat_2Eh_rat_REP_CLASS : \iota$ be given. Assume the following.

$$c_2Eh_rat_2Eh_rat_REP_CLASS \in ((2^{(ty_2Epair_2Eprod\ ty_2Enum_2Enum\ ty_2Enum_2Enum)})^{ty_2Eh_rat_2Eh_rat}) \quad (14)$$

Definition 15 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A$.if $(\exists x \in A.p (ap\ P\ x))$ then (the $(\lambda x.x \in A \wedge p\ x)$ of type $\iota \Rightarrow \iota$).

Definition 16 We define $c_2Eh_rat_2Eh_rat_REP$ to be $\lambda V0a \in ty_2Eh_rat_2Eh_rat$.(ap $(c_2Emin_2E_40\ (ty_2Eh_rat_2Eh_rat\ a))$).

Let $c_2Eh_rat_2Etr_at_add : \iota$ be given. Assume the following.

$$c_2Eh_rat_2Etr_at_add \in (((ty_2Epair_2Eprod\ ty_2Enum_2Enum\ ty_2Enum_2Enum)^{(ty_2Epair_2Eprod\ ty_2Enum_2Enum)})^{(ty_2Eh_rat_2Eh_rat)}} \quad (15)$$

Definition 17 We define $c_2Eh_rat_2Eh_rat_add$ to be $\lambda V0T1 \in ty_2Eh_rat_2Eh_rat$. $\lambda V1T2 \in ty_2Eh_rat_2Eh_rat$.

Definition 18 We define $c_2Ebool_2E_3F$ to be $\lambda A_27a : \iota$.($\lambda V0P \in (2^{A_27a})$).(ap $V0P$ (ap $(c_2Emin_2E_40\ (ty_2Eh_rat_2Eh_rat\ P))$)).

Definition 19 We define $c_2Ehreal_2Eh_rat_lt$ to be $\lambda V0x \in ty_2Eh_rat_2Eh_rat$. $\lambda V1y \in ty_2Eh_rat_2Eh_rat$.

Definition 20 We define $c_2Ehreal_2Ecut_of_h_rat$ to be $\lambda V0x \in ty_2Eh_rat_2Eh_rat$.($\lambda V1y \in ty_2Eh_rat_2Eh_rat$).

Let $ty_2Ehreal_2Ehreal : \iota$ be given. Assume the following.

$$nonempty\ ty_2Ehreal_2Ehreal \quad (16)$$

Let $c_2Ehreal_2Ehreal : \iota$ be given. Assume the following.

$$c_2Ehreal_2Ehreal \in (ty_2Ehreal_2Ehreal^{(2^{ty_2Eh_rat_2Eh_rat})}) \quad (17)$$

Definition 21 We define $c_2Ehreal_2Ehreal_1$ to be (ap $c_2Ehreal_2Ehreal$ (ap $c_2Ehreal_2Ecut_of_h_rat$)).

Definition 22 We define $c_2Erealax_2Etre_al_0$ to be (ap (ap $(c_2Epair_2E_2C\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)$)).

Let $c_2Erealax_2Etre_al_eq : \iota$ be given. Assume the following.

$$c_2Erealax_2Etre_al_eq \in ((2^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)})^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal)}) \quad (18)$$

Let $c_2Erealax_2Ereal_ABS_CLASS : \iota$ be given. Assume the following.

$$c_2Erealax_2Ereal_ABS_CLASS \in (ty_2Erealax_2Ereal^{(2^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)})}) \quad (19)$$

Definition 23 We define $c_2Erealax_2Ereal_ABS$ to be $\lambda V0r \in (ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)$.

Definition 24 We define $c_2Erealax_2Ereal_0$ to be (ap $c_2Erealax_2Ereal_ABS$ $c_2Erealax_2Etre_al_0$).

Let $c_2Ehreal_2Ecut : \iota$ be given. Assume the following.

$$c_2Ehreal_2Ecut \in ((2^{ty_2Eh_rat_2Eh_rat})^{ty_2Ehreal_2Ehreal}) \quad (20)$$

Definition 25 We define $c_Ehreal_Ehreal_add$ to be $\lambda V0X \in ty_Ehreal_Ehreal.\lambda V1Y \in ty_Ehreal_Ehreal.$

Definition 26 We define $c_Erealax_Etreax_1$ to be $(ap (ap (c_Epair_E_2C ty_Ehreal_Ehreal ty_Ehreal_Ehreal$

Definition 27 We define $c_Erealax_Ereal_1$ to be $(ap c_Erealax_Ereal_ABS c_Erealax_Etreax_1)$.

Definition 28 We define c_Ebool_EF to be $(ap (c_Ebool_E_21 2) (\lambda V0t \in 2.V0t))$.

Definition 29 We define $c_Ebool_E_7E$ to be $(\lambda V0t \in 2.(ap (ap c_Emin_E_3D_3D_3E V0t) c_Ebool_E_21 2))$.

Assume the following.

$$(c_Erealax_Ereal_0 = (ap c_Ereal_Ereal_of_num c_Eenum_E0)) \quad (21)$$

Assume the following.

$$(c_Erealax_Ereal_1 = (ap c_Ereal_Ereal_of_num (ap c_Earithmic_ENUMERAL (ap c_Earithmic_EBIT1 c_Earithmic_EZERO)))) \quad (22)$$

Assume the following.

$$(\neg(c_Erealax_Ereal_1 = c_Erealax_Ereal_0)) \quad (23)$$

Theorem 1

$$(\neg((ap c_Ereal_Ereal_of_num (ap c_Earithmic_ENUMERAL (ap c_Earithmic_EBIT1 c_Earithmic_EZERO))) = (ap c_Ereal_Ereal_of_num c_Eenum_E0)))$$