

thm\_2Ereal\_2EREAL\_ABS\_POS  
(TMav2H2H2ttMjY1YoYwhGCxmxLbRmwF1rUB)

October 26, 2020

Let  $ty\_2Ehreal\_2Ehreal : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Ehreal\_2Ehreal \tag{1}$$

Let  $ty\_2Epair\_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty\_2Epair\_2Eprod\ A0\ A1) \tag{2}$$

Let  $ty\_2Erealax\_2Ereal : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Erealax\_2Ereal \tag{3}$$

Let  $c\_2Erealax\_2Ereal\_REP\_CLASS : \iota$  be given. Assume the following.

$$c\_2Erealax\_2Ereal\_REP\_CLASS \in ((2^{(ty\_2Epair\_2Eprod\ ty\_2Ehreal\_2Ehreal\ ty\_2Ehreal\_2Ehreal)})^{ty\_2Erealax\_2Ereal}) \tag{4}$$

**Definition 1** We define  $c\_2Emin\_2E\_40$  to be  $\lambda A.\lambda P \in 2^A.$ if  $(\exists x \in A.p\ (ap\ P\ x))$  **then** (the  $(\lambda x.x \in A \wedge p)$  of type  $\iota \Rightarrow \iota$ ).

**Definition 2** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o\ (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 3** We define  $c\_2Ebool\_2E\_21$  to be  $(ap\ (ap\ (c\_2Emin\_2E\_3D\ (2^2))\ (\lambda V0x \in 2.V0x))\ (\lambda V1x \in 2.V1x))$

**Definition 4** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A.27a : \iota.(\lambda V0P \in (2^{A-27a}).(ap\ (ap\ (c\_2Emin\_2E\_3D\ (2^{A-27a})))$

**Definition 5** We define  $c\_2Erealax\_2Ereal\_REP$  to be  $\lambda V0a \in ty\_2Erealax\_2Ereal.(ap\ (c\_2Emin\_2E\_40\ (ty\_2Erealax\_2Ereal\_REP\_CLASS\ a)))$

Let  $c\_2Erealax\_2Etrealm\_neg : \iota$  be given. Assume the following.

$$c\_2Erealax\_2Etrealm\_neg \in ((ty\_2Epair\_2Eprod\ ty\_2Ehreal\_2Ehreal\ ty\_2Ehreal\_2Ehreal)^{(ty\_2Epair\_2Eprod\ ty\_2Ehreal\_2Ehreal\ ty\_2Ehreal\_2Ehreal)}) \tag{5}$$

Let  $c\_2Erealax\_2Etreall\_eq : \iota$  be given. Assume the following.

$$c\_2Erealax\_2Etreall\_eq \in ((2^{(ty\_2Epair\_2Eprod\ ty\_2Ehreal\_2Ehreal\ ty\_2Ehreal\_2Ehreal)}) (ty\_2Epair\_2Eprod\ ty\_2Ehreal\_2Ehreal)) \quad (6)$$

Let  $c\_2Erealax\_2Ereal\_ABS\_CLASS : \iota$  be given. Assume the following.

$$c\_2Erealax\_2Ereal\_ABS\_CLASS \in (ty\_2Erealax\_2Ereal)^{(2^{(ty\_2Epair\_2Eprod\ ty\_2Ehreal\_2Ehreal\ ty\_2Ehreal\_2Ehreal)})} \quad (7)$$

**Definition 6** We define  $c\_2Erealax\_2Ereal\_ABS$  to be  $\lambda V0r \in (ty\_2Epair\_2Eprod\ ty\_2Ehreal\_2Ehreal\ ty\_2Ehreal\_2Ehreal)$

**Definition 7** We define  $c\_2Erealax\_2Ereal\_neg$  to be  $\lambda V0T1 \in ty\_2Erealax\_2Ereal.(ap\ c\_2Erealax\_2Ereal\_ABS\ T1)$

Let  $c\_2Enum\_2EZERO\_REP : \iota$  be given. Assume the following.

$$c\_2Enum\_2EZERO\_REP \in \omega \quad (8)$$

Let  $ty\_2Enum\_2Enum : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Enum\_2Enum \quad (9)$$

Let  $c\_2Enum\_2EABS\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EABS\_num \in (ty\_2Enum\_2Enum^{\omega}) \quad (10)$$

**Definition 8** We define  $c\_2Enum\_2E0$  to be  $(ap\ c\_2Enum\_2EABS\_num\ c\_2Enum\_2EZERO\_REP)$ .

Let  $c\_2Ereal\_2Ereal\_of\_num : \iota$  be given. Assume the following.

$$c\_2Ereal\_2Ereal\_of\_num \in (ty\_2Erealax\_2Ereal)^{ty\_2Enum\_2Enum} \quad (11)$$

Let  $c\_2Erealax\_2Etreall\_lt : \iota$  be given. Assume the following.

$$c\_2Erealax\_2Etreall\_lt \in ((2^{(ty\_2Epair\_2Eprod\ ty\_2Ehreal\_2Ehreal\ ty\_2Ehreal\_2Ehreal)}) (ty\_2Epair\_2Eprod\ ty\_2Ehreal\_2Ehreal)) \quad (12)$$

**Definition 9** We define  $c\_2Erealax\_2Ereal\_lt$  to be  $\lambda V0T1 \in ty\_2Erealax\_2Ereal.\lambda V1T2 \in ty\_2Erealax\_2Ereal$

**Definition 10** We define  $c\_2Ebool\_2EF$  to be  $(ap\ (c\_2Ebool\_2E21\ 2)\ (\lambda V0t \in 2.V0t))$ .

**Definition 11** We define  $c\_2Emin\_2E3D\_3D\_3E$  to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o\ (p\ P \Rightarrow p\ Q)$  of type  $\iota$ .

**Definition 12** We define  $c\_2Ebool\_2E7E$  to be  $(\lambda V0t \in 2.(ap\ (ap\ c\_2Emin\_2E3D\_3D\_3E\ V0t)\ c\_2Ebool\_2E21))$

**Definition 13** We define  $c\_2Ereal\_2Ereal\_lte$  to be  $\lambda V0x \in ty\_2Erealax\_2Ereal.\lambda V1y \in ty\_2Erealax\_2Ereal$

**Definition 14** We define  $c\_2Ebool\_2E2F\_5C$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap\ (c\_2Ebool\_2E21\ 2)\ (\lambda V2t \in 2.V2t))))$

**Definition 15** We define  $c\_2Ebool\_2ECOND$  to be  $\lambda A\_27a : \iota.(\lambda V0t \in 2.(\lambda V1t1 \in A\_27a.(\lambda V2t2 \in A\_27a.(A\_27a\ V2t2))))$

**Definition 16** We define `c_2Ereal_2Eabs` to be  $\lambda V0x \in ty\_2Erealax\_2Ereal.(ap (ap (ap (c\_2Ebool\_2ECON$

Assume the following.

$$\begin{aligned}
 & (\forall V0x \in ty\_2Erealax\_2Ereal.(p (ap (ap c\_2Ereal\_2Ereal\_lte \\
 & (ap c\_2Ereal\_2Ereal\_of\_num c\_2Enum\_2E0)) (ap c\_2Ereal\_2Eabs \\
 & V0x)))) \quad (13)
 \end{aligned}$$

**Theorem 1**

$$\begin{aligned}
 & (\forall V0x \in ty\_2Erealax\_2Ereal.(p (ap (ap c\_2Ereal\_2Ereal\_lte \\
 & (ap c\_2Ereal\_2Ereal\_of\_num c\_2Enum\_2E0)) (ap c\_2Ereal\_2Eabs \\
 & V0x))))
 \end{aligned}$$