

# thm\_2Ereal\_2EREAL\_\_DIV\_\_LMUL (TMW- MUsDL1WZu7zhDap85pBzFoQ8BGWpzQQQ)

October 26, 2020

**Definition 1** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define  $c\_2Ebool\_2E\_2T$  to be  $(ap (ap (c\_2Emin\_2E\_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Let  $c\_2Enum\_2E\_ZERO\_REP : \iota$  be given. Assume the following.

$$c\_2Enum\_2E\_ZERO\_REP \in \omega \tag{1}$$

Let  $ty\_2Enum\_2Enum : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Enum\_2Enum \tag{2}$$

Let  $c\_2Enum\_2E\_ABS\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2E\_ABS\_num \in (ty\_2Enum\_2Enum^{\omega}) \tag{3}$$

**Definition 3** We define  $c\_2Enum\_2E\_0$  to be  $(ap\ c\_2Enum\_2E\_ABS\_num\ c\_2Enum\_2E\_ZERO\_REP)$ .

**Definition 4** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A.27a : \iota.(\lambda V0P \in (2^{A-27a}).(ap (ap (c\_2Emin\_2E\_3D (2^{A-27a}))$

**Definition 5** We define  $c\_2Ebool\_2E\_2F$  to be  $(ap (c\_2Ebool\_2E\_21\ 2) (\lambda V0t \in 2.V0t))$ .

**Definition 6** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o (p\ P \Rightarrow p\ Q)$  of type  $\iota$ .

**Definition 7** We define  $c\_2Ebool\_2E\_27E$  to be  $(\lambda V0t \in 2.(ap (ap\ c\_2Emin\_2E\_3D\_3D\_3E\ V0t)\ c\_2Ebool\_2E\_2F))$

Let  $ty\_2Ehreal\_2Ehreal : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Ehreal\_2Ehreal \tag{4}$$

Let  $ty\_2Epair\_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty\_2Epair\_2Eprod\ A0\ A1) \tag{5}$$

Let  $ty\_2Erealax\_2Ereal : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Erealax\_2Ereal \quad (6)$$

Let  $c\_2Erealax\_2Ereal\_REP\_CLASS : \iota$  be given. Assume the following.

$$c\_2Erealax\_2Ereal\_REP\_CLASS \in ((2^{(ty\_2Epair\_2Eprod\ ty\_2Ehreal\_2Ehreal\ ty\_2Ehreal\_2Ehreal)})^{ty\_2Erealax\_2Ereal}) \quad (7)$$

**Definition 8** We define  $c\_2Emin\_2E\_40$  to be  $\lambda A.\lambda P \in 2^A.$ if  $(\exists x \in A.p (ap\ P\ x))$  then (the  $(\lambda x.x \in A \wedge p$  of type  $\iota \Rightarrow \iota$ ).

**Definition 9** We define  $c\_2Erealax\_2Ereal\_REP$  to be  $\lambda V0a \in ty\_2Erealax\_2Ereal.(ap\ (c\_2Emin\_2E\_40\ (ty$

Let  $c\_2Erealax\_2Etrealm\_inv : \iota$  be given. Assume the following.

$$c\_2Erealax\_2Etrealm\_inv \in ((ty\_2Epair\_2Eprod\ ty\_2Ehreal\_2Ehreal\ ty\_2Ehreal\_2Ehreal)^{(ty\_2Epair\_2Eprod\ ty\_2Ehreal\_2Ehreal\ ty\_2Ehreal\_2Ehreal)}) \quad (8)$$

Let  $c\_2Erealax\_2Etrealm\_eq : \iota$  be given. Assume the following.

$$c\_2Erealax\_2Etrealm\_eq \in ((2^{(ty\_2Epair\_2Eprod\ ty\_2Ehreal\_2Ehreal\ ty\_2Ehreal\_2Ehreal)})^{(ty\_2Epair\_2Eprod\ ty\_2Ehreal\_2Ehreal\ ty\_2Ehreal\_2Ehreal)}) \quad (9)$$

Let  $c\_2Erealax\_2Ereal\_ABS\_CLASS : \iota$  be given. Assume the following.

$$c\_2Erealax\_2Ereal\_ABS\_CLASS \in (ty\_2Erealax\_2Ereal)^{(2^{(ty\_2Epair\_2Eprod\ ty\_2Ehreal\_2Ehreal\ ty\_2Ehreal\_2Ehreal)})} \quad (10)$$

**Definition 10** We define  $c\_2Erealax\_2Ereal\_ABS$  to be  $\lambda V0r \in (ty\_2Epair\_2Eprod\ ty\_2Ehreal\_2Ehreal\ ty$

**Definition 11** We define  $c\_2Erealax\_2Einv$  to be  $\lambda V0T1 \in ty\_2Erealax\_2Ereal.(ap\ c\_2Erealax\_2Ereal\_ABS$

Let  $c\_2Erealax\_2Etrealm\_mul : \iota$  be given. Assume the following.

$$c\_2Erealax\_2Etrealm\_mul \in (((ty\_2Epair\_2Eprod\ ty\_2Ehreal\_2Ehreal\ ty\_2Ehreal\_2Ehreal)^{(ty\_2Epair\_2Eprod\ ty\_2Ehreal\_2Ehreal\ ty\_2Ehreal\_2Ehreal)})^{(ty\_2Epair\_2Eprod\ ty\_2Ehreal\_2Ehreal\ ty\_2Ehreal\_2Ehreal)}) \quad (11)$$

**Definition 12** We define  $c\_2Erealax\_2Ereal\_mul$  to be  $\lambda V0T1 \in ty\_2Erealax\_2Ereal.\lambda V1T2 \in ty\_2Erealax$

**Definition 13** We define  $c\_2Ereal\_2E\_2F$  to be  $\lambda V0x \in ty\_2Erealax\_2Ereal.\lambda V1y \in ty\_2Erealax\_2Ereal.($

**Definition 14** We define  $c\_2Earithmic\_2EZERO$  to be  $c\_2Enum\_2E0$ .

Let  $c\_2Enum\_2EREP\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EREP\_num \in (\omega^{ty\_2Enum\_2Enum}) \quad (12)$$

Let  $c\_2Enum\_2ESUC\_REP : \iota$  be given. Assume the following.

$$c\_2Enum\_2ESUC\_REP \in (\omega^{\omega}) \quad (13)$$

**Definition 15** We define  $c\_2Enum\_2ESUC$  to be  $\lambda V0m \in ty\_2Enum\_2Enum.(ap\ c\_2Enum\_2EABS\_num$   
Let  $c\_2Earithmetic\_2E\_2B : \iota$  be given. Assume the following.

$$c\_2Earithmetic\_2E\_2B \in ((ty\_2Enum\_2Enum^{ty\_2Enum\_2Enum})^{ty\_2Enum\_2Enum}) \quad (14)$$

**Definition 16** We define  $c\_2Earithmetic\_2EBIT1$  to be  $\lambda V0n \in ty\_2Enum\_2Enum.(ap\ (ap\ c\_2Earithmetic$

**Definition 17** We define  $c\_2Earithmetic\_2ENUMERAL$  to be  $\lambda V0x \in ty\_2Enum\_2Enum.V0x$ .

Let  $c\_2Ereal\_2Ereal\_of\_num : \iota$  be given. Assume the following.

$$c\_2Ereal\_2Ereal\_of\_num \in (ty\_2Erealax\_2Ereal^{ty\_2Enum\_2Enum}) \quad (15)$$

Assume the following.

$$True \quad (16)$$

Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow (\forall V0x \in A\_27a.((V0x = V0x) \Leftrightarrow True)) \quad (17)$$

Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow (\forall V0x \in A\_27a.(\forall V1y \in A\_27a.((V0x = V1y) \Leftrightarrow (V1y = V0x)))) \quad (18)$$

Assume the following.

$$(\forall V0x \in ty\_2Erealax\_2Ereal.(\forall V1y \in ty\_2Erealax\_2Ereal. ((ap\ (ap\ c\_2Erealax\_2Ereal\_mul\ V0x)\ V1y) = (ap\ (ap\ c\_2Erealax\_2Ereal\_mul\ V1y)\ V0x)))) \quad (19)$$

Assume the following.

$$(\forall V0x \in ty\_2Erealax\_2Ereal.(\forall V1y \in ty\_2Erealax\_2Ereal. (\forall V2z \in ty\_2Erealax\_2Ereal.((ap\ (ap\ c\_2Erealax\_2Ereal\_mul\ V0x)\ (ap\ (ap\ c\_2Erealax\_2Ereal\_mul\ V1y)\ V2z)) = (ap\ (ap\ c\_2Erealax\_2Ereal\_mul\ (ap\ (ap\ c\_2Erealax\_2Ereal\_mul\ V0x)\ V1y))\ V2z)))))) \quad (20)$$

Assume the following.

$$(\forall V0x \in ty\_2Erealax\_2Ereal.((\neg(V0x = (ap\ c\_2Ereal\_2Ereal\_of\_num\ c\_2Enum\_2E0))) \Rightarrow ((ap\ (ap\ c\_2Erealax\_2Ereal\_mul\ (ap\ c\_2Erealax\_2Einv\ V0x))\ V0x) = (ap\ c\_2Ereal\_2Ereal\_of\_num\ (ap\ c\_2Earithmetic\_2ENUMERAL\ (ap\ c\_2Earithmetic\_2EBIT1\ c\_2Earithmetic\_2EZERO)))))) \quad (21)$$

Assume the following.

$$(\forall V0x \in ty\_2Erealax\_2Ereal.((ap\ (ap\ c\_2Erealax\_2Ereal\_mul\ V0x)\ (ap\ c\_2Ereal\_2Ereal\_of\_num\ (ap\ c\_2Earithmetic\_2ENUMERAL\ (ap\ c\_2Earithmetic\_2EBIT1\ c\_2Earithmetic\_2EZERO)))) = V0x)) \quad (22)$$

**Theorem 1**

$$\begin{aligned} & (\forall V0x \in \text{ty\_2Erealax\_2Ereal}.\forall V1y \in \text{ty\_2Erealax\_2Ereal}. \\ & ((\neg(V1y = (\text{ap } c\_2Ereal\_2Ereal\_of\_num \ c\_2Enum\_2E0)))) \Rightarrow ((\text{ap } ( \\ & \text{ap } c\_2Erealax\_2Ereal\_mul \ V1y) (\text{ap } (\text{ap } c\_2Ereal\_2E\_2F \ V0x) \ V1y)) = \\ & \quad V0x)))) \end{aligned}$$