

thm_2Ereal_2EREAL__DIV__LZERO (TMdViaHRdBq2Fcc35zz8mBsCUgQpeg78oWz)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define $c_2Ebool_2E_21$ to be $(ap (ap (c_2Emin_2E_3D (2^2))) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Let $ty_2Ehreal_2Ehreal : \iota$ be given. Assume the following.

$$nonempty\ ty_2Ehreal_2Ehreal \tag{1}$$

Let $ty_2Epair_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty_2Epair_2Eprod\ A0\ A1) \tag{2}$$

Let $ty_2Erealax_2Ereal : \iota$ be given. Assume the following.

$$nonempty\ ty_2Erealax_2Ereal \tag{3}$$

Let $c_2Erealax_2Ereal_REP_CLASS : \iota$ be given. Assume the following.

$$c_2Erealax_2Ereal_REP_CLASS \in ((2^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)})^{ty_2Erealax_2Ereal}) \tag{4}$$

Definition 3 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A.if (\exists x \in A.p (ap\ P\ x))$ then $(the (\lambda x.x \in A \wedge p (ap\ P\ x)))$ of type $\iota \Rightarrow \iota$.

Definition 4 We define $c_2Ebool_2E_21$ to be $\lambda A.27a : \iota.(\lambda V0P \in (2^{A-27a}).(ap (ap (c_2Emin_2E_3D (2^{A-27a}))) (\lambda V1P \in 2.V1P)))$

Definition 5 We define $c_2Erealax_2Ereal_REP$ to be $\lambda V0a \in ty_2Erealax_2Ereal.(ap (c_2Emin_2E_40 (ty_2Epair_2Eprod (ty_2Erealax_2Ereal (ap (c_2Emin_2E_3D (2^{2^a}))) (\lambda V1x \in 2.V1x)) (\lambda V2x \in 2.V2x))))$

Let $c_2Erealax_2Etreall_inv : \iota$ be given. Assume the following.

$$c_2Erealax_2Etreall_inv \in ((ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)}) \tag{5}$$

Let $c_2Erealax_2Etrealm_eq : \iota$ be given. Assume the following.

$$c_2Erealax_2Etrealm_eq \in ((2^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)})(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal)) \quad (6)$$

Let $c_2Erealax_2Ereal_ABS_CLASS : \iota$ be given. Assume the following.

$$c_2Erealax_2Ereal_ABS_CLASS \in (ty_2Erealax_2Ereal)^{(2^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)}} \quad (7)$$

Definition 6 We define $c_2Erealax_2Ereal_ABS$ to be $\lambda V0r \in (ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)$

Definition 7 We define $c_2Erealax_2Einv$ to be $\lambda V0T1 \in ty_2Erealax_2Ereal.(ap\ c_2Erealax_2Ereal_ABS)$

Let $c_2Erealax_2Etrealm_mul : \iota$ be given. Assume the following.

$$c_2Erealax_2Etrealm_mul \in (((ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal))^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)}) \quad (8)$$

Definition 8 We define $c_2Erealax_2Ereal_mul$ to be $\lambda V0T1 \in ty_2Erealax_2Ereal.\lambda V1T2 \in ty_2Erealax_2Ereal$

Definition 9 We define c_2Ereal_2E2F to be $\lambda V0x \in ty_2Erealax_2Ereal.\lambda V1y \in ty_2Erealax_2Ereal.(a$

Let $c_2Enum_2EZERO_REP : \iota$ be given. Assume the following.

$$c_2Enum_2EZERO_REP \in \omega \quad (9)$$

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \quad (10)$$

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum)^{\omega} \quad (11)$$

Definition 10 We define c_2Enum_2E0 to be $(ap\ c_2Enum_2EABS_num\ c_2Enum_2EZERO_REP)$.

Let $c_2Ereal_2Ereal_of_num : \iota$ be given. Assume the following.

$$c_2Ereal_2Ereal_of_num \in (ty_2Erealax_2Ereal)^{ty_2Enum_2Enum} \quad (12)$$

Assume the following.

$$True \quad (13)$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0x \in A_27a.((V0x = V0x) \Leftrightarrow True)) \quad (14)$$

Assume the following.

$$\begin{aligned} & (\forall V0x \in ty_2Erealax_2Ereal.((ap\ (ap\ c_2Erealax_2Ereal_mul \\ & (ap\ c_2Ereal_2Ereal_of_num\ c_2Enum_2E0))\ V0x) = (ap\ c_2Ereal_2Ereal_of_num \\ & c_2Enum_2E0))) \end{aligned} \quad (15)$$

Theorem 1

$(\forall V0x \in ty_2Erealax_2Ereal.((ap (ap c_2Ereal_2E_2F (ap$
 $c_2Ereal_2Ereal_of_num c_2Enum_2E0)) V0x) = (ap c_2Ereal_2Ereal_of_num$
 $c_2Enum_2E0)))$