

thm\_2Ereal\_2EREAL\_\_DIV\_\_REFL  
(TMQ7BGoyxMTViWXuuPbCqZSo4NZ1NNCESTG)

October 26, 2020

**Definition 1** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define  $c\_2Ebool\_2E\_2T$  to be  $(ap (ap (c\_2Emin\_2E\_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

**Definition 3** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A.^{27a} : \iota.(\lambda V0P \in (2^{A-27a}).(ap (ap (c\_2Emin\_2E\_3D (2^{A-27a}))$

**Definition 4** We define  $c\_2Ebool\_2E\_2F$  to be  $(ap (c\_2Ebool\_2E\_21 2) (\lambda V0t \in 2.V0t))$ .

**Definition 5** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o (p P \Rightarrow p Q)$  of type  $\iota$ .

**Definition 6** We define  $c\_2Ebool\_2E\_2F\_5C$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_2Ebool\_2E\_21 2) (\lambda V2t \in 2.V2t)))$

Let  $ty\_2Ehreal\_2Ehreal : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Ehreal\_2Ehreal \tag{1}$$

Let  $ty\_2Epair\_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty\_2Epair\_2Eprod\ A0\ A1) \tag{2}$$

Let  $ty\_2Erealax\_2Ereal : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Erealax\_2Ereal \tag{3}$$

Let  $c\_2Erealax\_2Ereal\_REP\_CLASS : \iota$  be given. Assume the following.

$$c\_2Erealax\_2Ereal\_REP\_CLASS \in ((2^{(ty\_2Epair\_2Eprod\ ty\_2Ehreal\_2Ehreal\ ty\_2Ehreal\_2Ehreal)})\ ty\_2Erealax\_2Ereal) \tag{4}$$

**Definition 7** We define  $c\_2Emin\_2E\_40$  to be  $\lambda A.\lambda P \in 2^A.if (\exists x \in A.p (ap P x)) \mathbf{then} (the (\lambda x.x \in A \wedge p x))$  of type  $\iota \Rightarrow \iota$ .

**Definition 8** We define  $c\_2Erealax\_2Ereal\_REP$  to be  $\lambda V0a \in ty\_2Erealax\_2Ereal.(ap (c\_2Emin\_2E\_40 (ty\_2Erealax\_2Ereal\_REP\_CLASS a)))$



**Definition 15** We define  $c\_2Enum\_2ESUC$  to be  $\lambda V0m \in ty\_2Enum\_2Enum.(ap\ c\_2Enum\_2EABS\_num$

Let  $c\_2Earithmetic\_2E\_2B : \iota$  be given. Assume the following.

$$c\_2Earithmetic\_2E\_2B \in ((ty\_2Enum\_2Enum^{ty\_2Enum\_2Enum})^{ty\_2Enum\_2Enum}) \quad (14)$$

**Definition 16** We define  $c\_2Earithmetic\_2EBIT1$  to be  $\lambda V0n \in ty\_2Enum\_2Enum.(ap\ (ap\ c\_2Earithmetic$

**Definition 17** We define  $c\_2Earithmetic\_2ENUMERAL$  to be  $\lambda V0x \in ty\_2Enum\_2Enum.V0x$ .

Let  $c\_2Ereal\_2Ereal\_of\_num : \iota$  be given. Assume the following.

$$c\_2Ereal\_2Ereal\_of\_num \in (ty\_2Erealax\_2Ereal^{ty\_2Enum\_2Enum}) \quad (15)$$

**Definition 18** We define  $c\_2Ebool\_2E\_7E$  to be  $(\lambda V0t \in 2.(ap\ (ap\ c\_2Emin\_2E\_3D\_3D\_3E\ V0t)\ c\_2Ebool\_2E$

Assume the following.

$$True \quad (16)$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2.(((True \Leftrightarrow (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \Leftrightarrow True) \Leftrightarrow \\ & (p\ V0t)) \wedge (((False \Leftrightarrow (p\ V0t)) \Leftrightarrow \neg(p\ V0t)) \wedge (((p\ V0t) \Leftrightarrow False) \Leftrightarrow \neg( \\ & p\ V0t)))))) \end{aligned} \quad (17)$$

Assume the following.

$$\begin{aligned} & (\forall V0x \in ty\_2Erealax\_2Ereal.(\neg(V0x = (ap\ c\_2Ereal\_2Ereal\_of\_num \\ & c\_2Enum\_2E0))) \Rightarrow ((ap\ (ap\ c\_2Erealax\_2Ereal\_mul\ V0x)\ (ap\ c\_2Erealax\_2Einv \\ & V0x)) = (ap\ c\_2Ereal\_2Ereal\_of\_num\ (ap\ c\_2Earithmetic\_2ENUMERAL \\ & (ap\ c\_2Earithmetic\_2EBIT1\ c\_2Earithmetic\_2EZERO)))))) \end{aligned} \quad (18)$$

**Theorem 1**

$$\begin{aligned} & (\forall V0x \in ty\_2Erealax\_2Ereal.(\neg(V0x = (ap\ c\_2Ereal\_2Ereal\_of\_num \\ & c\_2Enum\_2E0))) \Rightarrow ((ap\ (ap\ c\_2Ereal\_2E\_2F\ V0x)\ V0x) = (ap\ c\_2Ereal\_2Ereal\_of\_num \\ & (ap\ c\_2Earithmetic\_2ENUMERAL\ (ap\ c\_2Earithmetic\_2EBIT1\ c\_2Earithmetic\_2EZERO)))))) \end{aligned}$$