

thm\_2Ereal\_2EREAL\_DIV\_RMUL  
(TMQP8L4kYtgnVCQTWU7tdpkTR5YdEEwEMjy)

October 26, 2020

Let  $ty\_2Ehreal\_2Ehreal : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Ehreal\_2Ehreal \tag{1}$$

Let  $ty\_2Epair\_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty\_2Epair\_2Eprod\ A0\ A1) \tag{2}$$

Let  $ty\_2Erealax\_2Ereal : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Erealax\_2Ereal \tag{3}$$

Let  $c\_2Erealax\_2Ereal\_REP\_CLASS : \iota$  be given. Assume the following.

$$c\_2Erealax\_2Ereal\_REP\_CLASS \in ((2^{(ty\_2Epair\_2Eprod\ ty\_2Ehreal\_2Ehreal\ ty\_2Ehreal\_2Ehreal)})^{ty\_2Erealax\_2Ereal}) \tag{4}$$

**Definition 1** We define  $c\_2Emin\_2E40$  to be  $\lambda A.\lambda P \in 2^A.$ if  $(\exists x \in A.p\ (ap\ P\ x))$  then (the  $(\lambda x.x \in A \wedge p)$  of type  $\iota \Rightarrow \iota$ ).

**Definition 2** We define  $c\_2Emin\_2E3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o\ (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 3** We define  $c\_2Ebool\_2E27$  to be  $(ap\ (ap\ (c\_2Emin\_2E3D\ (2^2))\ (\lambda V0x \in 2.V0x))\ (\lambda V1x \in 2.V1x))$

**Definition 4** We define  $c\_2Ebool\_2E21$  to be  $\lambda A.27a : \iota.(\lambda V0P \in (2^{A-27a}).(ap\ (ap\ (c\_2Emin\_2E3D\ (2^{A-27a})))$

**Definition 5** We define  $c\_2Erealax\_2Ereal\_REP$  to be  $\lambda V0a \in ty\_2Erealax\_2Ereal.(ap\ (c\_2Emin\_2E40\ (ty\_2Erealax\_2Ereal\ V0a)))$

Let  $c\_2Erealax\_2Etreall\_inv : \iota$  be given. Assume the following.

$$c\_2Erealax\_2Etreall\_inv \in ((ty\_2Epair\_2Eprod\ ty\_2Ehreal\_2Ehreal\ ty\_2Ehreal\_2Ehreal)^{(ty\_2Epair\_2Eprod\ ty\_2Ehreal\_2Ehreal\ ty\_2Ehreal\_2Ehreal)}) \tag{5}$$

Let  $c\_2Erealax\_2Etrealm\_eq : \iota$  be given. Assume the following.

$$c\_2Erealax\_2Etrealm\_eq \in ((2^{(ty\_2Epair\_2Eprod\ ty\_2Ehreal\_2Ehreal\ ty\_2Ehreal\_2Ehreal)})(ty\_2Epair\_2Eprod\ ty\_2Ehreal\_2Ehreal)) \quad (6)$$

Let  $c\_2Erealax\_2Ereal\_ABS\_CLASS : \iota$  be given. Assume the following.

$$c\_2Erealax\_2Ereal\_ABS\_CLASS \in (ty\_2Erealax\_2Ereal)^{(2^{(ty\_2Epair\_2Eprod\ ty\_2Ehreal\_2Ehreal\ ty\_2Ehreal\_2Ehreal)}} \quad (7)$$

**Definition 6** We define  $c\_2Erealax\_2Ereal\_ABS$  to be  $\lambda V0r \in (ty\_2Epair\_2Eprod\ ty\_2Ehreal\_2Ehreal\ ty\_2Ehreal\_2Ehreal)$

**Definition 7** We define  $c\_2Erealax\_2Einv$  to be  $\lambda V0T1 \in ty\_2Erealax\_2Ereal.(ap\ c\_2Erealax\_2Ereal\_ABS)$

Let  $c\_2Erealax\_2Etrealm\_mul : \iota$  be given. Assume the following.

$$c\_2Erealax\_2Etrealm\_mul \in (((ty\_2Epair\_2Eprod\ ty\_2Ehreal\_2Ehreal\ ty\_2Ehreal\_2Ehreal)(ty\_2Epair\_2Eprod\ ty\_2Ehreal\_2Ehreal\ ty\_2Ehreal\_2Ehreal))^{(ty\_2Epair\_2Eprod\ ty\_2Ehreal\_2Ehreal\ ty\_2Ehreal\_2Ehreal)}) \quad (8)$$

**Definition 8** We define  $c\_2Erealax\_2Ereal\_mul$  to be  $\lambda V0T1 \in ty\_2Erealax\_2Ereal.\lambda V1T2 \in ty\_2Erealax\_2Ereal.$

**Definition 9** We define  $c\_2Ereal\_2E\_2F$  to be  $\lambda V0x \in ty\_2Erealax\_2Ereal.\lambda V1y \in ty\_2Erealax\_2Ereal.(a$

Let  $c\_2Enum\_2EZERO\_REP : \iota$  be given. Assume the following.

$$c\_2Enum\_2EZERO\_REP \in \omega \quad (9)$$

Let  $ty\_2Enum\_2Enum : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Enum\_2Enum \quad (10)$$

Let  $c\_2Enum\_2EABS\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EABS\_num \in (ty\_2Enum\_2Enum)^{\omega} \quad (11)$$

**Definition 10** We define  $c\_2Enum\_2E0$  to be  $(ap\ c\_2Enum\_2EABS\_num\ c\_2Enum\_2EZERO\_REP)$ .

Let  $c\_2Ereal\_2Ereal\_of\_num : \iota$  be given. Assume the following.

$$c\_2Ereal\_2Ereal\_of\_num \in (ty\_2Erealax\_2Ereal)^{(ty\_2Enum\_2Enum)} \quad (12)$$

**Definition 11** We define  $c\_2Ebool\_2EF$  to be  $(ap\ (c\_2Ebool\_2E\_21\ 2)\ (\lambda V0t \in 2.V0t))$ .

**Definition 12** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o\ (p\ P \Rightarrow p\ Q)$  of type  $\iota$ .

**Definition 13** We define  $c\_2Ebool\_2E\_7E$  to be  $(\lambda V0t \in 2.(ap\ (ap\ c\_2Emin\_2E\_3D\_3D\_3E\ V0t)\ c\_2Ebool\_2E\_21\ 2))$

Assume the following.

$$\begin{aligned}
& (\forall V0x \in ty\_2Erealax\_2Ereal. (\forall V1y \in ty\_2Erealax\_2Ereal. \\
& ((ap (ap c\_2Erealax\_2Ereal\_mul V0x) V1y) = (ap (ap c\_2Erealax\_2Ereal\_mul \\
& \quad V1y) V0x))))
\end{aligned} \tag{13}$$

Assume the following.

$$\begin{aligned}
& (\forall V0x \in ty\_2Erealax\_2Ereal. (\forall V1y \in ty\_2Erealax\_2Ereal. \\
& ((\neg (V1y = (ap c\_2Ereal\_2Ereal\_of\_num c\_2Enum\_2E0)))) \Rightarrow ((ap ( \\
& ap c\_2Erealax\_2Ereal\_mul V1y) (ap (ap c\_2Ereal\_2E\_2F V0x) V1y)) = \\
& \quad V0x))))
\end{aligned} \tag{14}$$

**Theorem 1**

$$\begin{aligned}
& (\forall V0x \in ty\_2Erealax\_2Ereal. (\forall V1y \in ty\_2Erealax\_2Ereal. \\
& ((\neg (V1y = (ap c\_2Ereal\_2Ereal\_of\_num c\_2Enum\_2E0)))) \Rightarrow ((ap ( \\
& ap c\_2Erealax\_2Ereal\_mul (ap (ap c\_2Ereal\_2E\_2F V0x) V1y)) V1y) = \\
& \quad V0x))))
\end{aligned}$$