

thm_2Ereal_2EREAL__EQ__LMUL2
(TMe18Jtx3hTD4M4dEYZBrB5tdDZaqiBmGrG)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define $c_2Ebool_2E_2T$ to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 3 We define $c_2Ebool_2E_21$ to be $\lambda A.\lambda a : \iota.(\lambda V0P \in (2^{A-27a}).(ap (ap (c_2Emin_2E_3D (2^{A-27a}))$

Definition 4 We define $c_2Ebool_2E_2F$ to be $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$.

Definition 5 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow p Q)$ of type ι .

Definition 6 We define $c_2Ebool_2E_27E$ to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2E_2F$

Definition 7 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2$

Let $c_2Enum_2EZERO_REP : \iota$ be given. Assume the following.

$$c_2Enum_2EZERO_REP \in \omega \tag{1}$$

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \tag{2}$$

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum^{\omega}) \tag{3}$$

Definition 8 We define c_2Enum_2E0 to be $(ap c_2Enum_2EABS_num c_2Enum_2EZERO_REP)$.

Let $ty_2Erealx_2Ereal : \iota$ be given. Assume the following.

$$nonempty\ ty_2Erealx_2Ereal \tag{4}$$

Let $c_2Ereal_2Ereal_of_num : \iota$ be given. Assume the following.

$$c_2Ereal_2Ereal_of_num \in (ty_2Erealx_2Ereal^{ty_2Enum_2Enum}) \tag{5}$$

Definition 9 We define $c_Ebool_2E_5C_2F$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_Ebool_2E_21) 2) (\lambda V2t \in 2)))$.
Let $ty_2Ehreal_2Ehreal : \iota$ be given. Assume the following.

$$nonempty\ ty_2Ehreal_2Ehreal \quad (6)$$

Let $ty_2Epair_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty_2Epair_2Eprod\ A0\ A1) \quad (7)$$

Let $c_2Erealax_2Ereal_REP_CLASS : \iota$ be given. Assume the following.

$$c_2Erealax_2Ereal_REP_CLASS \in ((2^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)})^{ty_2Erealax}) \quad (8)$$

Definition 10 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A.\text{if } (\exists x \in A.p (ap\ P\ x)) \text{ then } (the\ (\lambda x.x \in A \wedge P\ x))$ of type $\iota \Rightarrow \iota$.

Definition 11 We define $c_2Erealax_2Ereal_REP$ to be $\lambda V0a \in ty_2Erealax_2Ereal.(ap (c_2Emin_2E_40) (ty_2Erealax_2Ereal_REP_CLASS\ a))$.

Let $c_2Erealax_2Etrealmul : \iota$ be given. Assume the following.

$$c_2Erealax_2Etrealmul \in (((ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)})^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)}) \quad (9)$$

Let $c_2Erealax_2Etrealeq : \iota$ be given. Assume the following.

$$c_2Erealax_2Etrealeq \in ((2^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)})^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)}) \quad (10)$$

Let $c_2Erealax_2Ereal_ABS_CLASS : \iota$ be given. Assume the following.

$$c_2Erealax_2Ereal_ABS_CLASS \in (ty_2Erealax_2Ereal)^{(2^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)})} \quad (11)$$

Definition 12 We define $c_2Erealax_2Ereal_ABS$ to be $\lambda V0r \in (ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)$.

Definition 13 We define $c_2Erealax_2Ereal_mul$ to be $\lambda V0T1 \in ty_2Erealax_2Ereal.\lambda V1T2 \in ty_2Erealax_2Ereal$.

Assume the following.

$$(\forall V0t1 \in 2.(\forall V1t2 \in 2.(((p\ V0t1) \Rightarrow (p\ V1t2)) \Rightarrow (((p\ V1t2) \Rightarrow (p\ V0t1)) \Rightarrow ((p\ V0t1) \Leftrightarrow (p\ V1t2)))))) \quad (12)$$

Assume the following.

$$(\forall V0t \in 2.(False \Rightarrow (p\ V0t))) \quad (13)$$

Assume the following.

$$(\forall V0t \in 2.(((True \vee (p\ V0t)) \Leftrightarrow True) \wedge (((p\ V0t) \vee True) \Leftrightarrow True) \wedge (((False \vee (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \vee False) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \vee (p\ V0t)) \Leftrightarrow (p\ V0t)))))) \quad (14)$$

Assume the following.

$$\begin{aligned}
& (\forall V0x \in ty_2Erealax_2Ereal. (\forall V1y \in ty_2Erealax_2Ereal. \\
& (\forall V2z \in ty_2Erealax_2Ereal. (((ap (ap c_2Erealax_2Ereal_mul \\
V0x) V1y) = (ap (ap c_2Erealax_2Ereal_mul V0x) V2z)) \Leftrightarrow ((V0x = (ap \\
c_2Ereal_2Ereal_of_num c_2Enum_2E0)) \vee (V1y = V2z))))))
\end{aligned} \tag{15}$$

Theorem 1

$$\begin{aligned}
& (\forall V0x \in ty_2Erealax_2Ereal. (\forall V1y \in ty_2Erealax_2Ereal. \\
& (\forall V2z \in ty_2Erealax_2Ereal. ((\neg (V0x = (ap c_2Ereal_2Ereal_of_num \\
c_2Enum_2E0))) \Rightarrow ((V1y = V2z) \Leftrightarrow ((ap (ap c_2Erealax_2Ereal_mul \\
V0x) V1y) = (ap (ap c_2Erealax_2Ereal_mul V0x) V2z))))))
\end{aligned}$$