

thm_2Ereal_2EREAL_EQ_LMUL_IMP
(TMVKBcKVVQL-
WMXC8KeFkEPdTANCwz82ShTB)

October 26, 2020

Let $ty_2Ehreal_2Ehreal : \iota$ be given. Assume the following.

$$nonempty\ ty_2Ehreal_2Ehreal \tag{1}$$

Let $ty_2Epair_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty_2Epair_2Eprod\ A0\ A1) \tag{2}$$

Let $ty_2Erealax_2Ereal : \iota$ be given. Assume the following.

$$nonempty\ ty_2Erealax_2Ereal \tag{3}$$

Let $c_2Erealax_2Ereal_REP_CLASS : \iota$ be given. Assume the following.

$$c_2Erealax_2Ereal_REP_CLASS \in (((ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)\ ty_2Erealax_2Ereal) \tag{4}$$

Definition 1 We define c_2Emin_2E40 to be $\lambda A.\lambda P \in 2^A$.if $(\exists x \in A.p\ (ap\ P\ x))$ then (the $(\lambda x.x \in A \wedge p\ x)$ of type $\iota \Rightarrow \iota$).

Definition 2 We define c_2Emin_2E3D to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o\ (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 3 We define c_2Ebool_2E27 to be $(ap\ (ap\ (c_2Emin_2E3D\ (2^2))\ (\lambda V0x \in 2.V0x))\ (\lambda V1x \in 2.V1x))$

Definition 4 We define c_2Ebool_2E21 to be $\lambda A.\lambda a : \iota.(\lambda V0P \in (2^{A \cdot 27a}).(ap\ (ap\ (c_2Emin_2E3D\ (2^{A \cdot 27a}))\ (\lambda V1x \in 2.V1x))\ (\lambda V2x \in 2.V2x))$

Definition 5 We define $c_2Erealax_2Ereal_REP$ to be $\lambda V0a \in ty_2Erealax_2Ereal.(ap\ (c_2Emin_2E40\ (ty_2Erealax_2Ereal\ a))\ (ap\ (c_2Emin_2E3D\ (2^2))\ (\lambda V1x \in 2.V1x))\ (\lambda V2x \in 2.V2x))$

Let $c_2Erealax_2Etrealmul : \iota$ be given. Assume the following.

$$c_2Erealax_2Etrealmul \in (((ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)\ (ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)\ ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal) \tag{5}$$

Let $c_2Erealax_2Etrealm_eq : \iota$ be given. Assume the following.

$$c_2Erealax_2Etrealm_eq \in ((2^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)})^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal)}) \quad (6)$$

Let $c_2Erealax_2Ereal_ABS_CLASS : \iota$ be given. Assume the following.

$$c_2Erealax_2Ereal_ABS_CLASS \in (ty_2Erealax_2Ereal)^{(2^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)})} \quad (7)$$

Definition 6 We define $c_2Erealax_2Ereal_ABS$ to be $\lambda V0r \in (ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)$

Definition 7 We define $c_2Erealax_2Ereal_mul$ to be $\lambda V0T1 \in ty_2Erealax_2Ereal.\lambda V1T2 \in ty_2Erealax_2Ereal$

Let $c_2Enum_2EZERO_REP : \iota$ be given. Assume the following.

$$c_2Enum_2EZERO_REP \in \omega \quad (8)$$

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \quad (9)$$

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum)^{\omega} \quad (10)$$

Definition 8 We define c_2Enum_2E0 to be $(ap\ c_2Enum_2EABS_num\ c_2Enum_2EZERO_REP)$.

Let $c_2Ereal_2Ereal_of_num : \iota$ be given. Assume the following.

$$c_2Ereal_2Ereal_of_num \in (ty_2Erealax_2Ereal)^{ty_2Enum_2Enum} \quad (11)$$

Definition 9 We define c_2Ebool_2EF to be $(ap\ (c_2Ebool_2E_21\ 2)\ (\lambda V0t \in 2.V0t))$.

Definition 10 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o\ (p\ P \Rightarrow p\ Q)$ of type ι .

Definition 11 We define $c_2Ebool_2E_7E$ to be $(\lambda V0t \in 2.(ap\ (ap\ c_2Emin_2E_3D_3D_3E\ V0t)\ c_2Ebool_2E_21\ 2))$

Definition 12 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap\ (c_2Ebool_2E_21\ 2)\ (\lambda V2t \in 2.V2t))))$

Assume the following.

$$\begin{aligned} & (\forall V0x \in ty_2Erealax_2Ereal.(\forall V1y \in ty_2Erealax_2Ereal. \\ & ((ap\ (ap\ c_2Erealax_2Ereal_mul\ V0x)\ V1y) = (ap\ (ap\ c_2Erealax_2Ereal_mul \\ & \quad V1y)\ V0x)))) \end{aligned} \quad (12)$$

Assume the following.

$$\begin{aligned} & (\forall V0x \in ty_2Erealax_2Ereal.(\forall V1y \in ty_2Erealax_2Ereal. \\ & (\forall V2z \in ty_2Erealax_2Ereal.(((\neg(V2z = (ap\ c_2Ereal_2Ereal_of_num \\ & \quad c_2Enum_2E0)))) \wedge ((ap\ (ap\ c_2Erealax_2Ereal_mul\ V0x)\ V2z) = (ap \\ & \quad (ap\ c_2Erealax_2Ereal_mul\ V1y)\ V2z))) \Rightarrow (V0x = V1y)))) \end{aligned} \quad (13)$$

Theorem 1

$(\forall V0x \in ty_2Erealax_2Ereal. (\forall V1y \in ty_2Erealax_2Ereal. (\forall V2z \in ty_2Erealax_2Ereal. (((\neg(V0x = (ap\ c_2Ereal_2Ereal_of_num\ c_2Enum_2E0))) \wedge ((ap\ (ap\ c_2Erealax_2Ereal_mul\ V0x)\ V1y) = (ap\ (ap\ c_2Erealax_2Ereal_mul\ V0x)\ V2z))) \Rightarrow (V1y = V2z))))))$