

thm_2Ereal_2EREAL__EQ__MUL__LCANCEL
(TMX7zPxuQYEmGnWV1L1ShZ4j6H4hyEtfPXp)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define c_2Ebool_2ET to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Let $c_2Enum_2EZERO_REP : \iota$ be given. Assume the following.

$$c_2Enum_2EZERO_REP \in \omega \tag{1}$$

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \tag{2}$$

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum^{\omega}) \tag{3}$$

Definition 3 We define c_2Enum_2E0 to be $(ap\ c_2Enum_2EABS_num\ c_2Enum_2EZERO_REP)$.

Let $ty_2Erealx_2Ereal : \iota$ be given. Assume the following.

$$nonempty\ ty_2Erealx_2Ereal \tag{4}$$

Let $c_2Ereal_2Ereal_of_num : \iota$ be given. Assume the following.

$$c_2Ereal_2Ereal_of_num \in (ty_2Erealx_2Ereal^{ty_2Enum_2Enum}) \tag{5}$$

Definition 4 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p \Rightarrow q)$ of type ι .

Definition 5 We define $c_2Ebool_2E_21$ to be $\lambda A.\lambda a : \iota.(\lambda V0P \in (2^{A-27a}).(ap (ap (c_2Emin_2E_3D (2^{A-27a}))$

Definition 6 We define $c_2Ebool_2E_5C_2F$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2$

Let $ty_2Ehreal_2Ehreal : \iota$ be given. Assume the following.

$$nonempty\ ty_2Ehreal_2Ehreal \quad (6)$$

Let $ty_2Epair_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty_2Epair_2Eprod\ A0\ A1) \quad (7)$$

Let $c_2Erealax_2Ereal_REP_CLASS : \iota$ be given. Assume the following.

$$c_2Erealax_2Ereal_REP_CLASS \in ((2^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)})^{ty_2Erealax_2Ereal_REP_CLASS}) \quad (8)$$

Definition 7 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A.$ if $(\exists x \in A.p\ (ap\ P\ x))$ then (the $(\lambda x.x \in A \wedge p$ of type $\iota \Rightarrow \iota$).

Definition 8 We define $c_2Erealax_2Ereal_REP$ to be $\lambda V0a \in ty_2Erealax_2Ereal.(ap\ (c_2Emin_2E_40\ (ty$

Let $c_2Erealax_2Etrealmul : \iota$ be given. Assume the following.

$$c_2Erealax_2Etrealmul \in (((ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)})^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)}) \quad (9)$$

Let $c_2Erealax_2Etrealeq : \iota$ be given. Assume the following.

$$c_2Erealax_2Etrealeq \in ((2^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)})^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)}) \quad (10)$$

Let $c_2Erealax_2Ereal_ABS_CLASS : \iota$ be given. Assume the following.

$$c_2Erealax_2Ereal_ABS_CLASS \in (ty_2Erealax_2Ereal)^{(2^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)})} \quad (11)$$

Definition 9 We define $c_2Erealax_2Ereal_ABS$ to be $\lambda V0r \in (ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)$

Definition 10 We define $c_2Erealax_2Ereal_mul$ to be $\lambda V0T1 \in ty_2Erealax_2Ereal.\lambda V1T2 \in ty_2Erealax_2Ereal$

Assume the following.

$$True \quad (12)$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0t \in 2.((\forall V1x \in A_27a.(p\ V0t) \Leftrightarrow (p\ V0t)))) \quad (13)$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0x \in A_27a.((V0x = V0x) \Leftrightarrow True)) \quad (14)$$

Assume the following.

$$\begin{aligned}
& (\forall V0x \in ty_2Erealax_2Ereal. (\forall V1y \in ty_2Erealax_2Ereal. \\
& (\forall V2z \in ty_2Erealax_2Ereal. (((ap (ap c_2Erealax_2Ereal_mul \\
V0x) V1y) = (ap (ap c_2Erealax_2Ereal_mul V0x) V2z)) \Leftrightarrow ((V0x = (ap \\
c_2Ereal_2Ereal_of_num c_2Enum_2E0)) \vee (V1y = V2z))))))
\end{aligned} \tag{15}$$

Theorem 1

$$\begin{aligned}
& (\forall V0x \in ty_2Erealax_2Ereal. (\forall V1y \in ty_2Erealax_2Ereal. \\
& (\forall V2z \in ty_2Erealax_2Ereal. (((ap (ap c_2Erealax_2Ereal_mul \\
V0x) V1y) = (ap (ap c_2Erealax_2Ereal_mul V0x) V2z)) \Leftrightarrow ((V0x = (ap \\
c_2Ereal_2Ereal_of_num c_2Enum_2E0)) \vee (V1y = V2z))))))
\end{aligned}$$