

thm_2Ereal_2EREAL__IMP__MIN__LE2
(TMF7QQXLagTaQNa5rn7sKRdrWNp5yMDUTXT)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define $c_2Ebool_2E_ET$ to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 3 We define $c_2Ebool_2E_21$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap (ap (c_2Emin_2E_3D (2^{A_27a}))$

Definition 4 We define $c_2Ebool_2E_EF$ to be $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$.

Definition 5 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow p Q)$ of type ι .

Definition 6 We define $c_2Ebool_2E_7E$ to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2E_EF$

Definition 7 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t)$

Definition 8 We define $c_2Ebool_2E_5C_2F$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t)$

Let $ty_2Ehreal_2Ehreal : \iota$ be given. Assume the following.

$$nonempty\ ty_2Ehreal_2Ehreal \tag{1}$$

Let $ty_2Epair_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty_2Epair_2Eprod\ A0\ A1) \tag{2}$$

Let $ty_2Erealax_2Ereal : \iota$ be given. Assume the following.

$$nonempty\ ty_2Erealax_2Ereal \tag{3}$$

Let $c_2Erealax_2Ereal_REP_CLASS : \iota$ be given. Assume the following.

$$c_2Erealax_2Ereal_REP_CLASS \in ((2^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)})ty_2Erealax_2Ereal) \tag{4}$$

Assume the following.

$$\begin{aligned}
& (\forall V0z \in ty_2Erealax_2Ereal. (\forall V1x \in ty_2Erealax_2Ereal. \\
& (\forall V2y \in ty_2Erealax_2Ereal. ((p (ap (ap c_2Ereal_2Ereal_lte \\
V0z) (ap (ap c_2Ereal_2Ereal_emin V1x) V2y))) \Leftrightarrow ((p (ap (ap c_2Ereal_2Ereal_lte \\
V0z) V1x)) \wedge (p (ap (ap c_2Ereal_2Ereal_lte V0z) V2y)))))))
\end{aligned} \tag{12}$$

Assume the following.

$$\begin{aligned}
& (\forall V0z \in ty_2Erealax_2Ereal. (\forall V1x \in ty_2Erealax_2Ereal. \\
& (\forall V2y \in ty_2Erealax_2Ereal. ((p (ap (ap c_2Ereal_2Ereal_lte \\
(ap (ap c_2Ereal_2Ereal_emin V1x) V2y)) V0z)) \Leftrightarrow ((p (ap (ap c_2Ereal_2Ereal_lte \\
V1x) V0z)) \vee (p (ap (ap c_2Ereal_2Ereal_lte V2y) V0z)))))))
\end{aligned} \tag{13}$$

Theorem 1

$$\begin{aligned}
& (\forall V0x1 \in ty_2Erealax_2Ereal. (\forall V1x2 \in ty_2Erealax_2Ereal. \\
& (\forall V2y1 \in ty_2Erealax_2Ereal. (\forall V3y2 \in ty_2Erealax_2Ereal. \\
(((p (ap (ap c_2Ereal_2Ereal_lte V0x1) V2y1)) \wedge (p (ap (ap c_2Ereal_2Ereal_lte \\
V1x2) V3y2))) \Rightarrow (p (ap (ap c_2Ereal_2Ereal_lte (ap (ap c_2Ereal_2Ereal_emin \\
V0x1) V1x2)) (ap (ap c_2Ereal_2Ereal_emin V2y1) V3y2)))))))
\end{aligned}$$