

thm_2Ereal_2EREAL__LET__TOTAL
(TMEzbYo3ciQ6XxxLXwQLE8dPqpN7rLDma3E)

October 26, 2020

Definition 1 We define `c_2Emin_2E_3D_3D_3E` to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow p Q)$ of type ι .

Definition 2 We define `c_2Emin_2E_3D` to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 3 We define `c_2Ebool_2ET` to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 4 We define `c_2Ebool_2E_21` to be $\lambda A_{27a} : \iota.(\lambda V0P \in (2^{A-27a}).(ap (ap (c_2Emin_2E_3D (2^{A-27a}))$

Definition 5 We define `c_2Ebool_2E_5C_2F` to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t))$

Definition 6 We define `c_2Ebool_2EF` to be $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$.

Definition 7 We define `c_2Ebool_2E_2F_5C` to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t))$

Let `ty_2Ehreal_2Ehreal` : ι be given. Assume the following.

$$nonempty\ ty_2Ehreal_2Ehreal \tag{1}$$

Let `ty_2Epair_2Eprod` : $\iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty_2Epair_2Eprod\ A0\ A1) \tag{2}$$

Let `ty_2Erealax_2Ereal` : ι be given. Assume the following.

$$nonempty\ ty_2Erealax_2Ereal \tag{3}$$

Let `c_2Erealax_2Ereal__REP__CLASS` : ι be given. Assume the following.

$$c_2Erealax_2Ereal_REP_CLASS \in ((2^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)})ty_2Erealax) \tag{4}$$

Definition 8 We define `c_2Emin_2E_40` to be $\lambda A.\lambda P \in 2^A.if (\exists x \in A.p (ap P x))$ **then** (the $(\lambda x.x \in A \wedge p$ of type $\iota \Rightarrow \iota$).

Definition 9 We define $c_2Erealax_2Ereal_REP$ to be $\lambda V0a \in ty_2Erealax_2Ereal.(ap (c_2Emin_2E40 (ty$
 Let $c_2Erealax_2Etreall_lt : \iota$ be given. Assume the following.

$$c_2Erealax_2Etreall_lt \in ((2^{(ty_2Epair_2Eprod ty_2Ehreal_2Ehreal ty_2Ehreal_2Ehreal)})(ty_2Epair_2Eprod ty_2Ehreal_2Ehreal)) \quad (5)$$

Definition 10 We define $c_2Erealax_2Ereal_lt$ to be $\lambda V0T1 \in ty_2Erealax_2Ereal.\lambda V1T2 \in ty_2Erealax_2Ereal.$

Definition 11 We define c_2Ebool_2E7E to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E3D_3D_3E V0t) c_2Ebool_2E7E$

Definition 12 We define $c_2Ereal_2Ereal_lte$ to be $\lambda V0x \in ty_2Erealax_2Ereal.\lambda V1y \in ty_2Erealax_2Ereal.$

Assume the following.

$$True \quad (6)$$

Assume the following.

$$(\forall V0t \in 2.(((True \vee (p V0t)) \Leftrightarrow True) \wedge (((p V0t) \vee True) \Leftrightarrow True) \wedge \\ (((False \vee (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \vee False) \Leftrightarrow (p V0t)) \wedge (((p V0t) \vee \\ (p V0t)) \Leftrightarrow (p V0t)))))) \quad (7)$$

Assume the following.

$$((\forall V0t \in 2.((\neg(\neg(p V0t))) \Leftrightarrow (p V0t))) \wedge ((\neg True) \Leftrightarrow False) \wedge \\ ((\neg False) \Leftrightarrow True))) \quad (8)$$

Theorem 1

$$(\forall V0x \in ty_2Erealax_2Ereal.(\forall V1y \in ty_2Erealax_2Ereal. \\ ((p (ap (ap c_2Ereal_2Ereal_lte V0x) V1y)) \vee (p (ap (ap c_2Erealax_2Ereal_lt \\ V1y) V0x))))))$$