

thm_2Ereal_2EREAL__LE__NEG2 (TMYfbQma- Tod1N3rBJbR7Qqjne5MPMSKBDNV)

October 26, 2020

Let $ty_2Ehreal_2Ehreal : \iota$ be given. Assume the following.

$$nonempty\ ty_2Ehreal_2Ehreal \tag{1}$$

Let $ty_2Epair_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty_2Epair_2Eprod\ A0\ A1) \tag{2}$$

Let $ty_2Erealax_2Ereal : \iota$ be given. Assume the following.

$$nonempty\ ty_2Erealax_2Ereal \tag{3}$$

Let $c_2Erealax_2Ereal_REP_CLASS : \iota$ be given. Assume the following.

$$c_2Erealax_2Ereal_REP_CLASS \in ((2^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)})_{ty_2Erealax_2Ereal}) \tag{4}$$

Definition 1 We define c_2Emin_2E40 to be $\lambda A.\lambda P \in 2^A.$ if $(\exists x \in A.p\ (ap\ P\ x))$ **then** (the $(\lambda x.x \in A \wedge p)$ of type $\iota \Rightarrow \iota$).

Definition 2 We define c_2Emin_2E3D to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o\ (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 3 We define c_2Ebool_2E27 to be $(ap\ (ap\ (c_2Emin_2E3D\ (2^2))\ (\lambda V0x \in 2.V0x))\ (\lambda V1x \in 2.V1x))$

Definition 4 We define c_2Ebool_2E21 to be $\lambda A.27a : \iota.(\lambda V0P \in (2^{A-27a}).(ap\ (ap\ (c_2Emin_2E3D\ (2^{A-27a}))\ (\lambda V1x \in 2.V1x))\ (\lambda V2x \in 2.V2x))$

Definition 5 We define $c_2Erealax_2Ereal_REP$ to be $\lambda V0a \in ty_2Erealax_2Ereal.(ap\ (c_2Emin_2E40\ (ty_2Erealax_2Ereal\ a)))$

Let $c_2Erealax_2Etrealm_neg : \iota$ be given. Assume the following.

$$c_2Erealax_2Etrealm_neg \in ((ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)_{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)}) \tag{5}$$

Let $c_2Erealax_2Etreall_eq : \iota$ be given. Assume the following.

$$c_2Erealax_2Etreall_eq \in ((2^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)}) (ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal)) \quad (6)$$

Let $c_2Erealax_2Ereal_ABS_CLASS : \iota$ be given. Assume the following.

$$c_2Erealax_2Ereal_ABS_CLASS \in (ty_2Erealax_2Ereal)^{(2^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)})} \quad (7)$$

Definition 6 We define $c_2Erealax_2Ereal_ABS$ to be $\lambda V0r \in (ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)$

Definition 7 We define $c_2Erealax_2Ereal_neg$ to be $\lambda V0T1 \in ty_2Erealax_2Ereal.(ap\ c_2Erealax_2Ereal_neg)$

Let $c_2Erealax_2Etreall_lt : \iota$ be given. Assume the following.

$$c_2Erealax_2Etreall_lt \in ((2^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)}) (ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal)) \quad (8)$$

Definition 8 We define $c_2Erealax_2Ereal_lt$ to be $\lambda V0T1 \in ty_2Erealax_2Ereal.\lambda V1T2 \in ty_2Erealax_2Ereal$

Definition 9 We define c_2Ebool_2EF to be $(ap\ (c_2Ebool_2E21\ 2))\ (\lambda V0t \in 2.V0t)$.

Definition 10 We define $c_2Emin_2E3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o\ (p\ P \Rightarrow p\ Q)$ of type ι .

Definition 11 We define c_2Ebool_2E7E to be $(\lambda V0t \in 2.(ap\ (ap\ c_2Emin_2E3D_3D_3E\ V0t)\ c_2Ebool_2E7E))$

Definition 12 We define $c_2Ereal_2Ereal_lte$ to be $\lambda V0x \in ty_2Erealax_2Ereal.\lambda V1y \in ty_2Erealax_2Ereal$

Assume the following.

$$\begin{aligned} & (\forall V0x \in ty_2Erealax_2Ereal.(\forall V1y \in ty_2Erealax_2Ereal. \\ & ((p\ (ap\ (ap\ c_2Ereal_2Ereal_lte\ (ap\ c_2Erealax_2Ereal_neg\ V0x)) \\ & (ap\ c_2Erealax_2Ereal_neg\ V1y)))) \Leftrightarrow (p\ (ap\ (ap\ c_2Ereal_2Ereal_lte \\ & V1y)\ V0x)))) \end{aligned} \quad (9)$$

Theorem 1

$$\begin{aligned} & (\forall V0x \in ty_2Erealax_2Ereal.(\forall V1y \in ty_2Erealax_2Ereal. \\ & ((p\ (ap\ (ap\ c_2Ereal_2Ereal_lte\ (ap\ c_2Erealax_2Ereal_neg\ V0x)) \\ & (ap\ c_2Erealax_2Ereal_neg\ V1y)))) \Leftrightarrow (p\ (ap\ (ap\ c_2Ereal_2Ereal_lte \\ & V1y)\ V0x)))) \end{aligned}$$