

thm_2Ereal_2EREAL__LE__NEGR
(TMRqTbcyLk8G5ifBTcLLVFSCVe4rgpiz7yh)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define c_2Ebool_2ET to be $(ap (ap (c_2Emin_2E_3D (2^2))) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Let $c_2Enum_2EZERO_REP : \iota$ be given. Assume the following.

$$c_2Enum_2EZERO_REP \in \omega \tag{1}$$

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \tag{2}$$

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum^{\omega}) \tag{3}$$

Definition 3 We define c_2Enum_2E0 to be $(ap\ c_2Enum_2EABS_num\ c_2Enum_2EZERO_REP)$.

Let $ty_2Erealx_2Ereal : \iota$ be given. Assume the following.

$$nonempty\ ty_2Erealx_2Ereal \tag{4}$$

Let $c_2Ereal_2Ereal_of_num : \iota$ be given. Assume the following.

$$c_2Ereal_2Ereal_of_num \in (ty_2Erealx_2Ereal^{ty_2Enum_2Enum}) \tag{5}$$

Let $ty_2Ehreal_2Ehreal : \iota$ be given. Assume the following.

$$nonempty\ ty_2Ehreal_2Ehreal \tag{6}$$

Let $ty_2Epair_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty_2Epair_2Eprod\ A0\ A1) \tag{7}$$

Let $c_2Erealx_2Ereal_REP_CLASS : \iota$ be given. Assume the following.

$$c_2Erealx_2Ereal_REP_CLASS \in ((2^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)})^{ty_2Erealx_2Ereal}) \tag{8}$$

Definition 4 We define `c_2Emin_2E_40` to be $\lambda A. \lambda P \in 2^A. \mathbf{if} (\exists x \in A. p (ap P x)) \mathbf{then}$ (the $(\lambda x. x \in A \wedge p$ of type $\iota \Rightarrow \iota$).

Definition 5 We define `c_2Ebool_2E_21` to be $\lambda A_{27a} : \iota. (\lambda V0P \in (2^{A_{27a}}). (ap (ap (c_2Emin_2E_3D (2^{A_{27a}})$

Definition 6 We define `c_2Erealx_2Ereal__REP` to be $\lambda V0a \in ty_2Erealx_2Ereal. (ap (c_2Emin_2E_40 (ty$

Let `c_2Erealx_2Etrealm__neg` : ι be given. Assume the following.

$$c_2Erealx_2Etrealm_neg \in ((ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal) (ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)) \quad (9)$$

Let `c_2Erealx_2Etrealm__eq` : ι be given. Assume the following.

$$c_2Erealx_2Etrealm_eq \in ((2^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)} (ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal)) (ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal)) \quad (10)$$

Let `c_2Erealx_2Ereal__ABS__CLASS` : ι be given. Assume the following.

$$c_2Erealx_2Ereal_ABS_CLASS \in (ty_2Erealx_2Ereal)^{(2^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)} (ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal))} \quad (11)$$

Definition 7 We define `c_2Erealx_2Ereal__ABS` to be $\lambda V0r \in (ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)$

Definition 8 We define `c_2Erealx_2Ereal__neg` to be $\lambda V0T1 \in ty_2Erealx_2Ereal. (ap\ c_2Erealx_2Ereal_neg$

Let `c_2Erealx_2Etrealm__lt` : ι be given. Assume the following.

$$c_2Erealx_2Etrealm_lt \in ((2^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)} (ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal)) (ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal)) \quad (12)$$

Definition 9 We define `c_2Erealx_2Ereal__lt` to be $\lambda V0T1 \in ty_2Erealx_2Ereal. \lambda V1T2 \in ty_2Erealx_2Ereal.$

Definition 10 We define `c_2Ebool_2E_21` to be $(ap (c_2Ebool_2E_21\ 2) (\lambda V0t \in 2.V0t))$.

Definition 11 We define `c_2Emin_2E_3D_3D_3E` to be $\lambda P \in 2. \lambda Q \in 2. inj_o (p\ P \Rightarrow p\ Q)$ of type ι .

Definition 12 We define `c_2Ebool_2E_7E` to be $(\lambda V0t \in 2. (ap (ap\ c_2Emin_2E_3D_3D_3E\ V0t)\ c_2Ebool_2E_7E$

Definition 13 We define `c_2Ereal_2Ereal__lte` to be $\lambda V0x \in ty_2Erealx_2Ereal. \lambda V1y \in ty_2Erealx_2Ereal.$

Assume the following.

$$True \quad (13)$$

Assume the following.

$$\forall A_{27a}. nonempty\ A_{27a} \Rightarrow (\forall V0x \in A_{27a}. ((V0x = V0x) \Leftrightarrow True)) \quad (14)$$

Assume the following.

$$(\forall V0x \in ty_2Erealx_2Ereal. ((ap\ c_2Erealx_2Ereal_neg\ (ap\ c_2Erealx_2Ereal_neg\ V0x)) = V0x)) \quad (15)$$

Assume the following.

$$\begin{aligned}
 & (\forall V0x \in ty_2Erealax_2Ereal. ((p (ap (ap c_2Ereal_2Ereal_lte \\
 & (ap c_2Ereal_2Ereal_of_num c_2Enum_2E0)) (ap c_2Erealax_2Ereal_neg \\
 & V0x))) \Leftrightarrow (p (ap (ap c_2Ereal_2Ereal_lte V0x) (ap c_2Ereal_2Ereal_of_num \\
 & c_2Enum_2E0))))))
 \end{aligned} \tag{16}$$

Assume the following.

$$\begin{aligned}
 & (\forall V0x \in ty_2Erealax_2Ereal. ((p (ap (ap c_2Ereal_2Ereal_lte \\
 & (ap c_2Erealax_2Ereal_neg V0x)) V0x)) \Leftrightarrow (p (ap (ap c_2Ereal_2Ereal_lte \\
 & (ap c_2Ereal_2Ereal_of_num c_2Enum_2E0)) V0x))))))
 \end{aligned} \tag{17}$$

Theorem 1

$$\begin{aligned}
 & (\forall V0x \in ty_2Erealax_2Ereal. ((p (ap (ap c_2Ereal_2Ereal_lte \\
 & V0x) (ap c_2Erealax_2Ereal_neg V0x))) \Leftrightarrow (p (ap (ap c_2Ereal_2Ereal_lte \\
 & V0x) (ap c_2Ereal_2Ereal_of_num c_2Enum_2E0))))))
 \end{aligned}$$