

thm\_2Ereal\_2EREAL\_\_LE\_\_SQUARE  
(TMVT7pKZxCT4Rfv9VLX8d7ynKr1ujVU7Kff)

October 26, 2020

**Definition 1** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define  $c\_2Ebool\_2E\_7E$  to be  $(ap (ap (c\_2Emin\_2E\_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

**Definition 3** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A.27a : \iota.(\lambda V0P \in (2^{A-27a}).(ap (ap (c\_2Emin\_2E\_3D (2^{A-27a}))$

**Definition 4** We define  $c\_2Ebool\_2E\_7E$  to be  $(ap (c\_2Ebool\_2E\_21 2) (\lambda V0t \in 2.V0t))$ .

**Definition 5** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o (p P \Rightarrow p Q)$  of type  $\iota$ .

**Definition 6** We define  $c\_2Ebool\_2E\_7E$  to be  $(\lambda V0t \in 2.(ap (ap c\_2Emin\_2E\_3D\_3D\_3E V0t) c\_2Ebool\_2E\_7E))$

Let  $ty\_2Ehreal\_2Ehreal : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Ehreal\_2Ehreal \tag{1}$$

Let  $ty\_2Epair\_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty\_2Epair\_2Eprod\ A0\ A1) \tag{2}$$

Let  $ty\_2Erealax\_2Ereal : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Erealax\_2Ereal \tag{3}$$

Let  $c\_2Erealax\_2Ereal\_REP\_CLASS : \iota$  be given. Assume the following.

$$c\_2Erealax\_2Ereal\_REP\_CLASS \in ((2^{(ty\_2Epair\_2Eprod\ ty\_2Ehreal\_2Ehreal\ ty\_2Ehreal\_2Ehreal)})\ ty\_2Erealax\_2Ereal) \tag{4}$$

**Definition 7** We define  $c\_2Emin\_2E\_40$  to be  $\lambda A.\lambda P \in 2^A.if (\exists x \in A.p (ap P x)) \mathbf{then} (the (\lambda x.x \in A \wedge p))$  of type  $\iota \Rightarrow \iota$ .

**Definition 8** We define  $c\_2Erealax\_2Ereal\_REP$  to be  $\lambda V0a \in ty\_2Erealax\_2Ereal.(ap (c\_2Emin\_2E\_40 (ty\_2Erealax\_2Ereal\_REP\_CLASS V0a)))$

Let  $c\_2Erealax\_2Etreal\_neg : \iota$  be given. Assume the following.

$$c\_2Erealax\_2Etreal\_neg \in ((ty\_2Epair\_2Eprod\ ty\_2Ehreal\_2Ehreal\ ty\_2Ehreal\_2Ehreal)^{(ty\_2Epair\_2Eprod\ ty\_2Ehreal\_2Ehreal\ ty\_2Ehreal\_2Ehreal)}) \quad (5)$$

Let  $c\_2Erealax\_2Etreal\_eq : \iota$  be given. Assume the following.

$$c\_2Erealax\_2Etreal\_eq \in ((2^{(ty\_2Epair\_2Eprod\ ty\_2Ehreal\_2Ehreal\ ty\_2Ehreal\_2Ehreal)})^{(ty\_2Epair\_2Eprod\ ty\_2Ehreal\_2Ehreal)}) \quad (6)$$

Let  $c\_2Erealax\_2Ereal\_ABS\_CLASS : \iota$  be given. Assume the following.

$$c\_2Erealax\_2Ereal\_ABS\_CLASS \in (ty\_2Erealax\_2Ereal)^{(2^{(ty\_2Epair\_2Eprod\ ty\_2Ehreal\_2Ehreal\ ty\_2Ehreal\_2Ehreal)})} \quad (7)$$

**Definition 9** We define  $c\_2Erealax\_2Ereal\_ABS$  to be  $\lambda V0r \in (ty\_2Epair\_2Eprod\ ty\_2Ehreal\_2Ehreal\ ty\_2Ehreal\_2Ehreal)$

**Definition 10** We define  $c\_2Erealax\_2Ereal\_neg$  to be  $\lambda V0T1 \in ty\_2Erealax\_2Ereal.(ap\ c\_2Erealax\_2Ereal)$

**Definition 11** We define  $c\_2Ebool\_2E\_5C\_2F$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap\ (c\_2Ebool\_2E\_21\ 2)\ (\lambda V2t \in 2)))$

Let  $c\_2Erealax\_2Etreal\_mul : \iota$  be given. Assume the following.

$$c\_2Erealax\_2Etreal\_mul \in (((ty\_2Epair\_2Eprod\ ty\_2Ehreal\_2Ehreal\ ty\_2Ehreal\_2Ehreal)^{(ty\_2Epair\_2Eprod\ ty\_2Ehreal\_2Ehreal\ ty\_2Ehreal\_2Ehreal)})^{(ty\_2Epair\_2Eprod\ ty\_2Ehreal\_2Ehreal\ ty\_2Ehreal\_2Ehreal)}) \quad (8)$$

**Definition 12** We define  $c\_2Erealax\_2Ereal\_mul$  to be  $\lambda V0T1 \in ty\_2Erealax\_2Ereal.\lambda V1T2 \in ty\_2Erealax\_2Ereal$

Let  $c\_2Enum\_2EZERO\_REP : \iota$  be given. Assume the following.

$$c\_2Enum\_2EZERO\_REP \in \omega \quad (9)$$

Let  $ty\_2Enum\_2Enum : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Enum\_2Enum \quad (10)$$

Let  $c\_2Enum\_2EABS\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EABS\_num \in (ty\_2Enum\_2Enum)^{\omega} \quad (11)$$

**Definition 13** We define  $c\_2Enum\_2E0$  to be  $(ap\ c\_2Enum\_2EABS\_num\ c\_2Enum\_2EZERO\_REP)$ .

Let  $c\_2Ereal\_2Ereal\_of\_num : \iota$  be given. Assume the following.

$$c\_2Ereal\_2Ereal\_of\_num \in (ty\_2Erealax\_2Ereal)^{ty\_2Enum\_2Enum} \quad (12)$$

Let  $c\_2Erealax\_2Etreal\_lt : \iota$  be given. Assume the following.

$$c\_2Erealax\_2Etreal\_lt \in ((2^{(ty\_2Epair\_2Eprod\ ty\_2Ehreal\_2Ehreal\ ty\_2Ehreal\_2Ehreal)})^{(ty\_2Epair\_2Eprod\ ty\_2Ehreal\_2Ehreal)}) \quad (13)$$

**Definition 14** We define  $c\_2Erealax\_2Ereal\_lt$  to be  $\lambda V0T1 \in ty\_2Erealax\_2Ereal.\lambda V1T2 \in ty\_2Erealax\_2Ereal.$

**Definition 15** We define  $c\_2Ereal\_2Ereal\_lte$  to be  $\lambda V0x \in ty\_2Erealax\_2Ereal.\lambda V1y \in ty\_2Erealax\_2Ereal.$

**Definition 16** We define  $c\_2Ebool\_2E\_2F\_5C$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_2Ebool\_2E\_21 2) (\lambda V2t \in 2.$

Assume the following.

$$True \quad (14)$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2.(((True \Rightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Rightarrow True) \Leftrightarrow \\ & True) \wedge (((False \Rightarrow (p V0t)) \Leftrightarrow True) \wedge (((p V0t) \Rightarrow (p V0t)) \Leftrightarrow True) \wedge (( \\ & (p V0t) \Rightarrow False) \Leftrightarrow (\neg (p V0t)))))) \end{aligned} \quad (15)$$

Assume the following.

$$\begin{aligned} & \forall A\_27a.nonempty A\_27a \Rightarrow (\forall V0x \in A\_27a.(\forall V1y \in \\ & A\_27a.((V0x = V1y) \Leftrightarrow (V1y = V0x)))) \end{aligned} \quad (16)$$

Assume the following.

$$\begin{aligned} & (\forall V0x \in ty\_2Erealax\_2Ereal.(\forall V1y \in ty\_2Erealax\_2Ereal. \\ & ((ap c\_2Erealax\_2Ereal\_neg (ap (ap c\_2Erealax\_2Ereal\_mul V0x) \\ & V1y)) = (ap (ap c\_2Erealax\_2Ereal\_mul (ap c\_2Erealax\_2Ereal\_neg \\ & V0x)) V1y)))) \end{aligned} \quad (17)$$

Assume the following.

$$\begin{aligned} & (\forall V0x \in ty\_2Erealax\_2Ereal.(\forall V1y \in ty\_2Erealax\_2Ereal. \\ & ((ap c\_2Erealax\_2Ereal\_neg (ap (ap c\_2Erealax\_2Ereal\_mul V0x) \\ & V1y)) = (ap (ap c\_2Erealax\_2Ereal\_mul V0x) (ap c\_2Erealax\_2Ereal\_neg \\ & V1y)))) \end{aligned} \quad (18)$$

Assume the following.

$$\begin{aligned} & (\forall V0x \in ty\_2Erealax\_2Ereal.((ap c\_2Erealax\_2Ereal\_neg \\ & (ap c\_2Erealax\_2Ereal\_neg V0x)) = V0x)) \end{aligned} \quad (19)$$

Assume the following.

$$\begin{aligned} & (\forall V0x \in ty\_2Erealax\_2Ereal.((p (ap (ap c\_2Ereal\_2Ereal\_lte \\ & (ap c\_2Ereal\_2Ereal\_of\_num c\_2Enum\_2E0)) V0x)) \vee (p (ap (ap c\_2Ereal\_2Ereal\_lte \\ & (ap c\_2Ereal\_2Ereal\_of\_num c\_2Enum\_2E0)) (ap c\_2Erealax\_2Ereal\_neg \\ & V0x)))))) \end{aligned} \quad (20)$$

Assume the following.

$$\begin{aligned} & (\forall V0x \in ty\_2Erealax\_2Ereal.(\forall V1y \in ty\_2Erealax\_2Ereal. \\ & (((p (ap (ap c\_2Ereal\_2Ereal\_lte (ap c\_2Ereal\_2Ereal\_of\_num \\ & c\_2Enum\_2E0)) V0x)) \wedge (p (ap (ap c\_2Ereal\_2Ereal\_lte (ap c\_2Ereal\_2Ereal\_of\_num \\ & c\_2Enum\_2E0)) V1y))) \Rightarrow (p (ap (ap c\_2Ereal\_2Ereal\_lte (ap c\_2Ereal\_2Ereal\_of\_num \\ & c\_2Enum\_2E0)) (ap (ap c\_2Erealax\_2Ereal\_mul V0x) V1y)))))) \end{aligned} \quad (21)$$

**Theorem 1**

$(\forall V0x \in ty\_2Erealax\_2Ereal.(p (ap (ap c\_2Ereal\_2Ereal\_lte$   
 $(ap c\_2Ereal\_2Ereal\_of\_num c\_2Enum\_2E0)) (ap (ap c\_2Erealax\_2Ereal\_mul$   
 $V0x) V0x))))$