

thm_2Ereal_2EREAL_LT_HALF1
(TMd1iAPWvEFJS3Tq3jVZy13fuu6icSwJWG5)

October 26, 2020

Let $c_2Enum_2EZERO_REP : \iota$ be given. Assume the following.

$$c_2Enum_2EZERO_REP \in \omega \tag{1}$$

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \tag{2}$$

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum^{\omega}) \tag{3}$$

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define c_2Enum_2E0 to be $(ap\ c_2Enum_2EABS_num\ c_2Enum_2EZERO_REP)$.

Let $c_2Enum_2EREP_num : \iota$ be given. Assume the following.

$$c_2Enum_2EREP_num \in (\omega^{ty_2Enum_2Enum}) \tag{4}$$

Let $c_2Enum_2ESUC_REP : \iota$ be given. Assume the following.

$$c_2Enum_2ESUC_REP \in (\omega^{\omega}) \tag{5}$$

Definition 3 We define c_2Ebool_2ET to be $(ap\ (ap\ (c_2Emin_2E_3D\ (2^2))\ (\lambda V0x \in 2.V0x))\ (\lambda V1x \in 2.V1x))$

Definition 4 We define $c_2Ebool_2E_21$ to be $\lambda A.\lambda a : \iota.(\lambda V0P \in (2^{A-27a}).(ap\ (ap\ (c_2Emin_2E_3D\ (2^{A-27a}))\ (\lambda V1x \in 2.V1x))\ (\lambda V1x \in 2.V1x))$

Definition 5 We define c_2Enum_2ESUC to be $\lambda V0m \in ty_2Enum_2Enum.(ap\ c_2Enum_2EABS_num\ (c_2Enum_2ESUC_REP\ m))$

Let $c_2Earithmetic_2E_2B : \iota$ be given. Assume the following.

$$c_2Earithmetic_2E_2B \in ((ty_2Enum_2Enum^{ty_2Enum_2Enum})^{ty_2Enum_2Enum}) \tag{6}$$

Definition 6 We define `c_2Earithmic_2EBIT1` to be $\lambda V0n \in ty_2Enum_2Enum.(ap (ap c_2Earithmic_2EBIT1))$.

Definition 7 We define `c_2Earithmic_2EZERO` to be `c_2Enum_2E0`.

Definition 8 We define `c_2Earithmic_2EBIT2` to be $\lambda V0n \in ty_2Enum_2Enum.(ap (ap c_2Earithmic_2EBIT2))$.

Definition 9 We define `c_2Earithmic_2ENUMERAL` to be $\lambda V0x \in ty_2Enum_2Enum.V0x$.

Definition 10 We define `c_2Ebool_2EF` to be $(ap (c_2Ebool_2E_21)) (\lambda V0t \in 2.V0t)$.

Definition 11 We define `c_2Emin_2E_3D_3D_3E` to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow p Q)$ of type ι .

Definition 12 We define `c_2Ebool_2E_2F_5C` to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21)) (\lambda V2t \in 2.V0t)))$.

Let `ty_2Ehreal_2Ehreal` : ι be given. Assume the following.

$$nonempty\ ty_2Ehreal_2Ehreal \tag{7}$$

Let `ty_2Epair_2Eprod` : $\iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty_2Epair_2Eprod\ A0\ A1) \tag{8}$$

Let `ty_2Erealax_2Ereal` : ι be given. Assume the following.

$$nonempty\ ty_2Erealax_2Ereal \tag{9}$$

Let `c_2Erealax_2Ereal_2REP_2CLASS` : ι be given. Assume the following.

$$c_2Erealax_2Ereal_2REP_2CLASS \in ((2^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)})^{ty_2Erealax_2Ereal}) \tag{10}$$

Definition 13 We define `c_2Emin_2E_40` to be $\lambda A.\lambda P \in 2^A.if (\exists x \in A.p (ap P x)) then (the (\lambda x.x \in A \wedge P x))$ of type $\iota \Rightarrow \iota$.

Definition 14 We define `c_2Erealax_2Ereal_2REP` to be $\lambda V0a \in ty_2Erealax_2Ereal.(ap (c_2Emin_2E_40))$.

Let `c_2Erealax_2Etreall_inv` : ι be given. Assume the following.

$$c_2Erealax_2Etreall_inv \in ((ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)^{ty_2Erealax_2Ereal}) \tag{11}$$

Let `c_2Erealax_2Etreall_eq` : ι be given. Assume the following.

$$c_2Erealax_2Etreall_eq \in ((2^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)})^{ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal}) \tag{12}$$

Let `c_2Erealax_2Ereal_2ABS_2CLASS` : ι be given. Assume the following.

$$c_2Erealax_2Ereal_2ABS_2CLASS \in (ty_2Erealax_2Ereal)^{(2^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)})} \tag{13}$$

Definition 15 We define $c_2Erealax_2Ereal_ABS$ to be $\lambda V0r \in (ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)$

Definition 16 We define $c_2Erealax_2Einv$ to be $\lambda V0T1 \in ty_2Erealax_2Ereal.(ap\ c_2Erealax_2Ereal_ABS$

Let $c_2Erealax_2Etrealmul : \iota$ be given. Assume the following.

$$c_2Erealax_2Etrealmul \in (((ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)})^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)})^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)} \quad (14)$$

Definition 17 We define $c_2Erealax_2Ereal_mul$ to be $\lambda V0T1 \in ty_2Erealax_2Ereal.\lambda V1T2 \in ty_2Erealax_2Ereal.$

Definition 18 We define $c_2Ereal_2E_2F$ to be $\lambda V0x \in ty_2Erealax_2Ereal.\lambda V1y \in ty_2Erealax_2Ereal.$

Let $c_2Ereal_2Ereal_of_num : \iota$ be given. Assume the following.

$$c_2Ereal_2Ereal_of_num \in (ty_2Erealax_2Ereal^{ty_2Eenum_2Eenum}) \quad (15)$$

Let $c_2Erealax_2Etrealmult : \iota$ be given. Assume the following.

$$c_2Erealax_2Etrealmult \in ((2^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)}\ ty_2Ehreal_2Ehreal)^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)})^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)} \quad (16)$$

Definition 19 We define $c_2Erealax_2Ereal_lt$ to be $\lambda V0T1 \in ty_2Erealax_2Ereal.\lambda V1T2 \in ty_2Erealax_2Ereal.$

Definition 20 We define $c_2Ebool_2E_7E$ to be $(\lambda V0t \in 2.(ap\ (ap\ c_2Emin_2E_3D_3D_3E\ V0t)\ c_2Ebool_2E_7E$

Assume the following.

$$((ap\ c_2Earithmetic_2ENUMERAL\ (ap\ c_2Earithmetic_2EBIT2\ c_2Earithmetic_2EZERO)) = (ap\ c_2Eenum_2ESUC\ (ap\ c_2Earithmetic_2ENUMERAL\ (ap\ c_2Earithmetic_2EBIT1\ c_2Earithmetic_2EZERO)))) \quad (17)$$

Assume the following.

$$True \quad (18)$$

Assume the following.

$$(\forall V0t1 \in 2.(\forall V1t2 \in 2.(((p\ V0t1) \Rightarrow (p\ V1t2)) \Rightarrow (((p\ V1t2) \Rightarrow (p\ V0t1)) \Rightarrow ((p\ V0t1) \Leftrightarrow (p\ V1t2)))))) \quad (19)$$

Assume the following.

$$(\forall V0t \in 2.(False \Rightarrow (p\ V0t))) \quad (20)$$

Assume the following.

$$((\forall V0t \in 2.((\neg(\neg(p\ V0t))) \Leftrightarrow (p\ V0t))) \wedge (((\neg True) \Leftrightarrow False) \wedge ((\neg False) \Leftrightarrow True))) \quad (21)$$

Assume the following.

$$(\forall V0n \in ty_2Enum_2Enum. (\neg((ap\ c_2Enum_2ESUC\ V0n) = c_2Enum_2E0))) \quad (22)$$

Assume the following.

$$\begin{aligned} & (\forall V0n \in ty_2Enum_2Enum. (\forall V1d \in ty_2Erealax_2Ereal. \\ & ((\neg(V0n = c_2Enum_2E0)) \Rightarrow ((p\ (ap\ (ap\ c_2Erealax_2Ereal_lt\ (ap \\ & c_2Ereal_2Ereal_of_num\ c_2Enum_2E0))\ (ap\ (ap\ c_2Ereal_2E_2F \\ & V1d)\ (ap\ c_2Ereal_2Ereal_of_num\ V0n)))) \Leftrightarrow (p\ (ap\ (ap\ c_2Erealax_2Ereal_lt \\ & (ap\ c_2Ereal_2Ereal_of_num\ c_2Enum_2E0))\ V1d)))))) \end{aligned} \quad (23)$$

Theorem 1

$$\begin{aligned} & (\forall V0d \in ty_2Erealax_2Ereal. ((p\ (ap\ (ap\ c_2Erealax_2Ereal_lt \\ & (ap\ c_2Ereal_2Ereal_of_num\ c_2Enum_2E0))\ (ap\ (ap\ c_2Ereal_2E_2F \\ & V0d)\ (ap\ c_2Ereal_2Ereal_of_num\ (ap\ c_2Earithmetic_2ENUMERAL \\ & (ap\ c_2Earithmetic_2EBIT2\ c_2Earithmetic_2EZERO)))))) \Leftrightarrow (p\ (\\ & ap\ (ap\ c_2Erealax_2Ereal_lt\ (ap\ c_2Ereal_2Ereal_of_num\ c_2Enum_2E0)) \\ & V0d)))) \end{aligned}$$