

thm_2Ereal_2EREAL__LT__LMUL__IMP (TMXn- dAP2d4tpA4Laj6VCXG3LY9ZFiL9hSmg)

October 26, 2020

Definition 1 We define `c_2Emin_2E_3D` to be $\lambda A. \lambda x \in A. \lambda y \in A. \text{inj_o } (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define `c_2Ebool_2ET` to be $(\text{ap } (\text{ap } (\text{c_2Emin_2E_3D } (2^2))) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x)$

Definition 3 We define `c_2Ebool_2E_21` to be $\lambda A. 27a : \iota. (\lambda V0P \in (2^{A-27a}). (\text{ap } (\text{ap } (\text{c_2Emin_2E_3D } (2^{A-27a}))))$

Definition 4 We define `c_2Ebool_2EF` to be $(\text{ap } (\text{c_2Ebool_2E_21 } 2)) (\lambda V0t \in 2.V0t)$.

Definition 5 We define `c_2Emin_2E_3D_3D_3E` to be $\lambda P \in 2. \lambda Q \in 2. \text{inj_o } (p P \Rightarrow p Q)$ of type ι .

Definition 6 We define `c_2Ebool_2E_7E` to be $(\lambda V0t \in 2. (\text{ap } (\text{ap } (\text{c_2Emin_2E_3D_3D_3E } V0t) \text{ c_2Ebool_2EF})))$

Definition 7 We define `c_2Ebool_2E_2F_5C` to be $(\lambda V0t1 \in 2. (\lambda V1t2 \in 2. (\text{ap } (\text{c_2Ebool_2E_21 } 2)) (\lambda V2t \in 2.V2t)))$

Let `ty_2Ehreal_2Ehreal` : ι be given. Assume the following.

$$\text{nonempty } \text{ty_2Ehreal_2Ehreal} \tag{1}$$

Let `ty_2Epair_2Eprod` : $\iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0. \text{nonempty } A0 \Rightarrow \forall A1. \text{nonempty } A1 \Rightarrow \text{nonempty } (\text{ty_2Epair_2Eprod } A0 A1) \tag{2}$$

Let `ty_2Erealax_2Ereal` : ι be given. Assume the following.

$$\text{nonempty } \text{ty_2Erealax_2Ereal} \tag{3}$$

Let `c_2Erealax_2Ereal__REP__CLASS` : ι be given. Assume the following.

$$\text{c_2Erealax_2Ereal_REP_CLASS} \in ((2^{(\text{ty_2Epair_2Eprod } \text{ty_2Ehreal_2Ehreal } \text{ty_2Ehreal_2Ehreal}) \text{ty_2Erealax_2Ereal}})) \tag{4}$$

Definition 8 We define `c_2Emin_2E_40` to be $\lambda A. \lambda P \in 2^A. \text{if } (\exists x \in A. p (\text{ap } P x)) \text{ then } (\text{the } (\lambda x. x \in A \wedge p x))$ of type $\iota \Rightarrow \iota$.

Definition 9 We define $c_2Erealax_2Ereal_REP$ to be $\lambda V0a \in ty_2Erealax_2Ereal.(ap (c_2Emin_2E.40 (ty_2Erealax_2Ereal_mul : \iota$ be given. Assume the following.

$$c_2Erealax_2Ereal_mul \in (((ty_2Epair_2Eprod ty_2Ehreal_2Ehreal ty_2Ehreal_2Ehreal)(ty_2Epair_2Eprod ty_2Ehreal_2Ehreal ty_2Ehreal_2Ehreal)(ty_2Epair_2Eprod ty_2Ehreal_2Ehreal ty_2Ehreal_2Ehreal)) (5)$$

Let $c_2Erealax_2Ereal_eq : \iota$ be given. Assume the following.

$$c_2Erealax_2Ereal_eq \in ((2^{(ty_2Epair_2Eprod ty_2Ehreal_2Ehreal ty_2Ehreal_2Ehreal)})(ty_2Epair_2Eprod ty_2Ehreal_2Ehreal ty_2Ehreal_2Ehreal)) (6)$$

Let $c_2Erealax_2Ereal_ABS_CLASS : \iota$ be given. Assume the following.

$$c_2Erealax_2Ereal_ABS_CLASS \in (ty_2Erealax_2Ereal)^{(2^{(ty_2Epair_2Eprod ty_2Ehreal_2Ehreal ty_2Ehreal_2Ehreal)}} (7)$$

Definition 10 We define $c_2Erealax_2Ereal_ABS$ to be $\lambda V0r \in (ty_2Epair_2Eprod ty_2Ehreal_2Ehreal ty_2Ehreal_2Ehreal$

Definition 11 We define $c_2Erealax_2Ereal_mul$ to be $\lambda V0T1 \in ty_2Erealax_2Ereal.\lambda V1T2 \in ty_2Erealax_2Ereal$

Let $c_2Enum_2EZERO_REP : \iota$ be given. Assume the following.

$$c_2Enum_2EZERO_REP \in \omega (8)$$

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty ty_2Enum_2Enum (9)$$

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum)^{\omega} (10)$$

Definition 12 We define c_2Enum_2E0 to be $(ap c_2Enum_2EABS_num c_2Enum_2EZERO_REP)$.

Let $c_2Ereal_2Ereal_of_num : \iota$ be given. Assume the following.

$$c_2Ereal_2Ereal_of_num \in (ty_2Erealax_2Ereal)^{ty_2Enum_2Enum} (11)$$

Let $c_2Erealax_2Ereal_lt : \iota$ be given. Assume the following.

$$c_2Erealax_2Ereal_lt \in ((2^{(ty_2Epair_2Eprod ty_2Ehreal_2Ehreal ty_2Ehreal_2Ehreal)})(ty_2Epair_2Eprod ty_2Ehreal_2Ehreal ty_2Ehreal_2Ehreal)) (12)$$

Definition 13 We define $c_2Erealax_2Ereal_lt$ to be $\lambda V0T1 \in ty_2Erealax_2Ereal.\lambda V1T2 \in ty_2Erealax_2Ereal$

Assume the following.

$$True (13)$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2.(((True \Rightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Rightarrow True) \Leftrightarrow \\ & True) \wedge (((False \Rightarrow (p V0t)) \Leftrightarrow True) \wedge (((p V0t) \Rightarrow (p V0t)) \Leftrightarrow True) \wedge ((\\ & (p V0t) \Rightarrow False) \Leftrightarrow (\neg (p V0t)))))) (14) \end{aligned}$$

Assume the following.

$$\begin{aligned}
& (\forall V0x \in ty_2Erealax_2Ereal. (\forall V1y \in ty_2Erealax_2Ereal. \\
& (\forall V2z \in ty_2Erealax_2Ereal. ((p (ap (ap c_2Erealax_2Ereal_lt \\
& (ap c_2Ereal_2Ereal_of_num c_2Enum_2E0)) V0x)) \Rightarrow ((p (ap (ap \\
& c_2Erealax_2Ereal_lt (ap (ap c_2Erealax_2Ereal_mul V0x) V1y)) \\
& (ap (ap c_2Erealax_2Ereal_mul V0x) V2z))) \Leftrightarrow (p (ap (ap c_2Erealax_2Ereal_lt \\
& V1y) V2z)))))))))
\end{aligned} \tag{15}$$

Theorem 1

$$\begin{aligned}
& (\forall V0x \in ty_2Erealax_2Ereal. (\forall V1y \in ty_2Erealax_2Ereal. \\
& (\forall V2z \in ty_2Erealax_2Ereal. (((p (ap (ap c_2Erealax_2Ereal_lt \\
V1y) V2z)) \wedge (p (ap (ap c_2Erealax_2Ereal_lt (ap c_2Ereal_2Ereal_of_num \\
c_2Enum_2E0)) V0x))) \Rightarrow (p (ap (ap c_2Erealax_2Ereal_lt (ap (ap \\
c_2Erealax_2Ereal_mul V0x) V1y)) (ap (ap c_2Erealax_2Ereal_mul \\
V0x) V2z)))))))))
\end{aligned}$$