

# thm\_2Ereal\_2EREAL\_LT\_RMUL\_IMP (TM-SwhJK2DuxJgXM9sKpC9Fzt2a2Xxwq3Kb7)

October 26, 2020

**Definition 1** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define  $c\_2Ebool\_2E\_2T$  to be  $(ap (ap (c\_2Emin\_2E\_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

**Definition 3** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A.\lambda a : \iota.(\lambda V0P \in (2^{A-27a}).(ap (ap (c\_2Emin\_2E\_3D (2^{A-27a}))$

**Definition 4** We define  $c\_2Ebool\_2E\_2F$  to be  $(ap (c\_2Ebool\_2E\_21 2) (\lambda V0t \in 2.V0t))$ .

**Definition 5** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o (p P \Rightarrow p Q)$  of type  $\iota$ .

**Definition 6** We define  $c\_2Ebool\_2E\_27E$  to be  $(\lambda V0t \in 2.(ap (ap c\_2Emin\_2E\_3D\_3D\_3E V0t) c\_2Ebool\_2E\_2F))$

**Definition 7** We define  $c\_2Ebool\_2E\_2F\_5C$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_2Ebool\_2E\_21 2) (\lambda V2t \in 2.V2t)))$

Let  $ty\_2Ehreal\_2Ehreal : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Ehreal\_2Ehreal \tag{1}$$

Let  $ty\_2Epair\_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty\_2Epair\_2Eprod\ A0\ A1) \tag{2}$$

Let  $ty\_2Erealax\_2Ereal : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Erealax\_2Ereal \tag{3}$$

Let  $c\_2Erealax\_2Ereal\_REP\_CLASS : \iota$  be given. Assume the following.

$$c\_2Erealax\_2Ereal\_REP\_CLASS \in ((2^{(ty\_2Epair\_2Eprod\ ty\_2Ehreal\_2Ehreal\ ty\_2Ehreal\_2Ehreal)})\ ty\_2Erealax\_2Ereal) \tag{4}$$

**Definition 8** We define  $c\_2Emin\_2E\_40$  to be  $\lambda A.\lambda P \in 2^A.if (\exists x \in A.p (ap P x)) \mathbf{then} (the (\lambda x.x \in A \wedge p$  of type  $\iota \Rightarrow \iota$ .

**Definition 9** We define  $c\_2Erealax\_2Ereal\_REP$  to be  $\lambda V0a \in ty\_2Erealax\_2Ereal.(ap (c\_2Emin\_2E.40 (ty\_2Erealax\_2Ereal\_mul) a))$ .  
Let  $c\_2Erealax\_2Ereal\_mul : \iota$  be given. Assume the following.

$$c\_2Erealax\_2Ereal\_mul \in (((ty\_2Epair\_2Eprod ty\_2Ehreal\_2Ehreal ty\_2Ehreal\_2Ehreal) (ty\_2Epair\_2Eprod ty\_2Ehreal\_2Ehreal ty\_2Ehreal\_2Ehreal)) (ty\_2Epair\_2Eprod ty\_2Ehreal\_2Ehreal ty\_2Ehreal\_2Ehreal)) (ty\_2Epair\_2Eprod ty\_2Ehreal\_2Ehreal ty\_2Ehreal\_2Ehreal) \quad (5)$$

Let  $c\_2Erealax\_2Ereal\_eq : \iota$  be given. Assume the following.

$$c\_2Erealax\_2Ereal\_eq \in ((2^{(ty\_2Epair\_2Eprod ty\_2Ehreal\_2Ehreal ty\_2Ehreal\_2Ehreal)}) (ty\_2Epair\_2Eprod ty\_2Ehreal\_2Ehreal ty\_2Ehreal\_2Ehreal)) \quad (6)$$

Let  $c\_2Erealax\_2Ereal\_ABS\_CLASS : \iota$  be given. Assume the following.

$$c\_2Erealax\_2Ereal\_ABS\_CLASS \in (ty\_2Erealax\_2Ereal)^{(2^{(ty\_2Epair\_2Eprod ty\_2Ehreal\_2Ehreal ty\_2Ehreal\_2Ehreal)})} \quad (7)$$

**Definition 10** We define  $c\_2Erealax\_2Ereal\_ABS$  to be  $\lambda V0r \in (ty\_2Epair\_2Eprod ty\_2Ehreal\_2Ehreal ty\_2Ehreal\_2Ehreal)$ .

**Definition 11** We define  $c\_2Erealax\_2Ereal\_mul$  to be  $\lambda V0T1 \in ty\_2Erealax\_2Ereal.\lambda V1T2 \in ty\_2Erealax\_2Ereal$ .

Let  $c\_2Enum\_2EZERO\_REP : \iota$  be given. Assume the following.

$$c\_2Enum\_2EZERO\_REP \in \omega \quad (8)$$

Let  $ty\_2Enum\_2Enum : \iota$  be given. Assume the following.

$$\text{nonempty } ty\_2Enum\_2Enum \quad (9)$$

Let  $c\_2Enum\_2EABS\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EABS\_num \in (ty\_2Enum\_2Enum)^{\omega} \quad (10)$$

**Definition 12** We define  $c\_2Enum\_2E0$  to be  $(ap c\_2Enum\_2EABS\_num c\_2Enum\_2EZERO\_REP)$ .

Let  $c\_2Ereal\_2Ereal\_of\_num : \iota$  be given. Assume the following.

$$c\_2Ereal\_2Ereal\_of\_num \in (ty\_2Erealax\_2Ereal)^{ty\_2Enum\_2Enum} \quad (11)$$

Let  $c\_2Erealax\_2Ereal\_lt : \iota$  be given. Assume the following.

$$c\_2Erealax\_2Ereal\_lt \in ((2^{(ty\_2Epair\_2Eprod ty\_2Ehreal\_2Ehreal ty\_2Ehreal\_2Ehreal)}) (ty\_2Epair\_2Eprod ty\_2Ehreal\_2Ehreal ty\_2Ehreal\_2Ehreal)) \quad (12)$$

**Definition 13** We define  $c\_2Erealax\_2Ereal\_lt$  to be  $\lambda V0T1 \in ty\_2Erealax\_2Ereal.\lambda V1T2 \in ty\_2Erealax\_2Ereal$ .

Assume the following.

$$True \quad (13)$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2.(((True \Rightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Rightarrow True) \Leftrightarrow \\ & True) \wedge (((False \Rightarrow (p V0t)) \Leftrightarrow True) \wedge (((p V0t) \Rightarrow (p V0t)) \Leftrightarrow True) \wedge (( \\ & (p V0t) \Rightarrow False) \Leftrightarrow (\neg (p V0t)))))) \end{aligned} \quad (14)$$

Assume the following.

$$\begin{aligned}
& (\forall V0x \in ty\_2Erealax\_2Ereal. (\forall V1y \in ty\_2Erealax\_2Ereal. \\
& (\forall V2z \in ty\_2Erealax\_2Ereal. ((p (ap (ap c\_2Erealax\_2Ereal\_lt \\
& (ap c\_2Ereal\_2Ereal\_of\_num c\_2Enum\_2E0)) V2z)) \Rightarrow ((p (ap (ap \\
& c\_2Erealax\_2Ereal\_lt (ap (ap c\_2Erealax\_2Ereal\_mul V0x) V2z)) \\
& (ap (ap c\_2Erealax\_2Ereal\_mul V1y) V2z))) \Leftrightarrow (p (ap (ap c\_2Erealax\_2Ereal\_lt \\
& V0x) V1y)))))))))
\end{aligned} \tag{15}$$

**Theorem 1**

$$\begin{aligned}
& (\forall V0x \in ty\_2Erealax\_2Ereal. (\forall V1y \in ty\_2Erealax\_2Ereal. \\
& (\forall V2z \in ty\_2Erealax\_2Ereal. (((p (ap (ap c\_2Erealax\_2Ereal\_lt \\
V0x) V1y)) \wedge (p (ap (ap c\_2Erealax\_2Ereal\_lt (ap c\_2Ereal\_2Ereal\_of\_num \\
c\_2Enum\_2E0)) V2z))) \Rightarrow (p (ap (ap c\_2Erealax\_2Ereal\_lt (ap (ap \\
c\_2Erealax\_2Ereal\_mul V0x) V2z)) (ap (ap c\_2Erealax\_2Ereal\_mul \\
V1y) V2z)))))))))
\end{aligned}$$