

thm_2Ereal_2EREAL__LT__TRANS (TM- NeaVD1FXUSQR1EmfsVMuBK6K4Pb9GPbxw)

October 26, 2020

Let $ty_2Ehreal_2Ehreal : \iota$ be given. Assume the following.

$$nonempty\ ty_2Ehreal_2Ehreal \tag{1}$$

Let $ty_2Epair_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty_2Epair_2Eprod\ A0\ A1) \tag{2}$$

Let $ty_2Erealax_2Ereal : \iota$ be given. Assume the following.

$$nonempty\ ty_2Erealax_2Ereal \tag{3}$$

Let $c_2Erealax_2Ereal_REP_CLASS : \iota$ be given. Assume the following.

$$c_2Erealax_2Ereal_REP_CLASS \in ((2^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)})\ ty_2Erealax_2Ereal) \tag{4}$$

Definition 1 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A$.if $(\exists x \in A.p\ (ap\ P\ x))$ **then** (the $(\lambda x.x \in A \wedge p\ x)$ of type $\iota \Rightarrow \iota$).

Definition 2 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o\ (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 3 We define $c_2Ebool_2E_21$ to be $(ap\ (ap\ (c_2Emin_2E_3D\ (2^2)))\ (\lambda V0x \in 2.V0x))\ (\lambda V1x \in 2.V1x)$

Definition 4 We define $c_2Ebool_2E_21$ to be $\lambda A.\lambda P \in (2^{A-27a}).(ap\ (ap\ (c_2Emin_2E_3D\ (2^{A-27a})))\ P)$

Definition 5 We define $c_2Erealax_2Ereal_REP$ to be $\lambda V0a \in ty_2Erealax_2Ereal.(ap\ (c_2Emin_2E_40\ (ty_2Erealax_2Ereal_REP_CLASS\ a)))$

Let $c_2Erealax_2Etrealt_lt : \iota$ be given. Assume the following.

$$c_2Erealax_2Etrealt_lt \in ((2^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)})\ (ty_2Epair_2Eprod\ ty_2Erealax_2Ereal)) \tag{5}$$

Definition 6 We define $c_2Erealax_2Ereal_lt$ to be $\lambda V0T1 \in ty_2Erealax_2Ereal.\lambda V1T2 \in ty_2Erealax_2Ereal.c_2Etrealt_lt\ T1\ T2$

Definition 7 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow p Q)$ of type ι .

Definition 8 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.$

Assume the following.

$$\begin{aligned}
& (\forall V0x \in ty_2Erealax_2Ereal.(\forall V1y \in ty_2Erealax_2Ereal. \\
& (\forall V2z \in ty_2Erealax_2Ereal.(((p (ap (ap c_2Erealax_2Ereal_lt \\
& V0x) V1y)) \wedge (p (ap (ap c_2Erealax_2Ereal_lt V1y) V2z))) \Rightarrow (p (ap \\
& (ap c_2Erealax_2Ereal_lt V0x) V2z))))))
\end{aligned} \tag{6}$$

Theorem 1

$$\begin{aligned}
& (\forall V0x \in ty_2Erealax_2Ereal.(\forall V1y \in ty_2Erealax_2Ereal. \\
& (\forall V2z \in ty_2Erealax_2Ereal.(((p (ap (ap c_2Erealax_2Ereal_lt \\
& V0x) V1y)) \wedge (p (ap (ap c_2Erealax_2Ereal_lt V1y) V2z))) \Rightarrow (p (ap \\
& (ap c_2Erealax_2Ereal_lt V0x) V2z))))))
\end{aligned}$$