

thm\_2Ereal\_2EREAL\_\_MIN\_\_LE1  
(TMKxop61KPwbL6YDGYw6cZqPXwMJb81MGtB)

October 26, 2020

**Definition 1** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define  $c\_2Ebool\_2E\_2T$  to be  $(ap (ap (c\_2Emin\_2E\_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

**Definition 3** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A\_27a : \iota.(\lambda V0P \in (2^{A\_27a}).(ap (ap (c\_2Emin\_2E\_3D (2^{A\_27a})) (\lambda V1P \in 2.V1P)) (\lambda V2P \in 2.V2P)))$

**Definition 4** We define  $c\_2Ebool\_2E\_2F$  to be  $(ap (c\_2Ebool\_2E\_21 2) (\lambda V0t \in 2.V0t))$ .

**Definition 5** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o (p P \Rightarrow p Q)$  of type  $\iota$ .

**Definition 6** We define  $c\_2Ebool\_2E\_27E$  to be  $(\lambda V0t \in 2.(ap (ap c\_2Emin\_2E\_3D\_3D\_3E V0t) c\_2Ebool\_2E\_2F))$

**Definition 7** We define  $c\_2Ebool\_2E\_2F\_5C$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_2Ebool\_2E\_21 2) (\lambda V2t \in 2.V2t)))$

**Definition 8** We define  $c\_2Ebool\_2E\_5C\_2F$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_2Ebool\_2E\_21 2) (\lambda V2t \in 2.V2t)))$

Let  $ty\_2Ehreal\_2Ehreal : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Ehreal\_2Ehreal \tag{1}$$

Let  $ty\_2Epair\_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty\_2Epair\_2Eprod\ A0\ A1) \tag{2}$$

Let  $ty\_2Erealax\_2Ereal : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Erealax\_2Ereal \tag{3}$$

Let  $c\_2Erealax\_2Ereal\_REP\_CLASS : \iota$  be given. Assume the following.

$$c\_2Erealax\_2Ereal\_REP\_CLASS \in ((2^{(ty\_2Epair\_2Eprod\ ty\_2Ehreal\_2Ehreal\ ty\_2Ehreal\_2Ehreal)}\ ty\_2Erealax\_2Ereal)) \tag{4}$$

**Definition 9** We define  $c\_2Emin\_2E.40$  to be  $\lambda A.\lambda P \in 2^A.$ if  $(\exists x \in A.p (ap P x))$  then (the  $(\lambda x.x \in A \wedge p$  of type  $\iota \Rightarrow \iota$ .

**Definition 10** We define  $c\_2Erealax\_2Ereal\_REP$  to be  $\lambda V0a \in ty\_2Erealax\_2Ereal.(ap (c\_2Emin\_2E.40 ($

Let  $c\_2Erealax\_2Ereal\_lt : \iota$  be given. Assume the following.

$$c\_2Erealax\_2Ereal\_lt \in ((2^{(ty\_2Epair\_2Eprod ty\_2Ehreal\_2Ehreal ty\_2Ehreal\_2Ehreal)})(ty\_2Epair\_2Eprod ty\_2Ehreal\_2Ehreal)) \quad (5)$$

**Definition 11** We define  $c\_2Erealax\_2Ereal\_lt$  to be  $\lambda V0T1 \in ty\_2Erealax\_2Ereal.\lambda V1T2 \in ty\_2Erealax\_2Ereal.$

**Definition 12** We define  $c\_2Ereal\_2Ereal\_lte$  to be  $\lambda V0x \in ty\_2Erealax\_2Ereal.\lambda V1y \in ty\_2Erealax\_2Ereal.$

**Definition 13** We define  $c\_2Ebool\_2ECOND$  to be  $\lambda A.27a : \iota.(\lambda V0t \in 2.(\lambda V1t1 \in A.27a.(\lambda V2t2 \in A.27a.($

**Definition 14** We define  $c\_2Ereal\_2Emin$  to be  $\lambda V0x \in ty\_2Erealax\_2Ereal.\lambda V1y \in ty\_2Erealax\_2Ereal.$

Assume the following.

$$True \quad (6)$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2.(((True \vee (p V0t)) \Leftrightarrow True) \wedge (((p V0t) \vee True) \Leftrightarrow True) \wedge \\ & (((False \vee (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \vee False) \Leftrightarrow (p V0t)) \wedge (((p V0t) \vee \\ & (p V0t)) \Leftrightarrow (p V0t)))))) \end{aligned} \quad (7)$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2.(((True \Leftrightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Leftrightarrow True) \Leftrightarrow \\ & (p V0t)) \wedge (((False \Leftrightarrow (p V0t)) \Leftrightarrow \neg(p V0t)) \wedge (((p V0t) \Leftrightarrow False) \Leftrightarrow \neg( \\ & p V0t)))))) \end{aligned} \quad (8)$$

Assume the following.

$$(\forall V0x \in ty\_2Erealax\_2Ereal.(p (ap (ap c\_2Ereal\_2Ereal\_lte V0x) V0x))) \quad (9)$$

Assume the following.

$$\begin{aligned} & (\forall V0z \in ty\_2Erealax\_2Ereal.(\forall V1x \in ty\_2Erealax\_2Ereal. \\ & (\forall V2y \in ty\_2Erealax\_2Ereal.((p (ap (ap c\_2Ereal\_2Ereal\_lte \\ & (ap (ap c\_2Ereal\_2Emin V1x) V2y)) V0z)) \Leftrightarrow ((p (ap (ap c\_2Ereal\_2Ereal\_lte \\ & V1x) V0z)) \vee (p (ap (ap c\_2Ereal\_2Ereal\_lte V2y) V0z))))))))) \end{aligned} \quad (10)$$

**Theorem 1**

$$\begin{aligned} & (\forall V0x \in ty\_2Erealax\_2Ereal.(\forall V1y \in ty\_2Erealax\_2Ereal. \\ & (p (ap (ap c\_2Ereal\_2Ereal\_lte (ap (ap c\_2Ereal\_2Emin V0x) V1y)) \\ & V0x)))) \end{aligned}$$