

thm_2Ereal_2EREAL_NEG_RMUL
(TMGkoiGLvgUKt8azN6wT7BWdQhV6N9kGo3G)

October 26, 2020

Let $ty_2Ehreal_2Ehreal : \iota$ be given. Assume the following.

$$nonempty\ ty_2Ehreal_2Ehreal \tag{1}$$

Let $ty_2Epair_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty_2Epair_2Eprod\ A0\ A1) \tag{2}$$

Let $ty_2Erealax_2Ereal : \iota$ be given. Assume the following.

$$nonempty\ ty_2Erealax_2Ereal \tag{3}$$

Let $c_2Erealax_2Ereal_REP_CLASS : \iota$ be given. Assume the following.

$$c_2Erealax_2Ereal_REP_CLASS \in ((2^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)})_{ty_2Erealax_2Ereal}) \tag{4}$$

Definition 1 We define c_2Emin_2E40 to be $\lambda A.\lambda P \in 2^A$.if $(\exists x \in A.p\ (ap\ P\ x))$ then (the $(\lambda x.x \in A \wedge p\ x)$ of type $\iota \Rightarrow \iota$).

Definition 2 We define c_2Emin_2E3D to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o\ (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 3 We define c_2Ebool_2E27 to be $(ap\ (ap\ (c_2Emin_2E3D\ (2^2))\ (\lambda V0x \in 2.V0x))\ (\lambda V1x \in 2.V1x))$

Definition 4 We define c_2Ebool_2E21 to be $\lambda A.27a : \iota.(\lambda V0P \in (2^{A-27a}).(ap\ (ap\ (c_2Emin_2E3D\ (2^{A-27a}))\ (\lambda V1x \in 2.V1x))\ (\lambda V2x \in 2.V2x))$

Definition 5 We define $c_2Erealax_2Ereal_REP$ to be $\lambda V0a \in ty_2Erealax_2Ereal.(ap\ (c_2Emin_2E40\ (ty_2Erealax_2Ereal\ a)))$

Let $c_2Erealax_2Etrealmul : \iota$ be given. Assume the following.

$$c_2Erealax_2Etrealmul \in (((ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)_{ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal})_{ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal})_{ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal} \tag{5}$$

Let $c_2Erealax_2Etrealm_eq : \iota$ be given. Assume the following.

$$c_2Erealax_2Etrealm_eq \in ((2^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)})^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal)}) \quad (6)$$

Let $c_2Erealax_2Ereal_ABS_CLASS : \iota$ be given. Assume the following.

$$c_2Erealax_2Ereal_ABS_CLASS \in (ty_2Erealax_2Ereal)^{(2^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)})} \quad (7)$$

Definition 6 We define $c_2Erealax_2Ereal_ABS$ to be $\lambda V0r \in (ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)$

Definition 7 We define $c_2Erealax_2Ereal_mul$ to be $\lambda V0T1 \in ty_2Erealax_2Ereal.\lambda V1T2 \in ty_2Erealax_2Ereal$

Let $c_2Erealax_2Etrealm_neg : \iota$ be given. Assume the following.

$$c_2Erealax_2Etrealm_neg \in ((ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)}) \quad (8)$$

Definition 8 We define $c_2Erealax_2Ereal_neg$ to be $\lambda V0T1 \in ty_2Erealax_2Ereal.(ap\ c_2Erealax_2Ereal_mul$

Assume the following.

$$(\forall V0x \in ty_2Erealax_2Ereal.(\forall V1y \in ty_2Erealax_2Ereal. ((ap\ (ap\ c_2Erealax_2Ereal_mul\ V0x)\ V1y) = (ap\ (ap\ c_2Erealax_2Ereal_mul\ V1y)\ V0x)))) \quad (9)$$

Assume the following.

$$(\forall V0x \in ty_2Erealax_2Ereal.(\forall V1y \in ty_2Erealax_2Ereal. ((ap\ c_2Erealax_2Ereal_neg\ (ap\ (ap\ c_2Erealax_2Ereal_mul\ V0x)\ V1y)) = (ap\ (ap\ c_2Erealax_2Ereal_mul\ (ap\ c_2Erealax_2Ereal_neg\ V0x))\ V1y)))) \quad (10)$$

Theorem 1

$$(\forall V0x \in ty_2Erealax_2Ereal.(\forall V1y \in ty_2Erealax_2Ereal. ((ap\ c_2Erealax_2Ereal_neg\ (ap\ (ap\ c_2Erealax_2Ereal_mul\ V0x)\ V1y)) = (ap\ (ap\ c_2Erealax_2Ereal_mul\ V0x)\ (ap\ c_2Erealax_2Ereal_neg\ V1y))))))$$