

thm_2Ereal_2EREAL__POS__ID (TMdKvr- WVZ8B72fc8Y799ALRD5PLtmysRJ9F)

October 26, 2020

Definition 1 We define `c_2Emin_2E_3D` to be $\lambda A. \lambda x \in A. \lambda y \in A. \text{inj_o } (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define `c_2Ebool_2E_7E` to be $(\text{ap } (\text{ap } (\text{c_2Emin_2E_3D } (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 3 We define `c_2Ebool_2E_21` to be $\lambda A. 27a : \iota. (\lambda V0P \in (2^{A-27a}). (\text{ap } (\text{ap } (\text{c_2Emin_2E_3D } (2^{A-27a}))))$

Definition 4 We define `c_2Ebool_2E_2F` to be $(\text{ap } (\text{c_2Ebool_2E_21 } 2) (\lambda V0t \in 2.V0t))$.

Definition 5 We define `c_2Emin_2E_3D_3D_3E` to be $\lambda P \in 2. \lambda Q \in 2. \text{inj_o } (p \Rightarrow q)$ of type ι .

Definition 6 We define `c_2Ebool_2E_7E` to be $(\lambda V0t \in 2. (\text{ap } (\text{ap } (\text{c_2Emin_2E_3D_3D_3E } V0t) (\text{c_2Ebool_2E_2F}))))$

Definition 7 We define `c_2Ebool_2E_2F_5C` to be $(\lambda V0t1 \in 2. (\lambda V1t2 \in 2. (\text{ap } (\text{c_2Ebool_2E_21 } 2) (\lambda V2t \in 2.V2t))))$

Let `c_2Enum_2EZERO__REP` : ι be given. Assume the following.

$$c_2Enum_2EZERO_REP \in \text{omega} \tag{1}$$

Let `ty_2Enum_2Enum` : ι be given. Assume the following.

$$\text{nonempty } ty_2Enum_2Enum \tag{2}$$

Let `c_2Enum_2EABS__num` : ι be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum^{\text{omega}}) \tag{3}$$

Definition 8 We define `c_2Enum_2E0` to be $(\text{ap } (\text{c_2Enum_2EABS_num } c_2Enum_2EZERO_REP))$.

Let `ty_2Erealx_2Ereal` : ι be given. Assume the following.

$$\text{nonempty } ty_2Erealx_2Ereal \tag{4}$$

Let `c_2Ereal_2Ereal_of__num` : ι be given. Assume the following.

$$c_2Ereal_2Ereal_of_num \in (ty_2Erealx_2Ereal^{ty_2Enum_2Enum}) \tag{5}$$

Let $ty_2Ehreal_2Ehreal : \iota$ be given. Assume the following.

$$nonempty\ ty_2Ehreal_2Ehreal \quad (6)$$

Let $ty_2Epair_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty_2Epair_2Eprod\ A0\ A1) \quad (7)$$

Let $c_2Erealax_2Ereal_REP_CLASS : \iota$ be given. Assume the following.

$$c_2Erealax_2Ereal_REP_CLASS \in ((2^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)})^{ty_2Erealax_2Ereal_REP_CLASS}) \quad (8)$$

Definition 9 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A.$ if $(\exists x \in A.p\ (ap\ P\ x))$ then (the $(\lambda x.x \in A \wedge p\ x)$ of type $\iota \Rightarrow \iota$).

Definition 10 We define $c_2Erealax_2Ereal_REP$ to be $\lambda V0a \in ty_2Erealax_2Ereal.(ap\ (c_2Emin_2E_40\ a))$

Let $c_2Erealax_2Etreall_lt : \iota$ be given. Assume the following.

$$c_2Erealax_2Etreall_lt \in ((2^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)})^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal)}) \quad (9)$$

Definition 11 We define $c_2Erealax_2Ereal_lt$ to be $\lambda V0T1 \in ty_2Erealax_2Ereal.\lambda V1T2 \in ty_2Erealax_2Ereal.$

Definition 12 We define $c_2Ereal_2Ereal_lte$ to be $\lambda V0x \in ty_2Erealax_2Ereal.\lambda V1y \in ty_2Erealax_2Ereal.$

Definition 13 We define c_2Ebool_2ECOND to be $\lambda A_27a : \iota.(\lambda V0t \in 2.(\lambda V1t1 \in A_27a.(\lambda V2t2 \in A_27a.($

Definition 14 We define c_2Ereal_2Epos to be $\lambda V0x \in ty_2Erealax_2Ereal.(ap\ (ap\ (ap\ (c_2Ebool_2ECOND\ x))))$

Assume the following.

$$True \quad (10)$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2.(((True \Rightarrow (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \Rightarrow True) \Leftrightarrow \\ & True) \wedge (((False \Rightarrow (p\ V0t)) \Leftrightarrow True) \wedge (((p\ V0t) \Rightarrow (p\ V0t)) \Leftrightarrow True) \wedge ((\\ & (p\ V0t) \Rightarrow False) \Leftrightarrow (\neg (p\ V0t)))))) \end{aligned} \quad (11)$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0x \in A_27a.((V0x = V0x) \Leftrightarrow True)) \quad (12)$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0x \in A_27a.(\forall V1y \in A_27a.((V0x = V1y) \Leftrightarrow (V1y = V0x)))) \quad (13)$$

Assume the following.

$$\begin{aligned}
& (\forall V0t \in 2.(((True \Leftrightarrow (p \ V0t)) \Leftrightarrow (p \ V0t)) \wedge (((p \ V0t) \Leftrightarrow True) \Leftrightarrow \\
& (p \ V0t)) \wedge (((False \Leftrightarrow (p \ V0t)) \Leftrightarrow \neg(p \ V0t)) \wedge (((p \ V0t) \Leftrightarrow False) \Leftrightarrow \neg(\\
& p \ V0t))))))
\end{aligned} \tag{14}$$

Assume the following.

$$\begin{aligned}
& (\forall V0t1 \in 2.(\forall V1t2 \in 2.(\forall V2t3 \in 2.(((p \ V0t1) \Rightarrow \\
& ((p \ V1t2) \Rightarrow (p \ V2t3))) \Leftrightarrow (((p \ V0t1) \wedge (p \ V1t2)) \Rightarrow (p \ V2t3))))))
\end{aligned} \tag{15}$$

Assume the following.

$$\begin{aligned}
& (\forall V0x \in 2.(\forall V1x_27 \in 2.(\forall V2y \in 2.(\forall V3y_27 \in \\
& 2.(((p \ V0x) \Leftrightarrow (p \ V1x_27)) \wedge ((p \ V1x_27) \Rightarrow ((p \ V2y) \Leftrightarrow (p \ V3y_27)))) \Rightarrow \\
& ((p \ V0x) \Rightarrow (p \ V2y)) \Leftrightarrow ((p \ V1x_27) \Rightarrow (p \ V3y_27))))))
\end{aligned} \tag{16}$$

Assume the following.

$$\begin{aligned}
& \forall A_27a.nonempty \ A_27a \Rightarrow (\forall V0P \in 2.(\forall V1Q \in 2. \\
& (\forall V2x \in A_27a.(\forall V3x_27 \in A_27a.(\forall V4y \in A_27a. \\
& (\forall V5y_27 \in A_27a.(((p \ V0P) \Leftrightarrow (p \ V1Q)) \wedge ((p \ V1Q) \Rightarrow (V2x = V3x_27)) \wedge \\
& ((\neg(p \ V1Q)) \Rightarrow (V4y = V5y_27)))) \Rightarrow ((ap \ (ap \ (ap \ (c_2Ebool_2ECOND \ A_27a) \\
& V0P) \ V2x) \ V4y) = (ap \ (ap \ (ap \ (c_2Ebool_2ECOND \ A_27a) \ V1Q) \ V3x_27) \\
& V5y_27)))))))))
\end{aligned} \tag{17}$$

Assume the following.

$$\begin{aligned}
& \forall A_27a.nonempty \ A_27a \Rightarrow ((\forall V0t1 \in A_27a.(\forall V1t2 \in \\
& A_27a.((ap \ (ap \ (ap \ (c_2Ebool_2ECOND \ A_27a) \ c_2Ebool_2ET) \ V0t1) \\
& V1t2) = V0t1))) \wedge (\forall V2t1 \in A_27a.(\forall V3t2 \in A_27a.((ap \\
& (ap \ (ap \ (c_2Ebool_2ECOND \ A_27a) \ c_2Ebool_2EF) \ V2t1) \ V3t2) = V3t2))))))
\end{aligned} \tag{18}$$

Theorem 1

$$\begin{aligned}
& (\forall V0x \in ty_2Erealax_2Ereal.((p \ (ap \ (ap \ c_2Ereal_2Ereal_lte \\
& (ap \ c_2Ereal_2Ereal_of_num \ c_2Enum_2E0) \ V0x)) \Rightarrow ((ap \ c_2Ereal_2Epos \\
& V0x) = V0x)))
\end{aligned}$$