

thm_2Ereal_2EREAL__SUB__LE
(TMS9o9EuDTnR93RHayGS7wrbEodCZHmm7Sg)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define c_2Ebool_2ET to be $(ap (ap (c_2Emin_2E_3D (2^2))) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x)$

Let $c_2Enum_2EZERO_REP : \iota$ be given. Assume the following.

$$c_2Enum_2EZERO_REP \in \omega \tag{1}$$

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \tag{2}$$

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum^{\omega}) \tag{3}$$

Definition 3 We define c_2Enum_2E0 to be $(ap\ c_2Enum_2EABS_num\ c_2Enum_2EZERO_REP)$.

Let $ty_2Erealx_2Ereal : \iota$ be given. Assume the following.

$$nonempty\ ty_2Erealx_2Ereal \tag{4}$$

Let $c_2Ereal_2Ereal_of_num : \iota$ be given. Assume the following.

$$c_2Ereal_2Ereal_of_num \in (ty_2Erealx_2Ereal^{ty_2Enum_2Enum}) \tag{5}$$

Let $ty_2Ehreal_2Ehreal : \iota$ be given. Assume the following.

$$nonempty\ ty_2Ehreal_2Ehreal \tag{6}$$

Let $ty_2Epair_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty_2Epair_2Eprod\ A0\ A1) \tag{7}$$

Let $c_2Erealx_2Ereal_REP_CLASS : \iota$ be given. Assume the following.

$$c_2Erealx_2Ereal_REP_CLASS \in ((2^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)})^{ty_2Erealx_2Ereal}) \tag{8}$$

Definition 4 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A.$ **if** $(\exists x \in A.p (ap P x))$ **then** *(the $(\lambda x.x \in A \wedge p$ of type $\iota \Rightarrow \iota$).*

Definition 5 We define $c_2Ebool_2E_21$ to be $\lambda A.27a : \iota.(\lambda V0P \in (2^{A-27a}).(ap (ap (c_2Emin_2E_3D (2^{A-27a}$

Definition 6 We define $c_2Erealax_2Ereal_REP$ to be $\lambda V0a \in ty_2Erealax_2Ereal.(ap (c_2Emin_2E_40 (ty$

Let $c_2Erealax_2Etrealm_lt : \iota$ be given. Assume the following.

$$c_2Erealax_2Etrealm_lt \in ((2^{(ty_2Epair_2Eprod ty_2Ehreal_2Ehreal ty_2Ehreal_2Ehreal)})(ty_2Epair_2Eprod ty_2Ehreal_2Ehreal)) \quad (9)$$

Definition 7 We define $c_2Erealax_2Ereal_lt$ to be $\lambda V0T1 \in ty_2Erealax_2Ereal.\lambda V1T2 \in ty_2Erealax_2E$

Definition 8 We define $c_2Ebool_2E_2F$ to be $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$.

Definition 9 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow p Q)$ of type ι .

Definition 10 We define $c_2Ebool_2E_7E$ to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2E_2F$

Definition 11 We define $c_2Ereal_2Ereal_lte$ to be $\lambda V0x \in ty_2Erealax_2Ereal.\lambda V1y \in ty_2Erealax_2E$

Let $c_2Erealax_2Etrealm_neg : \iota$ be given. Assume the following.

$$c_2Erealax_2Etrealm_neg \in ((ty_2Epair_2Eprod ty_2Ehreal_2Ehreal ty_2Ehreal_2Ehreal)(ty_2Epair_2Eprod ty_2Ehreal_2Ehreal ty_2Ehreal_2Ehreal)) \quad (10)$$

Let $c_2Erealax_2Etrealm_eq : \iota$ be given. Assume the following.

$$c_2Erealax_2Etrealm_eq \in ((2^{(ty_2Epair_2Eprod ty_2Ehreal_2Ehreal ty_2Ehreal_2Ehreal)})(ty_2Epair_2Eprod ty_2Ehreal_2Ehreal)) \quad (11)$$

Let $c_2Erealax_2Ereal_ABS_CLASS : \iota$ be given. Assume the following.

$$c_2Erealax_2Ereal_ABS_CLASS \in (ty_2Erealax_2Ereal)^{(2^{(ty_2Epair_2Eprod ty_2Ehreal_2Ehreal ty_2Ehreal_2Ehreal)})(ty_2Epair_2Eprod ty_2Ehreal_2Ehreal ty_2Ehreal_2Ehreal)) \quad (12)$$

Definition 12 We define $c_2Erealax_2Ereal_ABS$ to be $\lambda V0r \in (ty_2Epair_2Eprod ty_2Ehreal_2Ehreal ty$

Definition 13 We define $c_2Erealax_2Ereal_neg$ to be $\lambda V0T1 \in ty_2Erealax_2Ereal.(ap c_2Erealax_2Ereal$

Let $c_2Erealax_2Etrealm_add : \iota$ be given. Assume the following.

$$c_2Erealax_2Etrealm_add \in (((ty_2Epair_2Eprod ty_2Ehreal_2Ehreal ty_2Ehreal_2Ehreal)(ty_2Epair_2Eprod ty_2Ehreal_2Ehreal ty_2Ehreal_2Ehreal)))(ty_2Epair_2Eprod ty_2Ehreal_2Ehreal ty_2Ehreal_2Ehreal)) \quad (13)$$

Definition 14 We define $c_2Erealax_2Ereal_add$ to be $\lambda V0T1 \in ty_2Erealax_2Ereal.\lambda V1T2 \in ty_2Erealax$

Definition 15 We define $c_2Ereal_2Ereal_sub$ to be $\lambda V0x \in ty_2Erealax_2Ereal.\lambda V1y \in ty_2Erealax_2E$

Assume the following.

$$True \quad (14)$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0x \in A_27a. ((V0x = V0x) \Leftrightarrow True)) \quad (15)$$

Assume the following.

$$(\forall V0x \in ty_2Erealax_2Ereal. ((ap (ap\ c_2Erealax_2Ereal_add (ap\ c_2Ereal_2Ereal_of_num\ c_2Enum_2E0))\ V0x) = V0x)) \quad (16)$$

Assume the following.

$$(\forall V0x \in ty_2Erealax_2Ereal. (\forall V1y \in ty_2Erealax_2Ereal. (\forall V2z \in ty_2Erealax_2Ereal. ((p (ap (ap\ c_2Ereal_2Ereal_lte (ap (ap\ c_2Erealax_2Ereal_add\ V0x)\ V2z)) (ap (ap\ c_2Erealax_2Ereal_add\ V1y)\ V2z))) \Leftrightarrow (p (ap (ap\ c_2Ereal_2Ereal_lte\ V0x)\ V1y)))))) \quad (17)$$

Assume the following.

$$(\forall V0x \in ty_2Erealax_2Ereal. (\forall V1y \in ty_2Erealax_2Ereal. ((ap (ap\ c_2Erealax_2Ereal_add (ap (ap\ c_2Ereal_2Ereal_sub\ V0x)\ V1y))\ V1y) = V0x))) \quad (18)$$

Theorem 1

$$(\forall V0x \in ty_2Erealax_2Ereal. (\forall V1y \in ty_2Erealax_2Ereal. ((p (ap (ap\ c_2Ereal_2Ereal_lte (ap\ c_2Ereal_2Ereal_of_num\ c_2Enum_2E0)) (ap (ap\ c_2Ereal_2Ereal_sub\ V0x)\ V1y))) \Leftrightarrow (p (ap (ap\ c_2Ereal_2Ereal_lte\ V1y)\ V0x))))))$$