

thm_2Ereal_2EREAL__SUP__SOMEPOS (TMd- fVmugtVxQ2hxcqBMJSafsMjCWbMjKuJK)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define $c_2Ebool_2E_2T$ to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 3 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p \Rightarrow q)$ of type ι .

Definition 4 We define $c_2Ebool_2E_21$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap (ap (c_2Emin_2E_3D (2^{A_27a}))$

Definition 5 We define $c_2Ebool_2E_5C_2E_2F$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t))$

Definition 6 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A.if (\exists x \in A.p (ap P x))$ then $(the (\lambda x.x \in A \wedge p x))$ of type $\iota \Rightarrow \iota$.

Definition 7 We define $c_2Ebool_2E_3F$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap V0P (ap (c_2Emin_2E_40 A_27a P))$

Let $c_2Enum_2EZERO_REP : \iota$ be given. Assume the following.

$$c_2Enum_2EZERO_REP \in \omega \tag{1}$$

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \tag{2}$$

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum^{\omega}) \tag{3}$$

Definition 8 We define c_2Enum_2E0 to be $(ap\ c_2Enum_2EABS_num\ c_2Enum_2EZERO_REP)$.

Let $ty_2Erealax_2Ereal : \iota$ be given. Assume the following.

$$nonempty\ ty_2Erealax_2Ereal \tag{4}$$

Let $c_2Ereal_2Ereal_of_num : \iota$ be given. Assume the following.

$$c_2Ereal_2Ereal_of_num \in (ty_2Erealax_2Ereal^{ty_2Enum_2Enum}) \tag{5}$$

Let $ty_2Ehreal_2Ehreal : \iota$ be given. Assume the following.

$$nonempty\ ty_2Ehreal_2Ehreal \quad (6)$$

Let $ty_2Epair_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty_2Epair_2Eprod\ A0\ A1) \quad (7)$$

Let $c_2Erealax_2Ereal_REP_CLASS : \iota$ be given. Assume the following.

$$c_2Erealax_2Ereal_REP_CLASS \in ((2^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)})^{ty_2Erealax_2Ereal_REP_CLASS}) \quad (8)$$

Definition 9 We define $c_2Erealax_2Ereal_REP$ to be $\lambda V0a \in ty_2Erealax_2Ereal.(ap\ (c_2Emin_2E_40\ (ty_2Erealax_2Ereal_REP_CLASS\ a)))$

Let $c_2Erealax_2Etreall_lt : \iota$ be given. Assume the following.

$$c_2Erealax_2Etreall_lt \in ((2^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)})^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)}) \quad (9)$$

Definition 10 We define $c_2Erealax_2Ereal_lt$ to be $\lambda V0T1 \in ty_2Erealax_2Ereal.\lambda V1T2 \in ty_2Erealax_2Ereal.(c_2Erealax_2Ereal_lt\ T1\ T2)$

Definition 11 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap\ (c_2Ebool_2E_21\ 2)\ (V0t1\ V1t2))))$

Definition 12 We define c_2Ebool_2EF to be $(ap\ (c_2Ebool_2E_21\ 2)\ (\lambda V0t \in 2.V0t))$.

Definition 13 We define $c_2Ebool_2E_7E$ to be $(\lambda V0t \in 2.(ap\ (ap\ c_2Emin_2E_3D_3D_3E\ V0t)\ c_2Ebool_2E_7E))$

Assume the following.

$$True \quad (10)$$

Assume the following.

$$(\forall V0t1 \in 2.(\forall V1t2 \in 2.(((p\ V0t1) \Rightarrow (p\ V1t2)) \Rightarrow (((p\ V1t2) \Rightarrow (p\ V0t1)) \Rightarrow ((p\ V0t1) \Leftrightarrow (p\ V1t2)))))) \quad (11)$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0t \in 2.((\forall V1x \in A_27a.(p\ V0t) \Leftrightarrow (p\ V0t)))) \quad (12)$$

Assume the following.

$$(\forall V0t \in 2.(((True \wedge (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \wedge True) \Leftrightarrow (p\ V0t)) \wedge (((False \wedge (p\ V0t)) \Leftrightarrow False) \wedge (((p\ V0t) \wedge False) \Leftrightarrow False) \wedge (((p\ V0t) \wedge (p\ V0t)) \Leftrightarrow (p\ V0t)))))) \quad (13)$$

Assume the following.

$$(\forall V0t \in 2.(((True \Rightarrow (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \Rightarrow True) \Leftrightarrow True) \wedge (((False \Rightarrow (p\ V0t)) \Leftrightarrow True) \wedge (((p\ V0t) \Rightarrow (p\ V0t)) \Leftrightarrow True) \wedge (((p\ V0t) \Rightarrow False) \Leftrightarrow (\neg (p\ V0t)))))) \quad (14)$$

Assume the following.

$$(\forall V0t \in 2.(((True \Leftrightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Leftrightarrow True) \Leftrightarrow (p V0t)) \wedge (((False \Leftrightarrow (p V0t)) \Leftrightarrow \neg(p V0t)) \wedge (((p V0t) \Leftrightarrow False) \Leftrightarrow \neg(p V0t)))))) \quad (15)$$

Assume the following.

$$(\forall V0x \in ty_2Erealx_2Ereal. (\forall V1y \in ty_2Erealx_2Ereal. ((V0x = V1y) \vee ((p (ap (ap c_2Erealx_2Ereal_lt V0x) V1y)) \vee (p (ap (ap c_2Erealx_2Ereal_lt V1y) V0x)))))) \quad (16)$$

Assume the following.

$$(\forall V0x \in ty_2Erealx_2Ereal. (\forall V1y \in ty_2Erealx_2Ereal. (\forall V2z \in ty_2Erealx_2Ereal. (((p (ap (ap c_2Erealx_2Ereal_lt V0x) V1y)) \wedge (p (ap (ap c_2Erealx_2Ereal_lt V1y) V2z))) \Rightarrow (p (ap (ap c_2Erealx_2Ereal_lt V0x) V2z)))))) \quad (17)$$

Assume the following.

$$(\forall V0P \in (2^{ty_2Erealx_2Ereal}). (((\forall V1x \in ty_2Erealx_2Ereal. ((p (ap V0P V1x)) \Rightarrow (p (ap (ap c_2Erealx_2Ereal_lt (ap c_2Ereal_2Ereal_of_num c_2Enum_2E0) V1x)))))) \wedge ((\exists V2x \in ty_2Erealx_2Ereal. (p (ap V0P V2x))) \wedge (\exists V3z \in ty_2Erealx_2Ereal. (\forall V4x \in ty_2Erealx_2Ereal. ((p (ap V0P V4x)) \Rightarrow (p (ap (ap c_2Erealx_2Ereal_lt V4x) V3z)))))) \Rightarrow (\exists V5s \in ty_2Erealx_2Ereal. (\forall V6y \in ty_2Erealx_2Ereal. ((\exists V7x \in ty_2Erealx_2Ereal. ((p (ap V0P V7x)) \wedge (p (ap (ap c_2Erealx_2Ereal_lt V6y) V7x)))) \Leftrightarrow (p (ap (ap c_2Erealx_2Ereal_lt V6y) V5s)))))) \quad (18)$$

Assume the following.

$$(\forall V0t \in 2. ((\neg(\neg(p V0t))) \Leftrightarrow (p V0t))) \quad (19)$$

Assume the following.

$$(\forall V0A \in 2. ((p V0A) \Rightarrow ((\neg(p V0A)) \Rightarrow False))) \quad (20)$$

Assume the following.

$$(\forall V0A \in 2. (((\neg(p V0A)) \Rightarrow False) \Rightarrow ((p V0A) \Rightarrow False))) \quad (21)$$

Assume the following.

$$(\forall V0p \in 2. (\forall V1q \in 2. ((\neg((p V0p) \Rightarrow (p V1q))) \Rightarrow (p V0p)))) \quad (22)$$

Assume the following.

$$(\forall V0p \in 2. (\forall V1q \in 2. ((\neg((p V0p) \Rightarrow (p V1q))) \Rightarrow (\neg(p V1q)))) \quad (23)$$

Theorem 1

$$\begin{aligned} & (\forall V0P \in (2^{ty_2Erealax_2Ereal}).((\exists V1x \in ty_2Erealax_2Ereal. \\ & ((p (ap V0P V1x)) \wedge (p (ap (ap c_2Erealax_2Ereal_lt (ap c_2Ereal_2Ereal_of_num \\ & c_2Enum_2E0)) V1x)))) \wedge (\exists V2z \in ty_2Erealax_2Ereal. (\forall V3x \in \\ & ty_2Erealax_2Ereal. ((p (ap V0P V3x)) \Rightarrow (p (ap (ap c_2Erealax_2Ereal_lt \\ & V3x) V2z)))))) \Rightarrow (\exists V4s \in ty_2Erealax_2Ereal. (\forall V5y \in \\ & ty_2Erealax_2Ereal. ((\exists V6x \in ty_2Erealax_2Ereal. ((p (\\ & ap V0P V6x)) \wedge (p (ap (ap c_2Erealax_2Ereal_lt V5y) V6x)))) \Leftrightarrow (p (\\ & ap (ap c_2Erealax_2Ereal_lt V5y) V4s)))))) \end{aligned}$$