

thm_2Ereal_2Ereal_lt

(TMFYSeKsPz3o9uetwEwp4MxPcJyC2Pzn4ax)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define c_2Ebool_2ET to be $(ap (ap (c_2Emin_2E_3D (2^2))) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x)$

Let $ty_2Ehreal_2Ehreal : \iota$ be given. Assume the following.

$$nonempty\ ty_2Ehreal_2Ehreal \tag{1}$$

Let $ty_2Epair_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty_2Epair_2Eprod\ A0\ A1) \tag{2}$$

Let $ty_2Erealax_2Ereal : \iota$ be given. Assume the following.

$$nonempty\ ty_2Erealax_2Ereal \tag{3}$$

Let $c_2Erealax_2Ereal_REP_CLASS : \iota$ be given. Assume the following.

$$c_2Erealax_2Ereal_REP_CLASS \in ((2^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)})\ ty_2Erealax) \tag{4}$$

Definition 3 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A.if (\exists x \in A.p (ap\ P\ x))$ then (the $(\lambda x.x \in A \wedge p$ of type $\iota \Rightarrow \iota$).

Definition 4 We define $c_2Ebool_2E_21$ to be $\lambda A.\lambda 27a : \iota.(\lambda V0P \in (2^{A-27a}).(ap (ap (c_2Emin_2E_3D (2^{A-27a})))$

Definition 5 We define $c_2Erealax_2Ereal_REP$ to be $\lambda V0a \in ty_2Erealax_2Ereal.(ap (c_2Emin_2E_40 (ty$

Let $c_2Erealax_2Etreallt : \iota$ be given. Assume the following.

$$c_2Erealax_2Etreallt \in ((2^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)})\ (ty_2Epair_2Eprod\ ty_2Ehreal)) \tag{5}$$

Definition 6 We define $c_2Erealax_2Ereal_lt$ to be $\lambda V0T1 \in ty_2Erealax_2Ereal.\lambda V1T2 \in ty_2Erealax_2Ereal$

Definition 7 We define c_2Ebool_2EF to be $(ap (c_2Ebool_2E_21\ 2) (\lambda V0t \in 2.V0t))$.

Definition 8 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow p Q)$ of type ι .

Definition 9 We define $c_2Ebool_2E_7E$ to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2EF$

Definition 10 We define $c_2Ereal_2Ereal_lte$ to be $\lambda V0x \in ty_2Erealax_2Ereal.\lambda V1y \in ty_2Erealax_2Er$

Definition 11 We define $c_2Ebool_2E_5C_2F$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21\ 2) (\lambda V2t \in$

Definition 12 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21\ 2) (\lambda V2t \in$

Assume the following.

$$True \tag{6}$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2.(((True \Leftrightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Leftrightarrow True) \Leftrightarrow \\ & (p V0t)) \wedge (((False \Leftrightarrow (p V0t)) \Leftrightarrow \neg(p V0t)) \wedge (((p V0t) \Leftrightarrow False) \Leftrightarrow \neg(\\ & p V0t)))))) \end{aligned} \tag{7}$$

Assume the following.

$$(\forall V0t \in 2.((\neg(\neg(p V0t))) \Leftrightarrow (p V0t))) \tag{8}$$

Assume the following.

$$(\forall V0A \in 2.((p V0A) \Rightarrow ((\neg(p V0A)) \Rightarrow False))) \tag{9}$$

Assume the following.

$$\begin{aligned} & (\forall V0A \in 2.(\forall V1B \in 2.(((\neg((\neg(p V0A)) \vee (p V1B))) \Rightarrow False) \Leftrightarrow \\ & (((p V0A) \Rightarrow False) \Rightarrow ((\neg(p V1B)) \Rightarrow False)))))) \end{aligned} \tag{10}$$

Assume the following.

$$\begin{aligned} & (\forall V0A \in 2.(\forall V1B \in 2.(((\neg((\neg(p V0A)) \vee (p V1B))) \Rightarrow False) \Leftrightarrow \\ & ((p V0A) \Rightarrow ((\neg(p V1B)) \Rightarrow False)))))) \end{aligned} \tag{11}$$

Assume the following.

$$(\forall V0A \in 2.(((\neg(p V0A)) \Rightarrow False) \Rightarrow (((p V0A) \Rightarrow False) \Rightarrow False))) \tag{12}$$

Assume the following.

$$\begin{aligned} & (\forall V0p \in 2.(\forall V1q \in 2.(\forall V2r \in 2.(((p V0p) \Leftrightarrow (\\ & (p V1q) \Leftrightarrow (p V2r))) \Leftrightarrow (((p V0p) \vee ((p V1q) \vee (p V2r))) \wedge (((p V0p) \vee ((\neg(\\ & p V2r)) \vee (\neg(p V1q)))) \wedge (((p V1q) \vee ((\neg(p V2r)) \vee (\neg(p V0p)))) \wedge ((p V2r) \vee \\ & ((\neg(p V1q)) \vee (\neg(p V0p)))))))))) \end{aligned} \tag{13}$$

Assume the following.

$$(\forall V0p \in 2. (\forall V1q \in 2. (((p V0p) \Leftrightarrow (\neg(p V1q))) \Leftrightarrow (((p V0p) \vee (p V1q)) \wedge ((\neg(p V1q)) \vee (\neg(p V0p))))))) \quad (14)$$

Assume the following.

$$(\forall V0p \in 2. (\forall V1q \in 2. ((\neg((p V0p) \Rightarrow (p V1q))) \Rightarrow (p V0p)))) \quad (15)$$

Assume the following.

$$(\forall V0p \in 2. (\forall V1q \in 2. ((\neg((p V0p) \Rightarrow (p V1q))) \Rightarrow (\neg(p V1q)))))) \quad (16)$$

Theorem 1

$$(\forall V0y \in ty_2Erealax_2Ereal. (\forall V1x \in ty_2Erealax_2Ereal. ((p (ap (ap c_2Erealax_2Ereal_lt V1x) V0y)) \Leftrightarrow (\neg(p (ap (ap c_2Ereal_2Ereal_lte V0y) V1x))))))$$