

thm\_2Ereal\_topo<sub>logy</sub>\_2EADD<sub>2</sub>SUB2 (TM-LYNWY9CEEeEfvwze5zSvp1VUhiBmnGDhL)

October 26, 2020

Let  $ty\_2Enum\_2Enum : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Enum\_2Enum \quad (1)$$

Let  $c\_2Earithmetic\_2E\_2B : \iota$  be given. Assume the following.

$$c\_2Earithmetic\_2E\_2B \in ((ty\_2Enum\_2Enum^{ty\_2Enum\_2Enum})^{ty\_2Enum\_2Enum}) \quad (2)$$

Let  $c\_2Earithmetic\_2E\_2D : \iota$  be given. Assume the following.

$$c\_2Earithmetic\_2E\_2D \in ((ty\_2Enum\_2Enum^{ty\_2Enum\_2Enum})^{ty\_2Enum\_2Enum}) \quad (3)$$

**Definition 1** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A. \lambda x \in A. \lambda y \in A. inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define  $c\_2Ebool\_2ET$  to be  $(ap (ap (c\_2Emin\_2E\_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x)))$

**Definition 3** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A\_27a : \iota. (\lambda V0P \in (2^{A\_27a}). (ap (ap (c\_2Emin\_2E\_3D (2^{A\_27a})) (\lambda V1x \in 2.V1x)))$

Assume the following.

$$(\forall V0m \in ty\_2Enum\_2Enum. (\forall V1n \in ty\_2Enum\_2Enum. (ap (ap c\_2Earithmetic\_2E\_2B V0m) V1n) = (ap (ap c\_2Earithmetic\_2E\_2B V1n) V0m))) \quad (4)$$

Assume the following.

$$(\forall V0a \in ty\_2Enum\_2Enum. (\forall V1c \in ty\_2Enum\_2Enum. (ap (ap c\_2Earithmetic\_2E\_2D (ap (ap c\_2Earithmetic\_2E\_2B V0a) V1c)) V1c) = V0a))) \quad (5)$$

**Theorem 1**

$$(\forall V0m \in ty\_2Enum\_2Enum. (\forall V1n \in ty\_2Enum\_2Enum. (ap (ap c\_2Earithmetic\_2E\_2D (ap (ap c\_2Earithmetic\_2E\_2B V0m) V1n)) V0m) = V1n)))$$