

thm\_2Ereal\_topology\_2EADD\_SUBR2 (TMMn-FKKAHdcMmKR7MpAHUHsMSRLktxceEG6)

October 26, 2020

Let  $ty\_2Enum\_2Enum : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Enum\_2Enum \quad (1)$$

Let  $c\_2Earithmetic\_2E\_2B : \iota$  be given. Assume the following.

$$c\_2Earithmetic\_2E\_2B \in ((ty\_2Enum\_2Enum^{ty\_2Enum\_2Enum})^{ty\_2Enum\_2Enum}) \quad (2)$$

**Definition 1** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A. \lambda x \in A. \lambda y \in A. inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define  $c\_2Ebool\_2ET$  to be  $(ap (ap (c\_2Emin\_2E\_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

**Definition 3** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A\_27a : \iota. (\lambda V0P \in (2^{A\_27a}).(ap (ap (c\_2Emin\_2E\_3D (2^{A\_27a})) (\lambda V1P \in 2.V1P)) (\lambda V2P \in 2.V2P)))$

**Definition 4** We define  $c\_2Ebool\_2EF$  to be  $(ap (c\_2Ebool\_2E\_21 2) (\lambda V0t \in 2.V0t))$ .

**Definition 5** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2. \lambda Q \in 2. inj\_o (p \Rightarrow p \ Q)$  of type  $\iota$ .

**Definition 6** We define  $c\_2Ebool\_2E\_7E$  to be  $(\lambda V0t \in 2. (ap (ap c\_2Emin\_2E\_3D\_3D\_3E V0t) c\_2Ebool\_2EF))$

**Definition 7** We define  $c\_2Ebool\_2E\_2F\_5C$  to be  $(\lambda V0t1 \in 2. (\lambda V1t2 \in 2. (ap (c\_2Ebool\_2E\_21 2) (\lambda V2t \in 2. inj\_o (V0t1 = V1t2))))$

Let  $c\_2Enum\_2EREP\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EREP\_num \in (\omega^{ty\_2Enum\_2Enum}) \quad (3)$$

Let  $c\_2Enum\_2ESUC\_REP : \iota$  be given. Assume the following.

$$c\_2Enum\_2ESUC\_REP \in (\omega^{\omega}) \quad (4)$$

Let  $c\_2Enum\_2EABS\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EABS\_num \in (ty\_2Enum\_2Enum^{\omega}) \quad (5)$$

**Definition 8** We define  $c\_2Enum\_2ESUC$  to be  $\lambda V0m \in ty\_2Enum\_2Enum.(ap c\_2Enum\_2EABS\_num ($

**Definition 9** We define  $c\_2Emin\_2E\_40$  to be  $\lambda A.\lambda P \in 2^A.\text{if } (\exists x \in A.p (ap P x)) \text{ then } (\lambda x.x \in A \wedge p$  of type  $\iota \Rightarrow \iota$ .

**Definition 10** We define  $c\_2Ebool\_2E\_3F$  to be  $\lambda A\_27a : \iota.(\lambda V0P \in (2^{A\_27a}).(ap V0P (ap (c\_2Emin\_2E\_40$

**Definition 11** We define  $c\_2Eprim\_rec\_2E\_3C$  to be  $\lambda V0m \in ty\_2Enum\_2Enum.\lambda V1n \in ty\_2Enum\_2Enum.$

**Definition 12** We define  $c\_2Ebool\_2E\_5C\_2F$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_2Ebool\_2E\_21 2) (\lambda V2t \in$

**Definition 13** We define  $c\_2Earithmetic\_2E\_3C\_3D$  to be  $\lambda V0m \in ty\_2Enum\_2Enum.\lambda V1n \in ty\_2Enum\_2Enum.$

Let  $c\_2Enum\_2EZERO\_REP : \iota$  be given. Assume the following.

$$c\_2Enum\_2EZERO\_REP \in \omega \quad (6)$$

**Definition 14** We define  $c\_2Enum\_2E0$  to be  $(ap c\_2Enum\_2EABS\_num c\_2Enum\_2EZERO\_REP).$

Let  $c\_2Earithmetic\_2E\_2D : \iota$  be given. Assume the following.

$$c\_2Earithmetic\_2E\_2D \in ((ty\_2Enum\_2Enum^{ty\_2Enum\_2Enum})^{ty\_2Enum\_2Enum}) \quad (7)$$

Assume the following.

$$\begin{aligned} & (\forall V0m \in ty\_2Enum\_2Enum.(\forall V1n \in ty\_2Enum\_2Enum. \\ & p (ap (ap c\_2Earithmetic\_2E\_3C\_3D V0m) (ap (ap c\_2Earithmetic\_2E\_2B \\ & V0m) V1n)))) \end{aligned} \quad (8)$$

Assume the following.

$$\begin{aligned} & (\forall V0m \in ty\_2Enum\_2Enum.(\forall V1n \in ty\_2Enum\_2Enum. \\ & ((ap (ap c\_2Earithmetic\_2E\_2D V0m) V1n) = c\_2Enum\_2E0) \Leftrightarrow (p (ap ( \\ & ap c\_2Earithmetic\_2E\_3C\_3D V0m) V1n)))) \end{aligned} \quad (9)$$

Assume the following.

$$True \quad (10)$$

Assume the following.

$$\forall A\_27a.\text{nonempty } A\_27a \Rightarrow (\forall V0t \in 2.((\forall V1x \in \\ A\_27a.(p V0t)) \Leftrightarrow (p V0t))) \quad (11)$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2.(((True \Leftrightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Leftrightarrow True) \Leftrightarrow \\ & (p V0t)) \wedge (((False \Leftrightarrow (p V0t)) \Leftrightarrow (\neg(p V0t))) \wedge (((p V0t) \Leftrightarrow False) \Leftrightarrow (\neg(p \\ & V0t))))))) \end{aligned} \quad (12)$$

### Theorem 1

$$\begin{aligned} & (\forall V0m \in ty\_2Enum\_2Enum.(\forall V1n \in ty\_2Enum\_2Enum. \\ & (ap (ap c\_2Earithmetic\_2E\_2D V0m) (ap (ap c\_2Earithmetic\_2E\_2B \\ & V0m) V1n)) = c\_2Enum\_2E0))) \end{aligned}$$