

thm_2Ereal__topology_2EALWAYS__EVENTUALLY (TMM4KbEWcBqEfHDje2oZeCVnMR4S546q9C1)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define $c_2Ebool_2E_ET$ to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Let $ty_2Ereal_topology_2E_enet : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty A0 \Rightarrow nonempty (ty_2Ereal_topology_2E_enet A0) \quad (1)$$

Let $c_2Ereal_topology_2E_enetord : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow c_2Ereal_topology_2E_enetord A_27a \in ((2^{A_27a})^{A_27a})^{(ty_2Ereal_topology_2E_enet A_27a)} \quad (2)$$

Definition 3 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A.if (\exists x \in A.p (ap P x))$ then (the $(\lambda x.x \in A \wedge p x)$) of type $\iota \Rightarrow \iota$.

Definition 4 We define $c_2Ebool_2E_3F$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap V0P (ap (c_2Emin_2E_40 A_27a P))))$

Definition 5 We define $c_2Ebool_2E_21$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap (ap (c_2Emin_2E_3D (2^{A_27a} P)) P)))$

Definition 6 We define $c_2Ebool_2E_EF$ to be $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$.

Definition 7 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow p Q)$ of type ι .

Definition 8 We define $c_2Ebool_2E_7E$ to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2E_EF))$

Definition 9 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t))))$

Definition 10 We define $c_2Ebool_2E_5C_2F$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t))))$

Definition 11 We define $c_2Ereal_topology_2E_trivial_limit$ to be $\lambda A_27a : \iota.\lambda V0net \in (ty_2Ereal_topology_2E_enet A_27a)$

Definition 12 We define $c_2Ereal_topology_2Eeventually$ to be $\lambda A_27a : \iota.\lambda V0p \in (2^{A_27a}).\lambda V1net \in (ty_2$

Assume the following.

$$True \tag{3}$$

Assume the following.

$$(\forall V0t1 \in 2.(\forall V1t2 \in 2.(((p V0t1) \Rightarrow (p V1t2)) \Rightarrow (((p V1t2) \Rightarrow (p V0t1)) \Rightarrow ((p V0t1) \Leftrightarrow (p V1t2)))))) \tag{4}$$

Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow (\forall V0t \in 2.((\forall V1x \in A_27a.(p V0t)) \Leftrightarrow (p V0t))) \tag{5}$$

Assume the following.

$$(\forall V0t \in 2.(((p V0t) \Rightarrow False) \Rightarrow (\neg(p V0t)))) \tag{6}$$

Assume the following.

$$(\forall V0t \in 2.((\neg(p V0t)) \Rightarrow ((p V0t) \Rightarrow False))) \tag{7}$$

Assume the following.

$$(\forall V0t \in 2.(((True \wedge (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \wedge True) \Leftrightarrow (p V0t)) \wedge (((False \wedge (p V0t)) \Leftrightarrow False) \wedge (((p V0t) \wedge False) \Leftrightarrow False) \wedge (((p V0t) \wedge (p V0t)) \Leftrightarrow (p V0t)))))) \tag{8}$$

Assume the following.

$$(\forall V0t \in 2.(((True \Rightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Rightarrow True) \Leftrightarrow True) \wedge (((False \Rightarrow (p V0t)) \Leftrightarrow True) \wedge (((p V0t) \Rightarrow (p V0t)) \Leftrightarrow True) \wedge ((p V0t) \Rightarrow False) \Leftrightarrow (\neg(p V0t)))))) \tag{9}$$

Assume the following.

$$((\forall V0t \in 2.((\neg(\neg(p V0t))) \Leftrightarrow (p V0t))) \wedge (((\neg True) \Leftrightarrow False) \wedge ((\neg False) \Leftrightarrow True))) \tag{10}$$

Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow (\forall V0x \in A_27a.(\forall V1y \in A_27a.((V0x = V1y) \Leftrightarrow (V1y = V0x)))) \tag{11}$$

Assume the following.

$$(\forall V0t \in 2.(((True \Leftrightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Leftrightarrow True) \Leftrightarrow (p V0t)) \wedge (((False \Leftrightarrow (p V0t)) \Leftrightarrow (\neg(p V0t))) \wedge (((p V0t) \Leftrightarrow False) \Leftrightarrow (\neg(p V0t)))))) \tag{12}$$

Assume the following.

$$\forall A.27a.nonempty A.27a \Rightarrow (\forall V0P \in (2^{A.27a}).((\neg(\forall V1x \in A.27a.(p (ap V0P V1x)))) \Leftrightarrow (\exists V2x \in A.27a.(\neg(p (ap V0P V2x)))))) \quad (13)$$

Assume the following.

$$\forall A.27a.nonempty A.27a \Rightarrow (\forall V0P \in (2^{A.27a}).((\neg(\exists V1x \in A.27a.(p (ap V0P V1x)))) \Leftrightarrow (\forall V2x \in A.27a.(\neg(p (ap V0P V2x)))))) \quad (14)$$

Assume the following.

$$\forall A.27a.nonempty A.27a \Rightarrow (\forall V0P \in (2^{A.27a}).(\forall V1Q \in (2^{A.27a}).((\forall V2x \in A.27a.((p (ap V0P V2x)) \wedge (p (ap V1Q V2x)))) \Leftrightarrow ((\forall V3x \in A.27a.(p (ap V0P V3x)) \wedge (\forall V4x \in A.27a.(p (ap V1Q V4x)))))))) \quad (15)$$

Assume the following.

$$\forall A.27a.nonempty A.27a \Rightarrow (\forall V0P \in (2^{A.27a}).(\forall V1Q \in (2^{A.27a}).((\exists V2x \in A.27a.((p (ap V0P V2x)) \vee (p (ap V1Q V2x)))) \Leftrightarrow ((\exists V3x \in A.27a.(p (ap V0P V3x)) \vee (\exists V4x \in A.27a.(p (ap V1Q V4x)))))))) \quad (16)$$

Assume the following.

$$\forall A.27a.nonempty A.27a \Rightarrow (\forall V0P \in 2.(\forall V1Q \in (2^{A.27a}).((p V0P) \vee (\exists V2x \in A.27a.(p (ap V1Q V2x)))) \Leftrightarrow (\exists V3x \in A.27a.((p V0P) \vee (p (ap V1Q V3x)))))) \quad (17)$$

Assume the following.

$$\forall A.27a.nonempty A.27a \Rightarrow (\forall V0P \in (2^{A.27a}).(\forall V1Q \in 2.((\exists V2x \in A.27a.((p (ap V0P V2x)) \wedge (p V1Q))) \Leftrightarrow ((\exists V3x \in A.27a.(p (ap V0P V3x)) \wedge (p V1Q)))))) \quad (18)$$

Assume the following.

$$\forall A.27a.nonempty A.27a \Rightarrow (\forall V0P \in 2.(\forall V1Q \in (2^{A.27a}).((\exists V2x \in A.27a.((p V0P) \wedge (p (ap V1Q V2x)))) \Leftrightarrow ((p V0P) \wedge (\exists V3x \in A.27a.(p (ap V1Q V3x)))))) \quad (19)$$

Assume the following.

$$(\forall V0A \in 2.(\forall V1B \in 2.(((\neg((p V0A) \wedge (p V1B))) \Leftrightarrow ((\neg(p V0A)) \vee (\neg(p V1B)))))) \wedge ((\neg((p V0A) \vee (p V1B))) \Leftrightarrow ((\neg(p V0A)) \wedge (\neg(p V1B)))))) \quad (20)$$

Assume the following.

$$\forall A.27a.nonempty A.27a \Rightarrow \forall A.27b.nonempty A.27b \Rightarrow (\forall V0P \in ((2^{A.27b})^{A.27a}).((\forall V1x \in A.27a.(\exists V2y \in A.27b.(p (ap (ap V0P V1x) V2y)))) \Leftrightarrow (\exists V3f \in (A.27b^{A.27a}).(\forall V4x \in A.27a.(p (ap (ap V0P V4x) (ap V3f V4x))))))) \quad (21)$$

Assume the following.

$$(\forall V0t \in 2.((\neg(\neg(p V0t))) \Leftrightarrow (p V0t))) \quad (22)$$

Assume the following.

$$(\forall V0A \in 2.((p V0A) \Rightarrow ((\neg(p V0A)) \Rightarrow \text{False}))) \quad (23)$$

Assume the following.

$$(\forall V0A \in 2.(\forall V1B \in 2.(((\neg((\neg(p V0A) \vee (p V1B))) \Rightarrow \text{False}) \Leftrightarrow ((p V0A) \Rightarrow \text{False}) \Rightarrow ((\neg(p V1B)) \Rightarrow \text{False})))))) \quad (24)$$

Assume the following.

$$(\forall V0A \in 2.(\forall V1B \in 2.(((\neg(\neg(p V0A) \vee (p V1B))) \Rightarrow \text{False}) \Leftrightarrow ((p V0A) \Rightarrow ((\neg(p V1B)) \Rightarrow \text{False})))))) \quad (25)$$

Assume the following.

$$(\forall V0A \in 2.(((\neg(p V0A)) \Rightarrow \text{False}) \Rightarrow (((p V0A) \Rightarrow \text{False}) \Rightarrow \text{False}))) \quad (26)$$

Assume the following.

$$(\forall V0p \in 2.(\forall V1q \in 2.(\forall V2r \in 2.(((p V0p) \Leftrightarrow (p V1q) \Leftrightarrow (p V2r))) \Leftrightarrow (((p V0p) \vee ((p V1q) \vee (p V2r))) \wedge (((p V0p) \vee ((\neg(p V2r)) \vee (\neg(p V1q)))) \wedge (((p V1q) \vee ((\neg(p V2r)) \vee (\neg(p V0p)))) \wedge ((p V2r) \vee ((\neg(p V1q)) \vee (\neg(p V0p)))))))))) \quad (27)$$

Assume the following.

$$(\forall V0p \in 2.(\forall V1q \in 2.(\forall V2r \in 2.(((p V0p) \Leftrightarrow (p V1q) \wedge (p V2r))) \Leftrightarrow (((p V0p) \vee ((\neg(p V1q)) \vee (\neg(p V2r)))) \wedge (((p V1q) \vee (\neg(p V0p))) \wedge ((p V2r) \vee (\neg(p V0p)))))))) \quad (28)$$

Assume the following.

$$(\forall V0p \in 2.(\forall V1q \in 2.(\forall V2r \in 2.(((p V0p) \Leftrightarrow (p V1q) \vee (p V2r))) \Leftrightarrow (((p V0p) \vee (\neg(p V1q))) \wedge (((p V0p) \vee (\neg(p V2r))) \wedge ((p V1q) \vee ((p V2r) \vee (\neg(p V0p)))))))) \quad (29)$$

Assume the following.

$$(\forall V0p \in 2.(\forall V1q \in 2.(\forall V2r \in 2.(((p V0p) \Leftrightarrow (p V1q) \Rightarrow (p V2r))) \Leftrightarrow (((p V0p) \vee (p V1q)) \wedge (((p V0p) \vee (\neg(p V2r))) \wedge ((\neg(p V1q)) \vee ((p V2r) \vee (\neg(p V0p)))))))) \quad (30)$$

Assume the following.

$$(\forall V0p \in 2.(\forall V1q \in 2.(((p V0p) \Leftrightarrow (\neg(p V1q))) \Leftrightarrow (((p V0p) \vee (p V1q)) \wedge ((\neg(p V1q)) \vee (\neg(p V0p)))))) \quad (31)$$

Theorem 1

$$\forall A_{27a}. \text{nonempty } A_{27a} \Rightarrow (\forall V_0 p \in (2^{A_{27a}}). (\forall V_1 \text{net} \in \\ (\text{ty_2Ereal_topology_2Enet } A_{27a}). ((\forall V_2 x \in A_{27a}. (p (\\ \text{ap } V_0 p \ V_2 x)))) \Rightarrow (p (\text{ap } (\text{ap } (\text{c_2Ereal_topology_2Eventually } A_{27a}) \\ V_0 p) \ V_1 \text{net}))))))$$